



Check List of Diptera (Insecta) of the Czech and Slovak Republics



KAROLINUM - CHARLES UNIVERSITY PRESS



Check List of Diptera (Insecta) of the Czech and Slovak Republics



Milan Chvála
editor

KAROLINUM
Charles University Press
Prague 1997

SCIENTIFIC ADVISERS
doc. RNDr. Zdeněk Laštůvka, CSc.
RNDr. Václav Skuhrový, CSc.

INTRODUCTION

Lists of insect faunas have a long tradition and are of outstanding importance as a basis for assessing the biodiversity of a particular geographic area. These lists of Diptera, or check lists as they are generally called, have a long history in Europe and provide a considerable stimulus for further dipterological and entomological studies. Even this year, just as a new revised edition of the Check List of Czech and Slovak flies is being made available, similar national lists are scheduled for publication in Switzerland, Germany and Great Britain.

Such lists are now essential tools for disseminating locally all the taxonomic and nomenclatural changes that appeared in the long-awaited Palaearctic Catalogue of Diptera, the thirteenth and final volume of which was published in 1994. Furthermore, since the first two volumes of the Catalogue were published as long ago as 1984 and the cut-off date for the inclusion described taxa for the whole series was 1982, some parts of the Catalogue are already out-of-date taxonomically and even nomenclaturally and are in urgent need of revision. Check Lists are therefore an important means of showing not only the national biodiversity but also the most recent corrections and additions in nomenclature, taxonomy and species composition.

Interest in a Check List of Diptera in the former Czechoslovakia dates from 1969, when the first meeting of Czech and Slovak dipterists, the so-called "Seminar of Czechoslovak dipterists", took place in Nitra, Slovakia. Since that time, regular dipterological seminars have been held every two years at different places, and the problem of the list of Diptera is one of the permanent tasks of these meetings. As early as 1971, a draft of the first working List was distributed to specialists for comment, but the final version ready for publication was not available until after the 8th seminar held in České Budějovice in 1984. The first Check List of Czechoslovak Diptera was published by the Prague National Museum in 1987 (see JEŽEK 1987). It included 6498 species in 104 families, and was compiled by 52 authors.

Since then, Czechoslovakia has split into two independent states, the Czech and Slovak Republics, but the arrangement of the present Check List has not been changed. It retains the format of the previous edition, including the division into the historical regions (from west to east): Bohemia (B), Moravia (M) and Slovakia (S). Species are always arranged alphabetically within genera or subgenera, whilst the genera may be listed in taxonomic order, showing the natural phylogenetic system, or in alphabetical order. There is, however, one substantial difference in the present Check List: authors were strongly urged to include only verified and authentic data; all unverified historical records or doubtful species have been excluded from this new, revised edition, and the rather common questionmarks of the 1987 edition have been completely omitted. The authors of the individual families were also asked to minimize the number of synonyms and to include only those that are really necessary for a correct understanding of the listed taxa. If names are given that are at variance with the Palaearctic Catalogue, the relevant taxonomic or nomenclatural changes will be found discussed in papers published by the authors of the families in question. This also applies to new or different faunistic data which are at variance with the original Check List of 1987. All the relevant literary records and other data published by Czech and Slovak authors are included in the bibliography of Czech and Slovak dipterological literature for the years 1986 to 1995 (see VAŇHARA & ROZKOŠNÝ 1997). All the papers published in 1996 and 1997, or papers submitted for publication, which contain data included in the present List, are included at the end in the References.

The present Check List of Diptera is a collaborative publication prepared by 43 authors, covering 7,640 species in 115 families. It must be obvious that the harmonisation of such a large number of manuscripts is a far from simple task. As a first step, several "supervisors" were asked to oversee larger units: L. JEDLIČKA and I. ORSZÁGH (Bratislava) were responsible for the Diptera Nematocera, M. CHVÁLA (Praha) for the Brachycera Orthorrhapha, J. HOLINKA (Pardubice) for the Cyclorrhapha Aschiza, J. ROHÁČEK (Opava) for the Schizophora Acalyptratae, and R. ROZKOŠNÝ (Brno) for the Schizophora Calyptratae. The classification of Diptera used in this List is based in general on that of the Palaearctic Catalogue, whilst the higher categories are mostly in accordance with the "New List of British and Irish Flies" organized by P. J. CHANDLER in Great Britain. If a different classification is accepted, it is based on specialist input in taxonomy and phylogeny by the individual authors, and further details may be found in the references.

The preparation of this new edition of the Check List of Czech and Slovak Diptera has been financially supported by Grant No. 206/96/0997 of the Grant Agency of the Czech Republic, by Grant No. 95/5195/069 of the Slovak Grant Agency, and by further additional grants supporting studies of individual families. The authors are also grateful to the authorities of the Charles University, Prague, for making possible the publication of this book, as well as to the two referees, Z. LAŠTŮVKA (Brno) and V. SKUHRAVÝ (Praha), for their careful reading of the manuscript and many useful suggestions. Very valuable editorial work has also been done by R. ROZKOŠNÝ (Brno). However, the fundamental and most meritorious work on the Check List has been carried out by J. VAŇHARA (Brno), whose editorial computer work on the many different manuscripts was crucial for printing this technically difficult publication.

Praha, 25th June 1997

M. CHVALA

CONTENTS

Higher Classification	9
TRICHOCERIDAE	(Jaroslav Starý & Jaroslav Martinovský) 11
LIMONIIDAE	(Jaroslav Starý) 11
PEDICIIDAE	(Jaroslav Starý) 15
CYLINDROTOMIDAE	(Jaroslav Starý) 15
TIPULIDAE	(Jaroslav Martinovský) 15
BLEPHARICERIDAE	(Jaroslav Starý & Ján Gulička) 17
AXYMYIIDAE	(Jaroslav Martinovský) 17
BIBIONIDAE	(Pavel Pecina) 17
PLECIIDAE	(Jaroslav Martinovský) 17
HESPERINIDAE	(Jaroslav Martinovský) 17
BOLITOPHILIDAE	(Vladimír Košel & Jaroslav Martinovský) 18
DIADOCIDIIDAE	(Vladimír Košel & Jaroslav Martinovský) 18
DITOMYIIDAE	(Vladimír Košel & Jaroslav Martinovský) 18
KEROPLATIDAE	(Vladimír Košel, Peter Chandler & Jaroslav Martinovský) 18
MACRO CERIDAE	(Vladimír Košel & Jaroslav Martinovský) 18
MYCETOPHILIDAE	(Vladimír Košel, Peter Chandler & Jaroslav Martinovský) 19
SCIARIDAE	(Hans-Georg Rudzinski & Vladimír Košel) 23
CECIDOMYIIDAE	(Marcela Skuhrová) 25
PSYCHODIDAE	(Jan Ježek) 30
ANISOPODIDAE	(Ivo Kovář) 32
MYCETOBIIDAE	(Jaroslav Martinovský) 32
SCATOPSIDAE	(Jaroslav Martinovský) 32
SYNNEURIDAE	(Jaroslav Martinovský) 33
CANTHYLOSCELIDAE	(Jaroslav Martinovský) 33
PTYCHOPTERIDAE	(Jaroslav Starý) 33
DIXIDAE	(Jan Knoz) 33
CHAOBORIDAE	(Ladislav Jedlička & Ivan Országh) 33
CULICIDAE	(Jan Minář & Jozef Halgoš) 34
CHIRONOMIDAE	(Peter Bitušik & Bohumil Losos) 34
CERATOPOGONIDAE	(Ivan Országh, Jan Knoz & Josef Chalupský) 39
SIMULIIDAE	(Jan Knoz, Ladislav Jedlička) 42
THAUMALEIDAE	(Jaroslav Martinovský) 42
XYLOPHAGIDAE	(Rudolf Rozkošný) 43
COENOMYIIDAE	(Rudolf Rozkošný) 43
ATHERICIDAE	(Rudolf Rozkošný & Karel Spitzer) 43
RHAGIONIDAE	(Rudolf Rozkošný & Karel Spitzer) 43
TABANIDAE	(Milan Chvála) 43
XYLOMYIDAE	(Rudolf Rozkošný) 44
STRATIOMYIDAE	(Rudolf Rozkošný) 44
ACROCERIDAE	(Milan Chvála) 45
MYTHICOMYIIDAE	(Jaroslav Bosák) 45
PHTHIRIIDAE	(Jaroslav Bosák) 45
BOMBYLIIDAE	(Jaroslav Bosák) 45
HILARIMORPHIDAE	(Rudolf Rozkošný) 46
THEREVIDAE	(Karel Spitzer) 46
SCENOPINIDAE	(Jaroslav Bosák) 46
ASILIDAE	(Jaroslav Bosák) 46
EMPIDIDAE	(Milan Chvála) 48
HYBOTIDAE	(Milan Chvála) 51
DOLICHOPODIDAE	(Jiří Olejníček) 54
ATELESTIDAE	(Milan Chvála) 57
OPETIIDAE	(Jaromír Vaňhara) 57
PLATYPEZIDAE	(Jaromír Vaňhara) 57
PHORIDAE	(Bohuslav Mocek) 58
LONCHOPTERIDAE	(Miroslav Barták) 60
SYRPHIDAE	(Jiří Holinka & Libor Mazánek) 60

CONTENTS

Higher Classification	9
TRICHOCERIDAE (Jaroslav Starý & Jaroslav Martinovský)	11
LIMONIIDAE (Jaroslav Starý)	11
PEDICIIDAE (Jaroslav Starý)	15
CYLINDROTOMIDAE (Jaroslav Starý)	15
TIPULIDAE (Jaroslav Martinovský)	15
BLEPHARICERIDAE (Jaroslav Starý & Ján Gulička)	17
AXYMYIIDAE (Jaroslav Martinovský)	17
BIBIONIDAE (Pavel Pecina)	17
PLECIIDAE (Jaroslav Martinovský)	17
HESPERINIDAE (Jaroslav Martinovský)	17
BOLITOPHILIDAE (Vladimír Košel & Jaroslav Martinovský)	18
DIADOCIDIIDAE (Vladimír Košel & Jaroslav Martinovský)	18
DITOMYIIDAE (Vladimír Košel & Jaroslav Martinovský)	18
KEROPLATIDAE (Vladimír Košel, Peter Chandler & Jaroslav Martinovský)	18
MACROCERIDAE (Vladimír Košel & Jaroslav Martinovský)	18
MYCETOPHILIDAE (Vladimír Košel, Peter Chandler & Jaroslav Martinovský)	19
SCIARIDAE (Hans-Georg Rudzinski & Vladimír Košel)	23
CECIDOMYIIDAE (Marcela Skuhrová)	25
PSYCHODIDAE (Jan Ježek)	30
ANISOPODIDAE (Ivo Kovář)	32
MYCETOBIIIDAE (Jaroslav Martinovský)	32
SCATOPSIDAE (Jaroslav Martinovský)	32
SYNNEURIDAE (Jaroslav Martinovský)	33
CANTHYLOSCELIDAE (Jaroslav Martinovský)	33
PTYCHOPTERIDAE (Jaroslav Starý)	33
DIXIDAE (Jan Knoz)	33
CHAOBORIDAE (Ladislav Jedlička & Ivan Országh)	33
CULICIDAE (Jan Míňák & Jozef Halgoš)	34
CHIRONOMIDAE (Peter Bitušík & Bohumil Losos)	34
CERATOPOGONIDAE (Ivan Országh, Jan Knoz & Josef Chalupský)	39
SIMULIIDAE (Jan Knoz, Ladislav Jedlička)	42
THAUMALEIDAE (Jaroslav Martinovský)	42
XYLOPHAGIDAE (Rudolf Rozkošný)	43
COENOMYIIDAE (Rudolf Rozkošný)	43
ATHERICIDAE (Rudolf Rozkošný & Karel Spitzer)	43
RHAGIONIDAE (Rudolf Rozkošný & Karel Spitzer)	43
TABANIDAE (Milan Chvála)	43
XYLOMYIDAE (Rudolf Rozkošný)	44
STRATIOMYIDAE (Rudolf Rozkošný)	44
ACROCERIDAE (Milan Chvála)	45
MYTHICOMYIIDAE (Jaroslav Bosák)	45
PHTHIRIIDAE (Jaroslav Bosák)	45
BOMBYLIIDAE (Jaroslav Bosák)	45
HILARIMORPHIDAE (Rudolf Rozkošný)	46
THEREVIDAE (Karel Spitzer)	46
SCENOPINIDAE (Jaroslav Bosák)	46
ASILIDAE (Jaroslav Bosák)	46
EMPIDIDAE (Milan Chvála)	48
HYBOTIDAE (Milan Chvála)	51
DOLICHOPODIDAE (Jiří Olejníček)	54
ATELESTIDAE (Milan Chvála)	57
OPETIIDAE (Jaromír Vaňhara)	57
PLATYPEZIDAE (Jaromír Vaňhara)	57
PHORIDAE (Bohuslav Mocek)	58
LONCHOPTERIDAE (Miroslav Barták)	60
SYRPHIDAE (Jiří Holinka & Libor Mazánek)	60

PIPUNCULIDAE	(Pavel Lauterer)	66
PSEUDOPOMYZIDAE	(Jindřich Roháček)	67
MICROPEZIDAE	(Jindřich Roháček)	67
TANYPEZIDAE	(Jindřich Roháček)	67
STRONGYLOPHTHALMYIIDAE	(Jindřich Roháček)	67
MEGAMERINIDAE	(Jindřich Roháček)	67
PSILIDAE	(Vladislav Martinek)	67
CONOPIDAE	(Milan Chvála)	68
LONCHAEIDAE	(Jan Máca)	68
PALLOPTERIDAE	(Vladislav Martinek)	69
PIOPHILIDAE	(Jan Zuska & Miroslav Barták)	69
OTITIDAE	(Vladislav Martinek)	70
ULIDIIDAE	(Jan Zuska)	70
PLATYSTOMATIDAE	(Vladislav Martinek)	70
TEPHRITIDAE	(Judita Kinkorová)	70
LAUXANIIDAE	(Vladislav Martinek)	72
CHAMAEMYIIDAE	(Vlastimil Lapáček)	73
CREMIFANIIDAE	(Jan Máca)	73
DRYOMYZIDAE	(Vladislav Martinek)	73
PHAEOMYIIDAE	(Rudolf Rozkošný)	73
SCIOMYZIDAE	(Rudolf Rozkošný)	73
SEPSIDAE	(Jan Zuska & Miroslav Barták)	74
CLUSIIDAE	(Jindřich Roháček)	75
ACARTOPHTHALMIDAE	(Jindřich Roháček)	75
ODINIIDAE	(Jan Máca)	75
AGROMYZIDAE	(Miloslav Vála & Miloš Černý)	75
OPOMYZIDAE	(Vladislav Martinek)	78
ANTHOMYZIDAE	(Jindřich Roháček)	78
AULACIGASTRIDAE	(Jan Máca)	79
PERISCCELIDAE	(Jan Máca & Jindřich Roháček)	79
STENOMICRIDAE	(Jindřich Roháček)	79
ASTEIIDAE	(Jindřich Roháček)	79
BRAULIDAE	(Jindřich Roháček)	79
CARNIDAE	(Jindřich Roháček)	79
TETHINIDAE	(Jindřich Roháček)	79
MILICHIIDAE	(Jindřich Roháček)	80
CHLOROPIDAE	(Jindřich Roháček)	80
HELEOMYZIDAE	(Vladislav Martinek)	82
TRIXOSCELIDAE	(Jindřich Roháček)	83
CHYROMYIDAE	(Jindřich Roháček)	83
SPHAEROCERIDAE	(Jindřich Roháček)	84
CURTONOTIDAE	(Jan Máca)	86
CAMILLIDAE	(Jan Máca)	86
DROSOPHILIDAE	(Jan Máca)	86
DIASTATIDAE	(Jindřich Roháček & Jan Máca)	87
EPHYDRIDAE	(Tadeusz Zatwarniecki)	87
SCATHOPHAGIDAE	(František Šifner)	89
ANTHOMYIIDAE	(Rudolf Rozkošný)	90
FANNIIDAE	(František Gregor & Rudolf Rozkošný)	93
MUSCIDAE	(František Gregor)	93
CALLIPHORIDAE	(František Gregor & Rudolf Rozkošný)	96
RHINOPHORIDAE	(Juraj Čepelák & Rudolf Rozkošný)	97
SARCOPHAGIDAE	(Dalibor Povolný)	98
TACHINIDAE	(Juraj Čepelák & Jaromír Vaňhara)	100
GASTEROPHILIDAE	(Jan Minář)	106
OESTRIDAE	(Jan Minář)	107
HYPODERMATIDAE	(Jan Minář)	107
HIPPOBOSCIDAE	(Josef Chalupský & Dalibor Povolný)	107
NYCTERIBIIDAE	(Karel Hůrka)	107
References		109
Index		113

suborder **NEMATOCERA**

TIPULOMORPHA

Tipuloidea

Trichoceridae
Limoniidae
Pediidae
Cylindrotomidae
Tipulidae

BLEPHARICEROMORPHA

Blepharicerioidea

Blephariceridae

AXYMYIOMORPHA

Axymyioidea

Axymyiidae

BIBIONOMORPHA

Bibionoidea

Bibionidae
Pleciidae
Hesperinidae

Sciaroidea

Bolitophilidae
Diadocidiidae
Ditomyiidae
Keroplidae
Macroceridae
Mycetophilidae
Sciariidae
Cecidomyiidae

PSYCHODOMORPHA

Psychodoidea

Psychodidae

Anisopodoidea

Anisopodidae
Mycetobiidae

Scatopsoidea

Scatopsidae
Synneuridae
Canthyluscelidae

PTYCHOPTEROMORPHA

Ptychopteroidea

Ptychopteridae

CULICOMORPHA

Culicoidea

Dixidae
Chaoboridae
Culicidae
Chironomoidea
Chironomidae
Ceratopogonidae
Simuliidae
Thaumaleidae

suborder **BRACHYCERA**

XYLOPHAGOMORPHA

Xylophagoidea

Xylophagidae
Coenomyiidae

TABANOMORPHA

Tabanoidea

Athericidae
Rhagionidae
Tabanidae
Stratiomyoidea
Xylomyidae
Stratiomyidae

ASILOMORPHA

Nemestrinoidea

Acroceridae

Asiloidea

Mythicomyiidae
Phthiriidae
Bombyliidae
Hilarimorphidae
Therevidae
Scenopinidae
Asilidae

MUSCOMORPHA

ORTHOGENYA

Empidoidea

Empididae
Hybotidae
Dolichopodidae
Atelestidae

CYCLORRHAPHA

ASCHIZA

Platypezoidea

Opetidae
Platypezidae
Phoridae

Lonchopteroidea

Lonchopteridae

Syrphoidea

Syrphidae
Pipunculidae

SCHIZOPHORA

ACALYPTRATAE

Nerioidea

Pseudopomyzidae
Micropezidae

Diopsoidea

Tanypezidae
Strongylophthalmyiidae
Megamerinidae
Psilidae

Conopoidea

Conopidae

Tephritoidea

Lonchacidae
Pallopteridae
Piophilidae
Otitidae
Ulidiidae

Platystomatidae

Tephritidae

Lauxanoidea

Lauxaniidae
Chamaemyiidae
Crenifaniidae

Sciomyzoidea

Dryomyzidae
Phacomyiidae
Sciomyzidae
Sepsidae

Opomyzoidea

Clusiidae
Acartophthalmidae
Odiniidae
Agromyzidae
Opomyzidae
Anthomyzidae
Aulacigastridae
Periscelididae
Stenomericidae
Asteiidae

Carnoidea

Braulidae
Carnidae
Tethinidae
Milichiidae
Chloropidae

Sphaeroceroidea

Heleomyzidae
Trixoscelididae
Chyromyzidae
Sphaeroceridae

Ephydroidea

Curionotidae
Camillidae
Drosophilidae
Diastatidae
Ephydriidae

CALYPTRATAE

Muscoidea

Scathophagidae
Anthomyiidae
Fanniidae
Muscidae

Calliphoroidea

Calliphoridae
Rhizophoridae
Sarcophagidae
Tachinidae

Oestroidea

Gasterophilidae
Oestridae
Hypodermatidae

Hippoboscidea

Hippoboscidae
Nycteribiidae

TRICHOCERIDAE

Jaroslav Starý & Jaroslav Martinovský

Diazosma Bergroth, 1913*hirtipenne* (Siebke, 1863) B M S*seclusum* Starý & Martinovský, 1993 M S*Trichocera* Meigen, 1803sg. *Metatrachocera* Dahl, 1966*candida* Dahl, 1976 S*dahlæ* Mendl, 1971 M S*forcipula* P. Nielsen, 1920 B M S*inexplorata* (Dahl, 1967) M*lutea* Becher, 1886 S*sibirica* Edwards, 1920 M Ssg. *Trichocera* s. str.*annulata* Meigen, 1818 B M S*carpathica* Starý & Martinovský, 1996 S*hiemalis* (De Geer, 1776) B M S*implicata* Dahl, 1976 M S*maculipennis* Meigen, 1818 B M S*major* Edwards, 1921 B M S*obtusa* Starý & Martinovský, 1996 M S*parva* Meigen, 1804 B M S*pubescens* Starý & Martinovský, 1996 B M S*regelationis* (Linnaeus, 1758) B M S*rufescens* Edwards, 1921 B M S*rufidentata* Edwards, 1938 B M S*saltator* (Harris, 1776) B M S*sparsa* Starý & Martinovský, 1996 B M S*trimaculata* (Zetterstedt, 1838) B M S*verralli* (Bergroth, 1912) M S*Epiphragma* Osten Sacken, 1860*ocellare* (Linnaeus, 1761) B M S*Euphyllidorea* Alexander, 1972*aperta* (Verrall, 1887) B M S*dispar* (Meigen, 1818) M S*lineola* (Meigen, 1804) B M S*meigenii* (Verrall, 1887) B M S*phaeostigma* (Schummel, 1829) B M S*Eutonia* van der Wulp, 1874*barbipes* (Meigen, 1804) B M S*Hexatoma* Latreille, 1809sg. *Coreozelia* Enderlein, 1936*cimicoides* (Scopoli, 1763) B M Ssg. *Eriocera* Macquart, 1838*chirothecata* (Scopoli, 1763) B M Ssg. *Hexatoma* s. str.*bicolor* (Meigen, 1818) B M S*fuscipennis* (Curtis, 1836) B M S*gaedii* (Meigen, 1830) B S*nubeculosa* (Burmeister, 1829) B S*vittata* (Meigen, 1830) B S*Idioptera* Macquart, 1834*linnei* Oosterbroek, 1992 B*pulchella* (Meigen, 1830) B M S*Limnophila* Macquart, 1834sg. *Limnophila* s. str.*arnoudi* Theowald, 1971 M S*pictipennis* (Meigen, 1818) B M S*schranksi* Oosterbroek, 1992 B M S*Neolimnomyia* Séguy, 1937sg. *Brachylimnophila* Alexander, 1966*nemoralis* (Meigen, 1818) B M Ssg. *Neolimnomyia* Séguy, 1937*batava* (Edwards, 1938) B M S*filata* (Walker, 1856) M S*Paradelphomyia* Alexander, 1936sg. *Oxyrhiza* de Meijere, 1946*czizekiana* Starý, 1971 B M S*ecalcarata* (Edwards, 1938) M S*fuscata* (Loew, 1873) B M S*nigrina* (Lackschewitz, 1940) B M*senilis* (Haliday, 1833) B M S*Phyllidorea* Bigot, 1854sg. *Paraphyllidorea* Savchenko, 1986*fulvonervosa* (Schummel, 1829) B M Ssg. *Phyllidorea* s. str.*abdominalis* (Staeger, 1840) B M S*bicolor* (Meigen, 1804) B M S*ferruginea* (Meigen, 1818) B M S*heterogyna* (Bergroth, 1913) B*longicornis* (Schummel, 1829) B M S*nervosa* (Schummel, 1829) B M*squalens* (Zetterstedt, 1838) B M S*Pilaria* Sintenis, 1889*decolor* (Zetterstedt, 1851) B M S

LIMONIIDAE

Jaroslav Starý

DACTYLOLABINAE

Dactylolabis Osten Sacken, 1860sg. *Dactylolabis* s. str.*denticulata* (Bergroth, 1891) S*sexmaculata* (Macquart, 1826) B S*subdilata* Starý, 1969 M*transversa* (Meigen, 1804) B M S*wodzickii* (Nowicki, 1867) S

LIMNOPHILINAE

Adelphomyia Bergroth, 1891*punctum* (Meigen, 1818) S*Austrolimnophila* Alexander, 1920sg. *Archilimnophila* Alexander in Curran, 1934*unica* (Osten Sacken, 1869) B M Ssg. *Austrolimnophila* s. str.*ochracea* (Meigen, 1804) B M S*Eloeophila* Rondani, 1856*apicata* (Loew, 1871) B M S*maculata* (Meigen, 1804) B M S*miliaria* (Egger, 1863) B M S*mundata* (Loew, 1871) M S*submarmorata* (Verrall, 1887) B M S

<i>discicollis</i> (Meigen, 1818)	B M S	<i>pederi</i> Tjeder, 1969	M
<i>fuscipennis</i> (Meigen, 1818)	B M S	<i>sordida</i> Zetterstedt, 1838	B M S
<i>meridiana</i> (Staeger, 1840)	B M S	<i>squalida</i> Loew, 1871	B M S
<i>nigropunctata</i> (Agrell, 1945)	M S	<i>tenuirama</i> Savchenko, 1972	S
<i>scutellata</i> (Staeger, 1840)	B M S	<i>verralli</i> Edwards, 1921	M S
Prionolabis Osten Sacken, 1860		sg. Mesocyphona Osten Sacken, 1869	
<i>hospes</i> (Egger, 1863)	B M S	<i>bivittata</i> (Loew, 1873)	B M S
Pseudolimnophila Alexander, 1919		<i>fossarum</i> (Loew, 1873)	B M S
<i>lucorum</i> (Meigen, 1818)	B M S	Gnophomyia Osten Sacken, 1860	
<i>sepium</i> (Verrall, 1886)	B M S	<i>elsneri</i> Starý, 1983	S
		<i>lugubris</i> (Zetterstedt, 1838)	B M S
CHIONEINAE		<i>viridipennis</i> (Gimmerthal, 1847)	M S
Arctoonopa Alexander, 1955		Gonempeda Alexander, 1924	
<i>melampodiv</i> (Loew, 1873)	M S	<i>flava</i> (Schummel, 1829)	B M S
Cheilotrichia Rossi, 1848		Gonomyia Meigen, 1818	
sg. Cheilotrichia s. str.		sg. Gonomyia s. str.	
<i>cinerea</i> (Strobl, 1898)	S	<i>bifida</i> Tonnoir in Goetghebuer	
<i>imbuta</i> (Meigen, 1818)	B M S	& Tonnoir, 1920	M S
<i>valai</i> Starý, 1992	S	<i>conoviensis</i> Barnes, 1924	B M S
sg. Empeda Osten Sacken, 1869		<i>dentata</i> de Meijere, 1920	M
<i>affinis</i> (Lackschewitz, 1927)	B M S	<i>lucidula</i> de Meijere, 1920	B M S
<i>caerulea</i> Starý, 1987	B M S	<i>recta</i> Tonnoir in Goetghebuer	
<i>cinerascens</i> (Meigen, 1804)	B M S	& Tonnoir, 1920	B M S
<i>minima</i> (Strobl, 1898)	M S	<i>simplex</i> Tonnoir in Goetghebuer	
<i>neglecta</i> (Lackschewitz, 1927)	B M S	& Tonnoir, 1920	B M S
<i>staryi</i> Mendl, 1973	B M S	<i>tenella</i> (Meigen, 1818)	B M S
Chionea Dalman, 1816		sg. Prolipophleps Savchenko, 1972	
sg. Chionea s. str.		<i>abbreviata</i> Loew, 1873	B M S
<i>araneoides</i> Dalman, 1816	B M S	<i>divergens</i> Bangerter, 1947	S
sg. Sphaeconophilus Becker, 1912		sg. Teuchogonomyia Alexander, 1968	
<i>botosaneanui</i> (Burghele-Bălăcescu, 1969)	M S	<i>edwardsi</i> Lackschewitz, 1925	S
<i>hutescens</i> Lundström, 1907	B M S	<i>ithyphallus</i> Lackschewitz, 1935	S
Crypteria Bergroth, 1913		Hoplolabis Osten Sacken, 1869	
<i>limnophiloides</i> Bergroth, 1913	B M S	sg. Lunaria Savchenko, 1982	
Ellipteroides Becker, 1907		<i>idiophallus</i> (Savchenko, 1973)	S
sg. Ellipteroides s. str.		sg. Parilisia Savchenko in Savchenko	
<i>lateralis</i> (Macquart, 1835)	M S	& Krivolutskaya, 1976	
sg. Protogonomyia Alexander, 1934		<i>areolata</i> (Siebke, 1872)	M S
<i>adrastea</i> Starý & Mendl, 1984	M	<i>mannheimi</i> (Mendl, 1974)	S
<i>alboscuteclatus</i> (von Roser, 1840)	M S	<i>punctigera obtusiapex</i> (Savchenko, 1982)	S
<i>limbatus</i> (von Roser, 1840)	M S	<i>sachalina</i> (Alexander, 1924)	M S
Erioconopa Starý, 1976		<i>sororcula</i> (Lackschewitz, 1940)	S
<i>diuturna</i> (Walker, 1848)	B M S	<i>spinosa</i> (P.Nielsen, 1953)	M S
<i>interposita</i> Starý, 1976	S	<i>subalpina</i> (Bangerter, 1947)	M S
<i>trivialis</i> (Meigen, 1818)	B M S	<i>vicina</i> (Tonnoir in Goetghebuer	
Erioptera Meigen, 1803		& Tonnoir, 1920)	B M S
sg. Erioptera s. str.		<i>yezoana</i> (Alexander, 1924)	M S
<i>divisa</i> (Walker, 1848)	B M S	Idiocera Dale, 1842	
<i>flavata</i> (Westhoff, 1882)	B M S	sg. Euptilostena Alexander, 1938	
<i>flavissima</i> Starý, 1972	M	<i>jucunda</i> (Loew, 1873)	M S
<i>fuscipennis</i> Meigen, 1818	B M S	sg. Idiocera s. str.	
<i>fusculenta</i> Edwards, 1938	B M S	<i>bradleyi</i> (Edwards, 1939)	M S
<i>griseipennis</i> Meigen, 1838	B M S	<i>pulchripennis</i> (Loew, 1856)	M S
<i>limbata</i> Loew, 1873	M S	<i>punctata</i> (Edwards, 1938)	M S
<i>lutea</i> Meigen, 1804	B M S	<i>sziladyi</i> (Lackschewitz, 1940)	M S
<i>minor distinguenda</i> Starý, 1983	S		
<i>minor minor</i> de Meijere, 1920	M S		

<i>Ilisia</i> Rondani, 1856				sg. <i>Ormosia</i> s. str.			
<i>inermis</i> Mendl, 1979			S	<i>aciculata</i> Edwards, 1921			S
<i>maculata</i> (Meigen, 1804)	B	M	S	<i>albitibia</i> Edwards, 1921		B	M S
<i>occocata</i> Edwards, 1936		M	S	<i>bicornis</i> (de Meijere, 1920)			M S
<i>Molophilus</i> Curtis, 1833				<i>bifida</i> (Lackschewitz, 1940)		B	M S
<i>appendiculatus</i> (Staeger, 1840)	B	M	S	<i>bihamata</i> Lackschewitz, 1935			M
<i>ater</i> (Meigen, 1804)	B	M	S	<i>clavata</i> (Tonnoir in Goetghebuer			
<i>bifidus</i> Goetghebuer in Goetghebuer				& Tonnoir, 1920)		B	M S
& Tonnoir, 1920	B	M	S	<i>depilata</i> Edwards, 1938		B	M S
<i>bihamatus</i> de Meijere, 1918	B	M	S	<i>egena</i> (Bergroth, 1891)		B	M S
<i>bischofi</i> Lackschewitz, 1940			M S	<i>fascipennis</i> (Zetterstedt, 1838)			M S
<i>brevihamatus</i> Bangerter, 1947	B	M	S	<i>hederae</i> (Curtis, 1835)		B	M S
<i>carstensis</i> Starý, 1971			M S	<i>lineata</i> (Meigen, 1804)		B	M S
<i>cinereifrons</i> de Meijere, 1920	B	M	S	<i>loxia</i> Starý, 1983			M S
<i>corniger</i> de Meijere, 1920			M S	<i>moravica</i> Starý, 1969			M S
<i>crassipygus</i> de Meijere, 1918	B	M	S	<i>nana</i> Starý, 1974			M
<i>curvatus</i> Tonnoir in Goetghebuer				<i>nodulosa</i> (Macquart, 1826)		B	M
& Tonnoir, 1920			M S	<i>pseudosimilis</i> (Lundström, 1912)		B	M
<i>czizeki</i> Lackschewitz in Czižek, 1931			M S	<i>rostrifera</i> Savchenko, 1973			M S
<i>flavus</i> Goetghebuer in Goetghebuer				<i>ruficauda</i> (Zetterstedt, 1838)		B	M S
& Tonnoir, 1920	B	M	S	<i>staegeriana</i> Alexander, 1953		B	M S
<i>fluviatilis</i> Bangerter, 1947			M S	<i>Phyllolabis</i> Osten Sacken, 1877			
<i>griseus</i> (Meigen, 1804)	B	M	S	<i>macroura</i> (Siebke, 1863)			S
<i>lackschewitzianus</i> Alexander, 1953			S	<i>Rhabdomastix</i> Skuse, 1890			
<i>maurus</i> Lackschewitz, 1925			S	sg. <i>Palaeogonomyia</i> Meunier, 1899			
<i>medius</i> de Meijere, 1918	B	M	S	<i>hirticornis</i> (Lackschewitz, 1940)			S
<i>niger</i> Goetghebuer in Goetghebuer				sg. <i>Sacandaga</i> Alexander, 1911			
& Tonnoir, 1920			M S	<i>laeta</i> (Loew, 1873)			M S
<i>nodicornis</i> Lackschewitz, 1935			M S	<i>lurida</i> (Loew, 1873)			S
<i>obscurus</i> (Meigen, 1818)	B	M	S	<i>subparva</i> Starý, 1971		B	M S
<i>occultus</i> de Meijere, 1918	B	M	S	<i>Rhypholophus</i> Kolenati, 1860			
<i>ochraceus</i> (Meigen, 1818)	B	M	S	<i>bifurcatus</i> Goetghebuer in Goetghebuer			
<i>pleuralis</i> de Meijere, 1920	B	M	S	& Tonnoir, 1920		B	M S
<i>priapoides</i> Starý, 1971			M S	<i>haemorrhoidalis</i> (Zetterstedt, 1838)		B	M S
<i>priapus</i> Lackschewitz, 1935	B	M	S	<i>lichtwardti</i> (Lackschewitz, 1935)			M S
<i>propinquus</i> (Egger, 1863)	B	M	S	<i>phryganopterus</i> Kolenati, 1860		B	M S
<i>pseudopropinquus</i> Mendl, 1973			S	<i>varius</i> (Meigen, 1818)		B	
<i>pullus</i> Lackschewitz, 1927			M S	<i>Scleroprocta</i> Edwards, 1938			
<i>repentinus</i> Starý, 1971			M S	<i>pentagonalis</i> (Loew, 1873)			M S
<i>savishenkoi</i> Starý, 1972			M S	<i>sororcula</i> (Zetterstedt, 1851)			M S
<i>scutellatus</i> Goetghebuer, 1929	B	M	S	<i>Symplecta</i> Meigen, 1830			
<i>serpentiger</i> Edwards, 1938			M S	sg. <i>Psiloconopa</i> Zetterstedt, 1838			
<i>stroblanus</i> P. Nielsen, 1953			M S	<i>bizarrea</i> Starý, 1992			S
<i>terravi</i> Starý, 1992			S	<i>meigeni</i> (Zetterstedt, 1838)		B	M
<i>tjederi</i> Starý, 1968			S	<i>pusilla</i> (Schiner, 1865)			M S
<i>undulatus</i> Tonnoir in Goetghebuer				<i>stictica</i> (Meigen, 1818)		B	M S
& Tonnoir, 1920			M S	sg. <i>Symplecta</i> s. str.			
<i>vafer</i> Lackschewitz, 1940			S	<i>chosenensis</i> (Alexander, 1940)			M S
<i>varispinus</i> Starý, 1971			M S	<i>grata</i> Loew, 1873			M S
<i>Neolimnophila</i> Alexander, 1920				<i>hybrida</i> (Meigen, 1804)		B	M S
<i>bergrothi</i> (Kuntze, 1919)			S	sg. <i>Trimicra</i> Osten Sacken, 1861			
<i>carteri</i> (Tonnoir in Goetghebuer				<i>pilipes</i> (Fabricius, 1787)		B	M S
& Tonnoir, 1921)	B	M	S	<i>Tasiocera</i> Skuse, 1890			
<i>placida</i> (Meigen, 1830)			M S	sg. <i>Dasymolophilus</i> Goetghebuer in			
<i>Ormosia</i> Rondani, 1856				Goetghebuer & Tonnoir, 1920			
sg. <i>Oreophila</i> Lackschewitz, 1935				<i>exigua</i> Savchenko, 1973			M S
<i>bergrothi</i> (Strobl, 1895)			S	<i>fuscescens</i> (Lackschewitz, 1940)			M S

<i>jenkinsoni</i> Freeman, 1951	M S	<i>nigristigma</i> P. Nielsen, 1919	B M S
<i>murina</i> (Meigen, 1818)	B M S	<i>stigmatica</i> (Meigen, 1830)	B M S
<i>robusta</i> (Bangerter, 1947)	M S	sg. <i>Melanolimonia</i> Alexander, 1965	
LIMONIINAE		<i>caledonica</i> Edwards, 1926	B M S
<i>Achyrolimonia</i> Alexander, 1965		<i>morio</i> (Fabricius, 1787)	B M S
<i>coeiana</i> (P. Nielsen, 1959)	M S	<i>occidua</i> Edwards, 1926	S
<i>decemmaculata</i> (Loew, 1873)	B M S	<i>rufiventris</i> (Strobl, 1901)	B M S
<i>neonebulosa</i> (Alexander, 1924)	B M S	<i>stylifera</i> Lackschewitz, 1928	S
<i>Antocha</i> Osten Sacken, 1860		sg. <i>Numantia</i> Bigot, 1854	
sg. <i>Antocha</i> s. str.		<i>fusca</i> (Meigen, 1804)	B M S
<i>vitripennis</i> (Meigen, 1830)	B M S	sg. <i>Sivalimnobia</i> Alexander, 1963	
sg. <i>Orimargula</i> Mik, 1883		<i>aquosa</i> Verrall, 1886	S
<i>alpigena</i> (Mik, 1883)	M S	<i>Dicranoptycha</i> Osten Sacken, 1860	
<i>Atypophthalmus</i> Brunetti, 1911		<i>cinerascens</i> (Meigen, 1818)	B M S
sg. <i>Atypophthalmus</i> s. str.		<i>fuscescens</i> (Schummel, 1829)	B M S
<i>imustus</i> (Meigen, 1818)	M S	<i>livescens</i> Loew, 1871	B M S
sg. <i>Microlimonia</i> Savchenko in Savchenko & Krivolutskaya, 1976		<i>paralivescens</i> Starý, 1972	M S
<i>machidai</i> (Alexander, 1921)	B M S	<i>pseudocinerea</i> Starý, 1972	S
<i>Dicranomyia</i> Stephens, 1829		<i>Discobola</i> Osten Sacken, 1865	
sg. <i>Dicranomyia</i> s. str.		<i>annulata</i> (Linnaeus, 1758)	B M S
<i>aperta</i> Wahlgren, 1904	S	<i>caesarea</i> (Osten Sacken, 1854)	M S
<i>autumnalis</i> (Staeger, 1840)	B M S	<i>parvispinula</i> (Alexander, 1947)	M S
<i>chorea</i> (Meigen, 1818)	B M S	<i>Elephantomyia</i> Osten Sacken, 1860	
<i>conchifera</i> (Strobl, 1901)	M S	<i>edwardsi</i> Lackschewitz, 1932	M S
<i>didyma</i> (Meigen, 1804)	B M S	<i>krivosheinae</i> Savchenko, 1976	S
<i>distendens</i> Lundström, 1912	B M S	<i>Elliptera</i> Schiner, 1863	
<i>frontalis</i> (Staeger, 1840)	B M S	<i>hungarica</i> Madarassy, 1881	S
<i>halterata</i> Osten Sacken, 1869	M	<i>omissa</i> Schiner, 1863	B M S
<i>handlirschi</i> Lackschewitz, 1928	B M S	<i>Helius</i> Lepeletier & Serville, 1828	
<i>incisurata</i> Lackschewitz, 1928	S	<i>flavus</i> (Walker, 1856)	B M S
<i>longipennis</i> (Schummel, 1829)	B M S	<i>longirostris</i> (Meigen, 1818)	B M S
<i>lucida</i> de Meijere, 1918	M S	<i>pallirostris</i> Edwards, 1921	M S
<i>luteipennis</i> Goetghebuer in Goetghebuer & Tonnoir, 1920	B M S	<i>Libnotes</i> Westwood, 1876	
<i>mitis</i> (Meigen, 1830)	B M S	sg. <i>Afrolimonia</i> Alexander, 1965	
<i>modesta</i> (Meigen, 1818)	B M S	<i>ladogensis</i> (Lackschewitz in Lackschewitz & Pagast, 1940)	S
<i>omissinervis</i> de Meijere, 1918	M S	<i>Limonia</i> Meigen, 1803	
<i>ornata</i> (Meigen, 1818)	B M S	<i>albifrons</i> (Meigen, 1818)	B M S
<i>radegasti</i> Starý, 1993	M	<i>alpicola</i> (Lackschewitz, 1928)	S
<i>sera</i> (Walker, 1848)	B M S	<i>aquilina</i> Starý, 1984	M
<i>strobl</i> Pagast in Lackschewitz & Pagast, 1941	M S	<i>flavipes</i> (Fabricius, 1787)	B M S
<i>ventralis</i> (Schummel, 1829)	B M S	<i>hercegovinae</i> (Strobl, 1898)	B M S
<i>zernyi</i> Lackschewitz, 1928	B	<i>interjecta</i> Starý, 1974	S
sg. <i>Glochina</i> Meigen, 1830		<i>macrostigma</i> (Schummel, 1829)	B M S
<i>kinensis</i> (Alexander, 1934)	M S	<i>nigropunctata</i> (Schummel, 1829)	B M S
<i>schineri</i> Lackschewitz, 1928	B M S	<i>mibeculosa</i> Meigen, 1804	B M S
<i>sericata</i> (Meigen, 1830)	B S	<i>pannonica</i> (Kowarz, 1868)	B S
<i>transsilvanica</i> Lackschewitz, 1928	M S	<i>phragmitidis</i> (Schränk, 1781)	B M S
<i>tristis</i> (Schummel, 1829)	B M S	<i>splendens</i> Kuntze, 1920	S
sg. <i>Idiopyga</i> Savchenko, 1987		<i>stigma</i> (Meigen, 1818)	B M S
<i>alpina</i> Bangerter, 1948	M S	<i>sylvicola</i> (Schummel, 1829)	B M S
<i>danica</i> Kuntze, 1919	B M S	<i>taurica</i> (Strobl, 1895)	B M S
<i>halterella</i> Edwards, 1921	B M S	<i>trivittata</i> (Schummel, 1829)	B M S
<i>lackschewitzi</i> Edwards, 1928	S	<i>Lipsothrix</i> Loew, 1873	
		<i>ecucullata</i> Edwards, 1938	M S
		<i>errans</i> (Walker, 1848)	B M S

<i>nobilis</i> Loew, 1873	B M S	<i>riedeli nielseni</i> (Slipka, 1955)	M S
<i>remota</i> (Walker, 1848)	B M S	<i>riedeli riedeli</i> (Lackschewitz, 1940)	M
<i>Metalimnobia</i> Matsumura, 1911		<i>straminea</i> (Meigen, 1838)	B M S
<i>bifasciata</i> (Schränk, 1781)	B M S	<i>zernyi pallens</i> Savchenko, 1978	M S
<i>quadrinotata</i> (Linnaeus, 1761)	B M S	sg. <i>Pedicia</i> s. str.	
<i>quadrinotata</i> (Meigen, 1818)	B M S	<i>rivosa</i> (Linnaeus, 1758)	B M S
<i>zetterstedti</i> (Tjeder, 1968)	B M S	<i>Tricyphona</i> Zetterstedt, 1837	
<i>Neolimonia</i> Alexander, 1964		<i>alpigena</i> (Strobl, 1909)	B M S
<i>dumetorum</i> (Meigen, 1804)	B M S	<i>alticola</i> Strobl, 1909	S
<i>Orimarga</i> Osten Sacken, 1869		<i>contraria</i> Bergroth, 1888	B M S
<i>attenuata</i> (Walker, 1848)	B M S	<i>immaculata</i> (Meigen, 1804)	B M S
<i>juvenilis</i> (Zetterstedt, 1851)	S	<i>livida</i> Madarassy, 1881	M S
<i>virgo</i> (Zetterstedt, 1851)	S	<i>schummeli</i> Edwards, 1921	B M S
<i>Rhipidia</i> Meigen, 1818		<i>unicolor</i> (Schummel, 1829)	B M S
<i>maculata</i> Meigen, 1818	B M S	<i>Ula</i> Haliday, 1833	
<i>punctipennis</i> Mik, 1887	S	<i>bolitophila</i> Loew, 1869	M S
<i>uniseriata</i> Schiner, 1864	M S	<i>ciscuncta</i> Stary, 1997	S
<i>Thaumastoptera</i> Mik, 1866		<i>mixta</i> Stary, 1983	B S
<i>calceata</i> Mik, 1866	M S	<i>mollissima</i> Haliday, 1833	B M S
		<i>sylvatica</i> (Meigen, 1818)	B M S

PEDICIIDAE

Jaroslav Stary

<i>Dicranota</i> Zetterstedt, 1838			
sg. <i>Dicranota</i> s. str.			
<i>bimaculata</i> (Schummel, 1829)	B M S		
<i>guerini</i> Zetterstedt, 1838	S		
sg. <i>Ludicia</i> Hutson & Vane-Wright, 1969			
<i>lucidipennis</i> (Edwards, 1921)	B M S		
sg. <i>Paradicranota</i> Alexander in Curran, 1934			
<i>brevicornis</i> Bergroth, 1891	B M S		
<i>brevitarsis</i> Bergroth, 1891	B M S		
<i>candelisequa</i> Stary, 1981	M S		
<i>consimilis</i> Mendl, 1987	S		
<i>flammatra</i> Stary, 1981	B M S		
<i>gracilipes</i> Wahlgren, 1905	B M S		
<i>landrocki</i> Cizik, 1931	M S		
<i>martinovskii</i> Stary, 1974	M S		
<i>mikiana</i> Lackschewitz, 1940	M		
<i>minuta</i> Lackschewitz, 1940	M S		
<i>pallens</i> Lackschewitz, 1940	M S		
<i>pavida</i> (Haliday, 1833)	B M S		
<i>robusta</i> Lundström, 1912	B M		
<i>schistacea</i> Lackschewitz, 1940	M S		
<i>simulans</i> Lackschewitz, 1940	B M S		
<i>subtilis</i> Loew, 1871	M S		
sg. <i>Rhaphidolabis</i> Osten Sacken, 1869			
<i>exclusa</i> (Walker, 1848)	B S		
<i>Nasiternella</i> Wahlgren, 1904			
<i>varinervis</i> (Zetterstedt, 1851)	S		
<i>Pedicia</i> Latreille, 1809			
sg. <i>Amalopsis</i> Haliday, 1856			
<i>occulta</i> (Meigen, 1830)	B M S		
sg. <i>Crunobia</i> Kolenati, 1859			
<i>littoralis</i> (Meigen, 1804)	B M S		

CYLINDROTOMIDAE

Jaroslav Stary

<i>Cylindrotoma</i> Macquart, 1834			
<i>distinctissima distinctissima</i> (Meigen, 1818)		B M S	
<i>Diogma</i> Edwards, 1938			
<i>glabrata glabrata</i> (Meigen, 1818)		B M S	
<i>Phalacrocer</i> Schiner, 1863			
<i>replicata</i> (Linnaeus, 1758)		B M	
<i>Triogma</i> Schiner, 1863			
<i>trisulcata</i> (Schummel, 1829)		B M S	

TIPULIDAE

Jaroslav Martinovský

DOLICHOPEZINAE

DOLICHOPEZINI

<i>Dolichopeza</i> Curtis, 1825			
<i>albipes</i> (Ström, 1768)		B M S	

TIPULINAE

PRIONOCERINI

<i>Prionocera</i> Loew, 1844			
<i>chosenicola</i> Alexander, 1945		M S	
= <i>tjederi</i> Mannheims, 1951			
<i>pubescens</i> Loew, 1844		B	
<i>turcica</i> (Fabricius, 1787)		B M S	
<i>subsericornis</i> (Zetterstedt, 1851)		M	

TIPULINI

<i>Nephrotoma</i> Meigen, 1803			
<i>aculeata</i> (Loew, 1871)		B M S	

<i>analis</i> (Schummel, 1833)	B M S	<i>pannonia</i> Loew, 1873	M S
<i>appendiculata</i> (Pierre, 1919)	B M S	<i>peliosigma</i> Schummel, 1833	B M S
<i>austriaca</i> (Mannheims & Theowald, 1959)	M S	<i>recticornis</i> Schummel, 1833	M
<i>cornicina</i> (Linnaeus, 1758)	B M S	<i>selene</i> Meigen, 1830	B M S
<i>crocata</i> (Linnaeus, 1758)	B M S	<i>soosi</i> Mannheims, 1954	B M S
<i>croceiventris lindneri</i> (Mannheims, 1951)	B M S	<i>truncata</i> Loew, 1873	B M S
<i>dorsalis</i> (Fabricius, 1782)	B M S	<i>vernalis</i> Meigen, 1804	B M S
<i>flavescens</i> (Linnaeus, 1758)	B M S	<i>verrucosa</i> Pierre, 1919	M
<i>guestfalica</i> (Westhoff, 1879)	B M S	sg. <i>Mediotipula</i> Pierre, 1924	
<i>lamellata</i> (Riedel, 1910)	B M S	<i>mikiana</i> Bergroth, 1888	B M S
<i>humilicornis</i> (Schummel, 1833)	B M S	<i>siebkei</i> Zetterstedt, 1852	S
<i>moravica</i> Martinovský, 1971	M S	<i>stigmatella</i> Schummel, 1833	B M S
<i>pratensis</i> (Linnaeus, 1758)	B M S	sg. <i>Odonatisca</i> Savchenko, 1956	
<i>quadrifaria</i> (Meigen, 1804)	B M S	<i>nodicornis</i> Meigen, 1818	B M S
<i>quadristriata</i> (Schummel, 1833)	M S	sg. <i>Platytipula</i> Matsumura, 1916	
<i>scalaris</i> (Meigen, 1818)	B M S	<i>autumnalis</i> Loew, 1864	B
<i>scurra</i> (Meigen, 1818)	B M S	<i>luteipennis</i> Meigen, 1830	B M S
<i>submaculosa</i> Edwards, 1928	M S	<i>melanoceros</i> Schummel, 1833	B M S
<i>tenuipes</i> (Riedel, 1910)	B M S	sg. <i>Pterelachisus</i> Rondani, 1842	
<i>Nigrotipula</i> Hutson & Vane-Wright, 1969		<i>apicispina</i> Alexander, 1934	S
<i>nigra</i> (Linnaeus, 1758)	B M S	<i>cinereocincta</i> Lundström, 1907	B M S
<i>Tipula</i> Linnaeus, 1758		<i>crassiventris</i> Riedel, 1913	M S
sg. <i>Acutipula</i> Alexander, 1924		<i>irrorata</i> Macquart, 1826	B M S
<i>bosnica</i> Strobl, 1898	M S	<i>laetibasis</i> Alexander, 1934	M
<i>fulvipennis</i> De Geer, 1776	B M S	<i>luridorostris</i> Schummel, 1833	B M S
<i>luna</i> Westhoff, 1879	B M S	<i>mayerduerii</i> Egger, 1863	S
<i>maxima balcanica</i> Vermoolen, 1983	S	<i>neurotica</i> Mannheims, 1966	B M S
<i>maxima maxima</i> Poda, 1761	B M S	<i>pabulina</i> Meigen, 1818	B M S
<i>tenuicornis</i> Schummel, 1833	B M S	<i>pauli</i> Mannheims, 1964	M
<i>vittata</i> Meigen, 1804	B M S	<i>plitviciensis</i> Simova, 1962	S
sg. <i>Beringotipula</i> Savchenko, 1961		<i>pseudocrassiventris</i> Theowald, 1980	M S
<i>unca</i> Wiedemann, 1817	B M S	<i>pseudoirrorata</i> Goetghebuer in Goetghebuer & Tonnoir, 1921	B M S
sg. <i>Dendrotipula</i> Savchenko, 1964		<i>pseudopruinosa</i> Strobl, 1895	S
<i>flavolineata</i> Meigen, 1804	B M S	<i>pseudovariipennis</i> Czižek, 1912	B M S
sg. <i>Emodotipula</i> Alexander, 1966		<i>submarmorata</i> Schummel, 1833	B M S
<i>saginata</i> Bergroth, 1891	B M S	<i>truncorum</i> Meigen, 1830	B M S
sg. <i>Lunatipula</i> Edwards, 1931		<i>varipennis</i> Meigen, 1818	B M S
<i>affinis</i> Schummel, 1833	B M S	sg. <i>Savtshenkia</i> Alexander, 1966	
<i>alpina</i> Loew, 1873	M S	<i>alpium</i> Bergroth, 1888	B M S
<i>borysthenica</i> Savchenko, 1954	S	<i>benesignata</i> Mannheims, 1954	M S
<i>bullata</i> Loew, 1873	S	<i>cheethami</i> Edwards, 1924	B M S
<i>cava</i> Riedel, 1913	B	<i>gimmerthali</i> Lackschewitz, 1925	B M
<i>cinerascens</i> Loew, 1873	S	<i>goriziensis</i> Strobl, 1893	S
<i>fascingulata</i> Mannheims, 1966	S	<i>grisescens</i> Zetterstedt, 1851	B M S
<i>fascipennis</i> Meigen, 1818	B M S	<i>interserta</i> Riedel, 1913	M S
<i>helvola</i> Loew, 1873	B M S	<i>invenusta subinvenusta</i> Slipka, 1950	S
<i>heros</i> Egger, 1863	S	<i>limbata</i> Zetterstedt, 1838	B M S
<i>humilis</i> Staeger, 1840	B S	<i>obsoleta</i> Meigen, 1818	B M S
<i>laetabilis</i> Zetterstedt, 1838	B M S	<i>pagana pagana</i> Meigen, 1818	B
<i>limitata</i> Schummel, 1833	B M S	<i>pagana holoptera</i> Edwards, 1939	B M S
<i>livida</i> Van der Wulp, 1858	B M S	<i>rufina</i> Meigen, 1818	B M S
<i>lunata</i> Linnaeus, 1758	B M S	<i>serrulifera</i> Alexander, 1942	M
<i>magnicauda</i> Strobl, 1895	B	<i>signata</i> Staeger, 1840	B M S
<i>mellea</i> Schummel, 1833	S	<i>simulans</i> Savchenko, 1966	M S
		<i>staegeri</i> P. Nielsen, 1922	M
		<i>subnodicornis</i> Zetterstedt, 1838	B M S

<i>subsignata</i> Lackschewitz, 1933	B M S	BLEPHARICERIDAE	
sg. <i>Schummelia</i> Edwards, 1931		Jaroslav Starý & Ján Gulička	
<i>variicornis</i> (Schummel, 1833)	B M S		
<i>zernyi</i> Mannheims, 1952	B M S	<i>Blepharicera</i> Macquart, 1843	
<i>zonaria</i> Goetghebuer in Goetghebuer & Tonnoir, 1921	M S	<i>fasciata</i> (Westwood, 1842)	B M S
sg. <i>Tipula</i> s. str.		<i>Liponeura</i> Loew, 1844	
<i>hungarica</i> Lackschewitz, 1930	M	<i>brevirostris</i> Loew, 1877	B M S
<i>oleracea</i> Linnaeus, 1758	B M S	<i>cinerascens cinerascens</i> Loew, 1844	B M
<i>orientalis</i> Lackschewitz, 1930	M S	<i>cinerascens minor</i> Bischoff, 1922	M S
<i>paludosa</i> Meigen, 1830	B M S	<i>cordata</i> Vimmer, 1916	B M S
<i>subcunctans</i> Alexander, 1921	B M S	<i>decipiens</i> Bezzi, 1913	M S
sg. <i>Vestiplex</i> Bezzi, 1924		<i>vimmeri</i> Mannheims, 1954	B M S
<i>excisa</i> Schummel, 1833	B M S		
<i>hemiptera strobliana</i> Mannheims, 1966	S	AXYMYIIDAE	
<i>hortorum</i> Linnaeus, 1758	B M S	Jaroslav Martinovský	
<i>montana</i> Curtis, 1834	S	<i>Mesaxymyia</i> Mamaev, 1968	
<i>nubeculosa</i> Meigen, 1804	B M S	<i>kerteszi</i> (Duda, 1930)	S
<i>scripta</i> Meigen, 1830	B M S		
sg. <i>Yamatotipula</i> Matsumura, 1916		BIBIONIDAE	
<i>caesia</i> Schummel, 1833	B M S	Pavel Pecina	
<i>coerulescens</i> Lackschewitz, 1923	M S	<i>Bibio</i> Geoffroy, 1762	
<i>couckeii</i> Tonnoir in Goetghebuer & Tonnoir, 1921	B M S	<i>clavipes</i> Meigen, 1818	B M
<i>fenestrella</i> Theowald, 1980	B M	<i>femoralis</i> Meigen, 1838	B M S
<i>lateralis</i> Meigen, 1804	B M S	<i>ferruginatus</i> (Linnaeus 1767)	B
<i>marginella</i> Theowald, 1980	B M S	<i>fulvipes</i> (Zetterstedt, 1838)	S
<i>montium</i> Egger, 1863	B M S	<i>hortulanus</i> (Linnaeus, 1758)	B M S
<i>pierrei</i> Tonnoir in Goetghebuer & Tonnoir, 1921	B M S	<i>johannis</i> (Linnaeus, 1767)	B M S
<i>pruinosa</i> Wiedemann, 1817	B M S	<i>lanigerus</i> Meigen, 1818	B M S
<i>submontium</i> Theowald & Oosterbroek, 1981	M S	<i>lautaretensis</i> Villeneuve, 1924	M S
		= <i>benesi</i> Pecina, 1962	
CTENOPHORINAE		<i>leucopterus</i> (Meigen, 1804)	B M S
CTENOPHORINI		<i>marci</i> (Linnaeus, 1758)	B M S
<i>Ctenophora</i> Meigen, 1803		<i>nigriventris</i> Haliday, 1833	B M S
sg. <i>Cnemoncosis</i> Enderlein, 1921		<i>pomoniae</i> (Fabricius, 1775)	B M S
<i>fastuosa</i> Loew, 1871	M	<i>reticulatus</i> Loew, 1846	B M S
<i>festiva</i> Meigen, 1804	B M S	<i>varipes</i> Meigen, 1830	B M S
<i>ornata</i> Meigen, 1818	B M S	<i>venosus</i> (Meigen, 1804)	B M S
sg. <i>Ctenophora</i> s. str.		<i>Dilophus</i> Meigen, 1803	
<i>elegans</i> Meigen, 1818	S	<i>antipedalis</i> Wiedemann in Meigen, 1818	M
<i>flaveolata</i> (Fabricius, 1794)	B M S	<i>febrilis</i> (Linnaeus, 1758)	B M S
<i>guttata</i> Meigen, 1818	B M S	<i>femoratus</i> Meigen, 1804	B M S
<i>pectinicornis</i> (Linnaeus, 1758)	B M S	<i>humeralis</i> Zetterstedt, 1850	B M
sg. <i>Xiphuromorpha</i> Savchenko, 1973		PLECIIDAE	
<i>sibirica</i> Portschiński, 1873	S	Jaroslav Martinovský	
<i>Dictenidia</i> Brullé, 1833		<i>Penthetria</i> Meigen, 1803	
<i>bimaculata</i> (Linnaeus, 1761)	B M S	<i>funebis</i> Meigen, 1804	B M S
TANYPTERINI		HESPERINIDAE	
<i>Tanyptera</i> Latreille, 1804		Jaroslav Martinovský	
<i>atrata</i> (Linnaeus, 1758)	B M S	<i>Hesperinus</i> Walker, 1848	
<i>nigricornis</i> (Meigen, 1818)	B M S	<i>imbecillus</i> (Loew, 1858)	not specified

BOLITOPHILIDAE

Vladimír Košel & Jaroslav Martinovský

Bolitophila Meigen, 1818sg. *Bolitophila* s. str.*austriaca* (Mayer, 1950) B M*basicornis* (Mayer, 1951) B M*cinerea* Meigen, 1818 B M S*saundersi* (Curtis, 1836) M S*tenella* Winnertz, 1863 B M Ssg. *Clipisa* Enderlein, 1936*aperta* Lundström, 1914 B M*bimaculata* Zetterstedt, 1838 B M*dubia* Siebke, 1861 M S*edwardsiana* Stackelberg, 1969 S*fumida* Edwards, 1941 B M S*glabrata* Loew, 1869 B M*hybrida* (Meigen, 1804) B M S*maculipennis* Walker, 1836 B M S*mayeri* Plassman, 1986*modesta* Lackschewitz, 1937 B S*occlusa* Edwards, 1913 B M S*pseudohybrida* Landrock, 1912 M*rectangulata* Lundström, 1913 S**DIADOCIDIIDAE**

Vladimír Košel & Jaroslav Martinovský

Diadocidia Ruhe, 1831sg. *Adidocidia* Laštovka & Matile, 1972*valida* Mik, 1874 B M Ssg. *Diadocidia* s. str.*ferruginosa* (Meigen, 1830) B M S*spinosula* Tollel, 1948 B M S**DITOMYIIDAE**

Vladimír Košel & Jaroslav Martinovský

Ditomyia Winnertz, 1846*fasciata* (Meigen, 1818) B M S*macroptera* Winnertz, 1852 B M*Symmerus* Walker, 1848*annulatus* (Meigen, 1830) B M S*nobilis* Lackschewitz, 1937 S**KEROPLATIDAE**

Vladimír Košel, Peter Chandler & Jaroslav Martinovský

Antlemon Loew, 1871sg. *Antlemonopsis* Tollel, 1953*brevimanum* (Loew, 1871) B M S*servulum* (Walker, 1836) M*Asindulum* Latreille, 1805*nigrum* Latreille, 1805 B*Cerotelion* Rondani, 1856*striatum* (Gmelin, 1790) B M*racovitzae* Matile & Burghelle
-Bălăcescu, 1969 B M S*Isoneuromyia* Brunetti, 1912*semirufa* (Meigen, 1818) B M S*Keroplatus* Bosc, 1792*recaumurii* (Dufour, 1839) B*testaceus* (Dalman, 1818) B S*tipuloides* Bosc, 1792 B M*Macrorrhyncha* Winnertz, 1846*collarti* Tollel, 1955 S*flava* Winnertz, 1846 M*Neoplatyura* Malloch, 1928*flava* (Macquart, 1826) M S*Orfelia* Costa, 1857*basalis* (Winnertz, 1863) S*discoloria* (Meigen, 1818) B M S*fasciata* (Meigen, 1804) B M S*nemoralis* (Meigen, 1818) B M S*nigricornis* (Fabricius, 1805) M*unicolor* (Staeger, 1840) B M*Platyura* Meigen, 1803*marginata* Meigen, 1804 B M S*Pyratula* Edwards, 1929*zonata* (Zetterstedt, 1855) B M S*Rocetelion* Matile, 1988*humale* (Zetterstedt, 1850) B M S*Rutylapa* Edwards, 1929*ruficornis* (Zetterstedt, 1851) M*Urytalpa* Edwards, 1929*ochracea* (Meigen, 1818) B M S*rhapsodica* Chandler, 1995 S**MACROCERIDAE**

Vladimír Košel & Jaroslav Martinovský

Macrocera Meigen, 1803*alpicola* Winnertz, 1863 M*anglica* Edwards, 1925 S*angulata* Meigen, 1818 B M S*centralis* Meigen, 1818 B M S*crassicornis* Winnertz, 1863 B M S*fasciata* Meigen, 1804 B M S*fascipennis* Staeger, 1840 S*fastuosa* Loew, 1869 B*inversa* Loew, 1869 M*longibrachiata* Landrock, 1917 M*lutea* Meigen, 1804 B M S*maculata* Meigen, 1818 M*nigricoxa* Winnertz, 1863 M S= *tusca* Loew, 1869*parva* Lundström, 1914 B M S*phalerata* Meigen, 1818 B M S*pilosa* Landrock, 1917 M*pusilla* Meigen, 1830 M

<i>stigma</i> Curtis, 1837	B M S	<i>affinis</i> (Staeger, 1840)	B S
<i>stigmoides</i> Edwards, 1925	B S	= <i>flava</i> (Winnertz, 1863)	
<i>sudetica</i> Landrock, 1923	M	<i>trilineata</i> (Zetterstedt, 1838)	B M S
<i>vittata</i> Meigen, 1830	B M S	sg. <i>Neomycomya</i> Väisänen, 1984	
		<i>fimbriata</i> (Meigen, 1818)	B M
MYCETOPHILIDAE		<i>Neoempheria</i> Osten-Sacken, 1878	
Vladimír Košel, Peter Chandler & Jaroslav Martinovský		<i>lineola</i> (Meigen, 1818)	B M S
		<i>pictipennis</i> (Haliday, 1833)	S
		<i>striata</i> (Meigen, 1818)	B M S
SCIOPHILINAE		SCIOPHILINI	
MYCOMYINI		<i>Acnemis</i> Winnertz, 1863	
<i>Mycomya</i> Rondani, 1856		<i>longipes</i> Winnertz, 1863	M
sg. <i>Cymomya</i> Väisänen, 1984		<i>nitidicollis</i> (Meigen, 1818)	B M S
<i>circumdata</i> (Staeger, 1840)	B M S	<i>Allocotocera</i> Mik, 1886	
sg. <i>Lycomya</i> Väisänen, 1984		<i>pulchella</i> (Curtis, 1837)	B M
<i>pectinifera</i> Edwards, 1924	S	<i>Anacileia</i> Meunier, 1904	
sg. <i>Mycomya</i> s. str.		<i>dispar</i> (Winnertz, 1863)	B
<i>annulata</i> (Meigen, 1818)	M S	<i>dziedzickii</i> (Landrock, 1911)	M
= <i>incisurata</i> (Zetterstedt, 1838)		<i>Azana</i> Walker, 1858	
<i>bicolor</i> (Dziedzicki, 1885)	B M	<i>anomala</i> (Staeger, 1840)	M
<i>brunnea</i> (Dziedzicki, 1885)	B	<i>Leptomorphus</i> Curtis, 1831	
<i>cinerascens</i> (Macquart, 1826)	B M S	<i>quadrimaculatus</i> (Matsumura, 1916)	B
<i>daniellae</i> Matile, 1972	B M S	<i>walkeri</i> Curtis, 1831	B M S
<i>denmax</i> Väisänen, 1979	B	<i>Megalopelma</i> Enderlein, 1911	
<i>dziedzickii</i> Väisänen, 1981	B M S	<i>nigroclavatum</i> (Strobl, 1910)	B
<i>egregia</i> (Dziedzicki, 1885)	B S	<i>Monoclona</i> Mik, 1886	
<i>flavicollis</i> (Zetterstedt, 1852)	B M	<i>braueri</i> (Strobl, 1895)	S
<i>griseovittata</i> (Zetterstedt, 1852)	S	<i>miki</i> Kertész, 1898	S
<i>hackmani</i> Väisänen, 1984	B	<i>orientalis</i> A. Zaitzev, 1983	M
<i>hebrardi</i> Väisänen & Matile, 1980	S	<i>rufilatera</i> (Walker, 1837)	B M
<i>hütsi</i> Väisänen, 1979	B M S	<i>silvatica</i> A. Zaitzev, 1983	M
<i>levis</i> (Dziedzicki, 1885)	M	<i>Neuratelia</i> Rondani, 1856	
= <i>pseudocinerascens</i> (Strobl, 1901)		<i>nemoralis</i> (Meigen, 1818)	B M S
<i>livida</i> (Dziedzicki, 1885)	B M	<i>Paratinia</i> Mik, 1874	
<i>marginata</i> (Meigen, 1818)	B M S	<i>sciarina</i> Mik, 1874	B M S
<i>neohyalinata</i> Väisänen, 1984	B M S	<i>Phthinia</i> Winnertz, 1863	
<i>nitida</i> (Zetterstedt, 1852)	B M	<i>humilis</i> Winnertz, 1863	B M S
= <i>exigua</i> (Winnertz, 1863)		<i>winnertzi</i> Mik, 1869	M S
<i>occultans</i> (Winnertz, 1863)	B M S	<i>Polylepta</i> Winnertz, 1863	
<i>ornata</i> (Meigen, 1818)	B M	<i>guttiventris</i> (Zetterstedt, 1852)	B M S
<i>parva</i> (Dziedzicki, 1885)	M S	<i>Sciophila</i> Meigen, 1818	
<i>prominens</i> (Lundström, 1913)	B M S	<i>dziedzickii</i> Edwards, 1925	B
<i>pseudoapicalis</i> Landrock, 1925	M	<i>fridolini</i> Stackelberg, 1943	B
<i>punctata</i> (Meigen, 1804)	B M	<i>hirta</i> Meigen, 1818	B S
<i>ruficollis</i> (Zetterstedt, 1852)	B M S	<i>interrupta</i> (Winnertz, 1863)	M
<i>shermani</i> Garrett, 1924	S	<i>lutea</i> Macquart, 1826	B M S
<i>siebecki</i> (Landrock, 1912)	M S	<i>quatriterga</i> Hutson, 1979	B M
<i>sigma</i> Johannsen, 1910	B M S	<i>rufa</i> Meigen, 1830	S
= <i>duplicata</i> Edwards, 1925		<i>thoracica</i> Staeger, 1840	M
<i>tenus</i> (Walker, 1856)	B M S	<i>Syntemna</i> Winnertz, 1863	
<i>trivittata</i> (Zetterstedt, 1838)	M	<i>morosa</i> Winnertz, 1863	B
<i>tonida</i> (Winnertz, 1863)	B M S	<i>relicta</i> (Lundström, 1912)	S
<i>vittiventris</i> (Zetterstedt, 1852)	B M S	<i>setigera</i> (Lundström, 1914)	S
<i>wankowiczii</i> (Dziedzicki, 1885)	B M S	GNORISTINI	
<i>winnertzi</i> (Dziedzicki, 1855)	B M S	<i>Apolephthisa</i> Grzegorzek, 1885	
sg. <i>Mycomyopsis</i> Väisänen, 1984		<i>subincana</i> (Curtis, 1837)	B S

<i>Boletina</i> Staeger, 1840			<i>referta</i> Plassmann, 1976	B
<i>anderschi</i> (Stannius, 1831)	M		<i>Greenomyia</i> Brunetti, 1912	
<i>basalis</i> (Meigen, 1818)	B M S		<i>borealis</i> (Winnertz, 1863)	M
<i>cincticornis</i> (Walker, 1848)	B M S		<i>Leia</i> Meigen, 1818	
<i>dispecta</i> Dziedzicki, 1885	S		<i>bilineata</i> (Winnertz, 1863)	M S
<i>dubia</i> (Meigen, 1804)	M		<i>bimaculata</i> (Meigen, 1804)	B M S
<i>gripha</i> Dziedzicki, 1885	B M S		<i>crucigera</i> Zetterstedt, 1838	M
<i>lundstroemi</i> Landrock, 1912	M		<i>cylindrica</i> (Winnertz, 1863)	M
<i>moravica</i> Landrock, 1912	M		<i>fascipennis</i> Meigen, 1818	B M S
<i>nigricans</i> Dziedzicki, 1885	M		<i>picta</i> Meigen, 1830	B M S
<i>nigricoxa</i> Staeger, 1840	B M		<i>piffardi</i> Edwards, 1925	M
<i>nitida</i> Grzegorzek, 1885	M		<i>subfasciata</i> (Meigen, 1818)	B M S
<i>oreadophila</i> Chandler, 1995	M		<i>winthemi</i> Lehmann, 1822	B M S
<i>plana</i> Walker, 1856	M S		<i>Megophthalmidia</i> Dziedzicki, 1889	
<i>rejecta</i> Edwards, 1941	S		<i>crassicornis</i> (Curtis, 1837)	M S
<i>sciarina</i> Staeger, 1840	B M S		<i>Rondaniella</i> Johannsen, 1909	
<i>trispinosa</i> Edwards, 1913	S		<i>dimidiata</i> (Meigen, 1804)	M
<i>trivittata</i> (Meigen, 1818)	B M S		<i>Tetragoneura</i> Winnertz, 1846	
<i>Coelophthiria</i> Edwards, 1941			<i>ambigua</i> , Grzegorzek, 1885	S
<i>thoracica</i> (Winnertz, 1863)	M		<i>sylvatica</i> (Curtis, 1837)	B M
<i>Coelosia</i> Winnertz, 1863				
<i>flava</i> (Staeger, 1840)	M		MYCETOPHILINAE	
<i>silvatica</i> Landrock, 1918	B M S		MYCETOPHILINI	
<i>truncata</i> Lundstroem, 1909	M		<i>Dynatosoma</i> Winnertz, 1863	
<i>Dziedzickia</i> Johannsen, 1909			<i>cochleare</i> Strobl, 1895	B M
<i>marginata</i> (Dziedzicki, 1885)	B		<i>fuscicorne</i> (Meigen, 1818)	B M S
<i>Gnoriste</i> Meigen, 1818			<i>majus</i> Landrock, 1912	M S
<i>apicalis</i> Meigen, 1818	M		<i>reciprocum</i> (Walker, 1848)	B M S
<i>bilineata</i> Zetterstedt, 1852	B M S		<i>thoracicum</i> (Zetterstedt, 1838)	B M
<i>Grzegorzekia</i> Edwards, 1941			= <i>rufithorax</i> Strobl, 1895	
<i>collaris</i> (Meigen, 1818)	B S		<i>Epicypta</i> Winnertz, 1863	
= <i>splendida</i> (Winnertz, 1863)			<i>aterrima</i> (Zetterstedt, 1852)	B
<i>Palaeodocosia</i> Meunier, 1904			<i>scatophora</i> (Perris, 1849)	S
<i>janickii</i> (Dziedzicki, 1923)	M		<i>Mycetophila</i> Meigen, 1803	
<i>Saigusaia</i> Vockeroth, 1980			<i>abbreviata</i> Landrock, 1914	M
<i>flaviventris</i> (Strobl, 1894)	M		<i>abiecta</i> (Laštovka, 1963)	B M
<i>Speolepta</i> Edwards, 1925			<i>adumbrata</i> Mik, 1884	M
<i>leptogaster</i> (Winnertz, 1863)	B M S		<i>alea</i> Laffoon, 1965	B M S
<i>Synapha</i> Meigen, 1818			<i>bialorussica</i> Dziedzicki, 1884	B
<i>fasciata</i> Meigen, 1818	B		<i>biusta</i> Meigen, 1818	B
<i>vitripennis</i> (Meigen, 1818)	M S		<i>blanda</i> Winnertz, 1863	M S
			<i>bohémica</i> (Laštovka, 1963)	B S
LEIINI			<i>brevitarsata</i> (Laštovka, 1963)	B S
<i>Clastobasis</i> Skuse, 1890			<i>caudata</i> Staeger, 1840	M
<i>alternans</i> (Winnertz, 1863)	M		<i>cingulum</i> Meigen, 1830	B M
= <i>unicolor</i> (Winnertz, 1863)			<i>confluens</i> Dziedzicki, 1884	M
<i>Docosia</i> Winnertz, 1863			<i>confusa</i> Dziedzicki, 1884	M S
<i>flavicoxa</i> Strobl, 1900	M		<i>curviseta</i> Lundström, 1911	M S
<i>fuscipes</i> (von Roser, 1840)	B M		<i>czizeki</i> Landrock, 1911	M
<i>gilvipes</i> (Walker, 1856)	B		<i>dentata</i> Lundström, 1913	M S
<i>moravica</i> Landrock, 1916	M		<i>dziedzickii</i> Chandler, 1977	B M S
<i>nigra</i> Landrock, 1928	B M S		<i>edwardsi</i> Lundström, 1913	M S
<i>pallipes</i> Edwards, 1941	S		<i>evanida</i> Laštovka, 1972	B S
<i>sciarina</i> (Meigen, 1830)	B M		<i>finlandica</i> Edwards, 1913	B M
<i>setosa</i> Landrock, 1916	M		<i>forcipata</i> Lundström, 1913	M
<i>Ectrepesthoneura</i> Enderlein, 1911			<i>formosa</i> Lundström, 1911	B M
<i>hirta</i> (Winnertz, 1846)	B M		<i>fraterna</i> Winnertz, 1863	M

<i>fungorum</i> (De Geer, 1776)	B M S	<i>cinerascens</i> Winnertz, 1863	M S
<i>gibbula</i> Edwards, 1925	B M S	<i>conformis</i> (Walker, 1865)	S
<i>gratiosa</i> Winnertz, 1863	M	<i>crassitarsus</i> Hackman, 1970	M
<i>hamata</i> Winnertz, 1863	M	<i>egregia</i> Dziedzicki, 1889	M
<i>hetschkoi</i> Landrock, 1918	M S	<i>electa</i> Dziedzicki, 1889	M
<i>ichneumonea</i> Say, 1823	B M S	<i>exigua</i> (Zetterstedt, 1852)	B M S
<i>idonea</i> Laštovka, 1972	B S	<i>flavipes</i> Winnertz, 1863	B M S
<i>immaculata</i> (Dziedzicki, 1884)	M	<i>forcipata</i> Winnertz, 1863	B M
<i>karpatica</i> Landrock, 1925	M	<i>forcipula</i> Winnertz, 1863	M S
<i>laeta</i> Walker, 1848	B	<i>humeralis</i> Winnertz, 1863	B S
<i>lamellata</i> Lundström, 1911	M S	<i>interstincta</i> Dziedzicki, 1889	M
<i>luciosa</i> Meigen, 1830	B M	<i>maculata</i> Dziedzicki, 1889	M S
<i>lunata</i> Meigen, 1804	B M S	<i>minuta</i> Landrock, 1928	M
<i>magnicauda</i> Strobl, 1895	M	<i>mutabilis</i> Dziedzicki, 1889	M
<i>marginata</i> Winnertz, 1863	B M S	<i>nigricornis</i> (Zetterstedt, 1852)	M S
<i>miki</i> Dziedzicki, 1884	M	<i>nitidiventris</i> (van der Wulp, 1858)	B M S
<i>montana</i> Landrock, 1925	M	<i>notata</i> Dziedzicki, 1889	S
<i>moravica</i> Landrock, 1925	M	<i>obtusa</i> Winnertz, 1863	M
<i>occultans</i> Lundström, 1913	M S	<i>peculiaris</i> Dziedzicki, 1889	M
<i>ocellus</i> Walker, 1848	B M S	<i>petulans</i> Dziedzicki, 1889	M S
<i>ornata</i> Stephens, 1829	B M S	<i>siebeckii</i> Dziedzicki, 1889	M
<i>pecinai</i> (Laštovka, 1963)	B S	<i>signata</i> Winnertz, 1863	M
<i>perpallida</i> Chandler, 1993	S	<i>strenua</i> Winnertz, 1863	B M S
<i>picula</i> Meigen, 1830	B M S	<i>sudetica</i> Dziedzicki, 1889	M
<i>pumila</i> Winnertz, 1863	B M S	<i>sylvatica</i> Dziedzicki, 1889	S
<i>rudis</i> Winnertz, 1863	M	<i>taczanowskyi</i> Dziedzicki, 1889	M
<i>ruficollis</i> Meigen, 1818	B S	<i>tenuis</i> Winnertz, 1863	B M S
<i>schnabli</i> (Dziedzicki, 1884)	B M	<i>tieffii</i> Dziedzicki, 1889	M
<i>sigillata</i> Dziedzicki, 1884	M S	<i>triangularis</i> Winnertz, 1863	M S
<i>signata</i> Meigen, 1830	B M S	<i>unica</i> Dziedzicki, 1889	M
<i>signatoides</i> Dziedzicki, 1884	B M S	<i>willistoni</i> Dziedzicki, 1889	M
<i>assimilis</i> Matile, 1967		Platurocypta Enderlein, 1910	
<i>sordida</i> van der Wulp, 1874	M S	<i>punctum</i> (Stannius, 1831)	M
<i>spectabilis</i> Winnertz, 1863	B M S	<i>testata</i> (Edwards, 1925)	B M S
<i>stolida</i> Walker, 1856	M S	Sceptonia Winnertz, 1863	
<i>strigata</i> Staeger, 1840	M	<i>concolor</i> Winnertz, 1863	M
<i>strigatoides</i> (Landrock, 1927)	S	<i>membranacea</i> Edwards, 1925	M S
<i>stroblii</i> Laštovka, 1972	B M S	<i>nigra</i> (Meigen, 1804)	B M
<i>stylata</i> (Dziedzicki, 1884)	M S	Trichonta Winnertz, 1863	
<i>stylatiformis</i> Landrock, 1925	M	<i>aberrans</i> Lundström, 1911	M
<i>sumavica</i> (Laštovka, 1963)	B S	<i>apicalis</i> Strobl, 1898	M S
<i>triangularis</i> Lundström, 1912	M	<i>atricauda</i> (Zetterstedt, 1852)	M S
<i>trinotata</i> Staeger, 1840	M S	<i>bezzii</i> Landrock, 1913	M S
<i>unguiculata</i> Lundström, 1913	M	<i>bicolor</i> Landrock, 1912	M
<i>unicolor</i> Stannius, 1831	M	<i>brevicauda</i> Lundström, 1906	S
<i>uninotata</i> Zetterstedt, 1852	B M S	<i>clavigera</i> Lundström, 1913	M S
<i>unipunctata</i> Meigen, 1818	B M S	<i>comis</i> Gagné, 1981	S
<i>v-nigrum</i> Lundström, 1913	M	<i>conjugens</i> Lundström, 1909	B M S
<i>vittipes</i> Zetterstedt, 1852	B S	<i>falcata</i> Lundström, 1911	M S
<i>xanthopyga</i> Winnertz, 1863	B M S	<i>flavicauda</i> Lundström, 1914	S
<i>zetterstedti</i> Lundström, 1906	B S	<i>fragilis</i> Gagné, 1981	S
Phronia Winnertz, 1863		<i>girschneri</i> Landrock, 1912	M S
<i>biarcuata</i> (Becker, 1908)	B M S	<i>hamata</i> Mik, 1880	B M S
<i>bicolor</i> Dziedzicki, 1889	M S	<i>melanura</i> (Staeger, 1840)	M
<i>fuscinervis</i> Van Duzee, 1928		<i>submaculata</i> (Staeger, 1840)	S
<i>braueri</i> Dziedzicki, 1889	M	<i>terminalis</i> (Walker, 1856)	M S
<i>caliginosa</i> Dziedzicki, 1889	M	<i>trivittata</i> Lundström, 1916	S

<i>venosa</i> (Staeger, 1840)	M S	<i>griseicollis</i> (Staeger, 1840)	B M
<i>vitta</i> (Meigen, 1830)	B M	<i>griseolum</i> (Zetterstedt, 1852)	S
<i>vulcani</i> (Dziedzicki, 1889)	M S	<i>luteum</i> (Landrock, 1925)	M
Zygomysia Winnertz, 1863		<i>proximum</i> (Staeger, 1840)	M
<i>humeralis</i> (Wiedemann, 1817)	M S	<i>radianum</i> (Lundström, 1911)	M
<i>notata</i> (Stannius, 1831)	B M S	<i>ruficorne</i> (Meigen, 1838)	M S
<i>pictipennis</i> (Staeger, 1840)	M	<i>sericoma</i> (Meigen, 1830)	B M S
<i>pseudohumeralis</i> Caspers, 1980	B	<i>spathulatum</i> (Lundström, 1911)	M
<i>semifusca</i> (Meigen, 1818)	M S	sg. Stigmatomeria Tuomikoski, 1966	
<i>valeriae</i> Chandler, 1991	S	<i>crassicornis</i> (Stannius, 1831)	B M S
<i>valida</i> Winnertz, 1863	B M	= <i>obscurum</i> (Winnertz, 1863)	
<i>vara</i> (Staeger, 1840)	B M	Cordyla Meigen, 1803	
EXECHIINI		<i>bicornuta</i> (Landrock, 1926)	M
Allodia Winnertz, 1863		<i>brevicornis</i> (Staeger, 1840)	M
sg. Allodia s. str.		<i>crassicornis</i> Meigen, 1818	B M S
<i>anglofennica</i> Edwards, 1921	B S	<i>fissa</i> Edwards, 1925	B M
<i>lugens</i> (Wiedemann, 1817)	B M S	<i>flaviceps</i> (Staeger, 1840)	M
<i>lundstroemi</i> Edwards 1921	B	<i>fusca</i> Meigen, 1804	B S
<i>ornaticollis</i> (Meigen, 1818)	B M S	<i>murina</i> Winnertz, 1863	B S
<i>pyxidiformis</i> Zaitzev, 1983	B	<i>nitens</i> Winnertz, 1863	M
sg. Brachycampta Winnertz, 1863		<i>nitidula</i> Edwards, 1925	B
<i>alternans</i> (Zetterstedt, 1838)	B M S	<i>parvipalpis</i> Edwards, 1925	M
<i>barbata</i> (Lundström, 1909)	M	<i>pusilla</i> Edwards, 1925	B M
<i>czernyi</i> (Landrock, 1912)	M	<i>semiflava</i> (Staeger, 1840)	M
<i>foliifera</i> (Strobl, 1910)	M	Exechia Winnertz, 1863	
<i>grata</i> (Meigen, 1830)	M	<i>bicincta</i> (Staeger, 1840)	B M
<i>pistillata</i> (Lundström, 1911)	M S	<i>chandleri</i> Caspers, 1987	S
<i>silvatica</i> (Landrock, 1912)	M	<i>cincta</i> Winnertz, 1863	B
<i>triangularis</i> (Strobl, 1895)	M	<i>confinis</i> Winnertz, 1863	B M
Allodiopsis Tuomikoski, 1966		<i>contaminata</i> Winnertz, 1863	B M
sg. Allodiopsis s. str.		<i>dizona</i> Edwards, 1924	B S
<i>domestica</i> (Meigen, 1830)	B M S	<i>dorsalis</i> (Staeger, 1840)	M S
<i>rustica</i> Edwards, 1941	B M S	<i>exigua</i> Lundström, 1909	M S
sg. Notolopha Tuomikoski, 1966		<i>fusca</i> (Meigen, 1804)	B M S
<i>cristata</i> (Staeger, 1840)	B M S	<i>maculipennis</i> (Stannius, 1831)	M
sg. Myrosia Tuomikoski, 1966		<i>nigroscutellata</i> Landrock, 1912	M
<i>maculosa</i> (Meigen, 1818)	B M	<i>papyracea</i> Stackelberg, 1948	S
Anatella Winnertz, 1863		<i>parva</i> Lundström, 1909	B M
<i>ciliata</i> Winnertz, 1863	B M S	<i>pseudocincta</i> Strobl, 1909	M S
<i>lenis</i> Dziedzicki, 1923	M S	<i>repanda</i> Johannsen, 1912	B
<i>longisetosa</i> Dziedzicki, 1923	M	<i>separata</i> Lundström, 1912	M S
<i>minuta</i> (Staeger, 1840)	M	<i>seriata</i> (Meigen, 1830)	M S
<i>novata</i> Dziedzicki, 1923	M	<i>spinuligera</i> Lundström, 1912	M S
<i>simpatia</i> Dziedzicki, 1923	B	Exechiopsis Tuomikoski, 1966	
Brachypeza Winnertz, 1863		sg. Exechiopsis s. str.	
sg. Brachypeza s. str.		<i>clypeata</i> (Lundström, 1911)	S
<i>armata</i> Winnertz, 1863	B S	<i>distendens</i> (Lackschewitz, 1937)	S
<i>bisignata</i> Winnertz, 1863	B	<i>dryaspagensis</i> Chandler, 1977	S
<i>radiata</i> Jenkinson, 1908	B	<i>dumitrescui</i> Burghel-Balacesco, 1972	S
sg. Paracordyla Tuomikoski, 1966		<i>fimbriata</i> (Lundström, 1909)	M S
<i>obscura</i> Winnertz, 1863	B	<i>furcata</i> (Lundström, 1911)	M S
Brevicornu Marshall, 1896		<i>hammi</i> (Edwards, 1925)	S
sg. Brevicornu s. str.		<i>indecisa</i> (Walker, 1856)	B M S
<i>fennicum</i> (Landrock, 1927)	M	<i>intersecta</i> (Meigen, 1818)	B M S
<i>fissicauda</i> (Lundström, 1911)	B	<i>januarii</i> (Lundström, 1913)	S
<i>fuscipenne</i> (Staeger, 1840)	M S	<i>lackschewitziana</i> (Stackelberg, 1948)	S
		<i>landrocki</i> (Lundström, 1912)	S

<i>ligulata</i> (Lundström, 1913)		S	<i>brevispina</i> Tuomikoski, 1960		M
<i>magnicauda</i> (Lundström, 1911)		M S	<i>brunnipes</i> (Meigen, 1804)		B M S
<i>olentica</i> (Burghele-Balacesco, 1965)		S	= <i>picipes</i> (Zetterstedt, 1838)		
<i>pseudindecis</i> Laštovka & Matile, 1974	B	S	<i>cinerascens</i> (Grzegorzek, 1884)		B
<i>pulchella</i> (Winnertz, 1863)		M	<i>desolata</i> Rudzinski, 1994		M
<i>sagittata</i> , Laštovka & Matile, 1974		S	<i>forficulata</i> (Bezzi, 1914)		not specified
<i>subulata</i> (Winnertz, 1863)	B M	S	<i>fungicola</i> (Winnertz, 1867)		M
<i>unguiculata</i> (Lundström, 1911)		S	<i>inuitata</i> Tuomikoski, 1960		S
<i>vizzavonensis</i> (Edwards, 1928)		S	<i>leptoptera</i> Tuomikoski, 1960		M
sg. <i>Xenexechia</i> Tuomikoski, 1966			<i>lobata</i> Hondru, 1968		M
<i>crucigera</i> (Lundström, 1909)	B		<i>lobulifera</i> Frey, 1948		not specified
<i>leptura</i> (Meigen, 1830)		M	<i>longicauda</i> Mohrig & Menzel, 1990		M
<i>pollicata</i> (Edwards, 1925)		S	<i>longistylia</i> Mohrig & Krivosheina, 1982		not specified
<i>Pseudexechia</i> Tuomikoski, 1966			<i>lutaria</i> (Winnertz, 1869)		B
<i>aurivernica</i> Chandler, 1978		S	<i>nervosa</i> (Meigen, 1818)		B M
<i>trisinata</i> (Edwards, 1913)		M S	<i>nitidicollis</i> (Meigen, 1818)		B M
<i>trivittata</i> (Staeger, 1840)		M	<i>nocturna</i> Tuomikoski, 1960		B M
<i>Pseudobrachypeza</i> Tuomikoski, 1966			<i>pallipes</i> (Fabricius, 1787)		B M
<i>helvetica</i> (Walker, 1856)		M S	<i>paranocturna</i> Rudzinski, 1994		M
<i>Pseudorymosia</i> Tuomikoski, 1966			<i>pauperata</i> (Winnertz, 1867)		B M S
<i>fovea</i> (Dziedzicki, 1910)		M S	<i>polonica</i> (Lengersdorf, 1929)		not specified
<i>optiva</i> (Dziedzicki, 1910)		M	<i>praecox</i> (Meigen, 1818)		B M S
<i>Rymosia</i> Winnertz, 1863			<i>pratricula</i> Tuomikoski, 1960		M
<i>affinis</i> Winnertz, 1863		M S	<i>reflexa</i> Tuomikoski, 1960		S
<i>bifida</i> Edwards, 1925	B		<i>rufescens</i> (Zetterstedt, 1852)		not specified
<i>connexa</i> Winnertz, 1863		M	<i>scabricornis</i> Tuomikoski, 1960		M S
<i>fasciata</i> (Meigen, 1804)		M S	<i>soudeki</i> Kratochvil, 1936		S
<i>fraudatrix</i> Dziedzicki, 1910		M S	<i>spinostyla</i> Mohrig & Menzel, 1990		M
<i>frenata</i> Dziedzicki, 1910		M	<i>strigata</i> (Staeger, 1840)		M
<i>lundstroemi</i> Dziedzicki, 1910		M	<i>submoesta</i> Mohrig & Krivosheina, 1989	B	
<i>placida</i> Winnertz, 1863		M	<i>subscabricornis</i> Mohrig & Menzel, 1990		M
<i>setiger</i> Dziedzicki, 1910		M S	<i>trivittata</i> (Staeger, 1840)		B M
<i>signatipes</i> (van der Wulp, 1859)		M	<i>vagans</i> (Winnertz, 1868)		B M
<i>spiniforceps</i> Matile, 1963	B	S	= <i>callicera</i> Frey, 1948		
<i>spinipes</i> Winnertz, 1863		S	<i>xenoreflexa</i> Mohrig & Barták, 1994		M
<i>virens</i> Dziedzicki, 1910		M S	<i>Camptochaeta</i> Hippa & Vilkamaa, 1994		
<i>Synplasta</i> Skuse, 1890			<i>ofenkaulis</i> (Lengersdorf, 1925)		M
<i>bayardi</i> Matile, 1971		S	<i>recurvispina</i> Freeman, 1987		M
<i>excogitata</i> (Dziedzicki, 1910)	B		<i>Chaetosciara</i> Frey, 1942		
<i>gracilis</i> (Winnertz, 1863)		M	<i>estlandica</i> (Lengersdorf, 1929)		B
<i>venosa</i> (Dziedzicki, 1910)		M	<i>Corynoptera</i> Winnertz, 1867		
<i>Tarnania</i> Tuomikoski, 1966			<i>blanda</i> (Winnertz, 1867)		B M
<i>dziedzickii</i> (Edwards, 1941)		S	<i>flavicauda</i> (Zetterstedt, 1855)		B
<i>fenestralis</i> (Meigen, 1818)	B M	S	<i>forcipata</i> (Winnertz, 1867)		M
<i>tarnanii</i> (Dziedzicki, 1910)		M S	<i>inexpectata</i> Tuomikoski, 1960		M
			<i>irmgardis</i> (Lengersdorf, 1930)		M
			<i>melanochaeta</i> Mohrig & Menzel, 1992		S
			<i>membranigera</i> (Kieffer, 1903)		S
			<i>minima</i> (Meigen, 1818)		M
			<i>montana</i> (Winnertz, 1869)	B	
			<i>nigrohalteralis</i> (Frey, 1948)	B	
			<i>perpusilla</i> Winnertz, 1867		M
			<i>postforcipata</i> Rudzinski, 1993		M
			<i>praeforcipata</i> Mohrig & Mamaev, 1937		M
			<i>saetistyla</i> Mohrig & Krivosheina, 1985		M
			<i>sphenoptera</i> Tuomikoski, 1960		M

SCIARIDAE

Hans-Georg Rudzinski & Vladimír Košel

Bradysia Winnertz, 1867

<i>affinis</i> (Zetterstedt, 1838)		M
<i>alpicola</i> (Winnertz, 1867)	B	S
<i>amoena</i> (Winnertz, 1867)	B M	S
<i>iriseriata</i> (Winnertz, 1867)		
<i>aprica</i> (Winnertz, 1867)		M S
<i>bicolor</i> (Meigen, 1818)	B M	S

<i>subflava</i> (Lengersdorf, 1941)	M	<i>mali</i> (Fitch, 1856)	B M
<i>vitella</i> Rudzinski & Drissner, 1992	M	= <i>solani</i> (Winnertz, 1871)	
<i>Cratyna</i> Winnertz, 1867		<i>micria</i> Mohrig & Menzel, 1990	S
sg. <i>Decembrina</i> Frey, 1942		<i>Mycosciara</i> Kieffer, 1903	
<i>ambigua</i> (Lengersdorf, 1934)	M S	<i>brevipalpis</i> Kieffer, 1903	S
<i>uliginosa</i> (Lengersdorf, 1929)	S	<i>Phytosciara</i> Frey, 1942	
sg. <i>Peyerimhoffia</i> Kieffer, 1903		sg. <i>Dolichosciara</i> Tuomikoski, 1960	
<i>vagabunda</i> (Winnertz, 1867)	B M	<i>flavipes</i> (Meigen, 1804)	B M S
= <i>brachyptera</i> (Kieffer, 1903)		sg. <i>Phytosciara</i> Frey, 1942	
sg. <i>Plastosciara</i> Berg, 1899		<i>halterata</i> (Lengersdorf, 1926)	M
<i>schineri</i> (Winnertz, 1867)	B	<i>macrotricha</i> (Lengersdorf, 1926)	S
sg. <i>Spathobdella</i> Frey, 1948		<i>oldenbergi</i> Mohrig & Menzel, 1994	S
<i>colei</i> (Freeman, 1990)	M	sg. <i>Prosciara</i> s. str.	
<i>falcifera</i> Lengersdorf, 1933	M S	<i>flexa</i> Rudzinski, 1996	S
<i>nobilis</i> (Winnertz, 1867)	M S	<i>prosciarioides</i> (Tuomikoski, 1960)	not specified
<i>perplexa</i> (Winnertz, 1867)	S	<i>ungulata</i> Winnertz, 1867	B S
= <i>socialis</i> (Winnertz, 1871)		<i>Scatopsiara</i> Edwards, 1927	
<i>Ctenosciara</i> Tuomikoski, 1960		<i>atomaria</i> (Zetterstedt, 1851)	B M
<i>hyalipennis</i> (Meigen, 1804)	B M	= <i>vivida</i> (Winnertz, 1867)	
<i>Epidapus</i> Haliday, 1851		<i>curvilinea</i> (Lengersdorf, 1934)	not specified
sg. <i>Epidapus</i> s. str.		<i>fluviatilis</i> (Lengersdorf, 1940)	not specified
<i>absoloni</i> Cizek, 1915	M	<i>multispina</i> (Bukowski & Lengersdorf, 1936)	not specified
<i>atomarius</i> (De Geer, 1778)	B M S	<i>nana</i> (Winnertz, 1871)	M
<i>gracilis</i> (Walker, 1848)	M	<i>tricuspidata</i> (Winnertz, 1867)	M
<i>schillei</i> (Börner, 1903)	M	<i>vitripennis</i> (Meigen, 1818)	B M S
= <i>titan</i> Frey, 1948		<i>Schwenckfeldina</i> Frey, 1942	
sg. <i>Lengersdorfia</i> Kratochvíl, 1936		<i>carbonaria</i> (Meigen, 1830)	B M S
<i>abscondita</i> (Vimmer, 1926)	M	<i>Sciara</i> Meigen, 1803	
<i>detrificola</i> (Kratochvíl, 1936)	M	<i>analisis</i> Schiner, 1864	B M S
sg. <i>Vimmeria</i> Kratochvíl, 1936		<i>flavimana</i> Zetterstedt, 1851	B
<i>microthorax</i> (Börner, 1903)	B M	<i>helvola</i> Winnertz, 1867	B
= <i>gracilicornis</i> (Lengersdorf, 1926)		<i>humeralis</i> Zetterstedt, 1851	S
<i>Leptosciarella</i> Tuomikoski, 1960		<i>laackschewitzi</i> (Lengersdorf, 1934)	B
<i>melanoma</i> (Mohrig & Menzel, 1990)	M	= <i>mendax</i> (Tuomikoski, 1960)	
<i>nudinervis</i> Tuomikoski, 1960	S	<i>militaris</i> Nowicki, 1868	B M S
<i>pilosa</i> (Staeger, 1840)	B S	<i>modesta</i> (Winnertz, 1867)	B
= <i>scutellata</i> (Staeger, 1840)		<i>ruficauda</i> Meigen, 1818	B
<i>rejecta</i> (Winnertz, 1867)	B S	<i>thomae</i> (Linnaeus, 1767)	B M S
<i>scutellata</i> (Staeger, 1840)	M	<i>Soudekia</i> Vimmer, 1928	
= <i>elegans</i> (Winnertz, 1867)		<i>soudeki</i> (Vimmer, 1926)	M
<i>trochanterata</i> (Zetterstedt, 1851)	M S	<i>Trichosia</i> Winnertz, 1867	
<i>yerburyi</i> (Freeman, 1983)	M	<i>acrotricha</i> Tuomikoski, 1960	not specified
<i>Lycoriella</i> Frey, 1942		<i>coarctata</i> (Winnertz, 1867)	M
sg. <i>Hemineurina</i> Frey, 1942		<i>edwardsi</i> (Lengersdorf, 1930)	S
<i>bruckii</i> (Winnertz, 1867)	M	<i>morio</i> (Fabricius, 1794)	B M S
<i>cochleata</i> (Rübsaamen, 1898)	M	= <i>caudata</i> (Walker, 1848)	
<i>inflata</i> (Winnertz, 1867)	not specified	<i>splendens</i> Winnertz, 1867	S
<i>nodulosa</i> (Mohrig & Krivosheina, 1985)	M	<i>Xylosciara</i> Tuomikoski, 1957	
<i>pabulum</i> (Frey, 1948)	M	<i>benulae</i> Tuomikoski, 1960	not specified
<i>secundaria</i> Mohrig & Menzel, 1990	M	<i>lignicola</i> (Winnertz, 1867)	S
<i>subbruckii</i> Mohrig & Hövemeyer, 1992	M	<i>misella</i> (Frey, 1948)	M
<i>venosa</i> (Staeger, 1840)	not specified	<i>separata</i> Rudzinski, 1996	S
sg. <i>Lycoriella</i> s. str.		<i>spectabilis</i> Rudzinski, 1992	S
<i>auripila</i> (Winnertz, 1867)	B	<i>steleocera</i> Tuomikoski, 1960	S
<i>cellaris</i> (Lengersdorf, 1934)	M	<i>Zygoneura</i> Meigen, 1830	
<i>felix</i> (Schmitz, 1919)	M	<i>sciarina</i> Meigen, 1830	S
<i>lundstroemi</i> (Frey, 1948)	M		

CECIDOMYIIDAE

Marcela Skuhravá

LESTREMIINAE

Campylomyza Meigen, 1818*flavipes* (Meigen, 1818)

B

Catocha Haliday, 1833*latipes* Haliday, 1833

B

Lestremia Macquart, 1826*leucophaea* (Meigen, 1818)

B

Micromya Rondani, 1840*lucorum* Rondani, 1840

B

Xylopriona Kieffer, 1913*atra* (Meigen, 1804)

B

PORRICONDYLINAE

Asynapta H. Loew, 1850*pectoralis* Winnertz, 1853

M

strobi (Kieffer, 1920)

B

thuraui Rübsaamen, 1893

B M

Camptomyia Kieffer, 1894*innotata* Kieffer, 1894

S

Dicerura Kieffer, 1898*iridis* (Kaltenbach, 1874)

M

Heteropeza Winnertz, 1846*pygmaea* Winnertz, 1846

B

Holoneurus Kieffer, 1895*marginatus* (De Meijere, 1901)

S

Parepidosis Kieffer, 1913*venusta* (Winnertz, 1853)

B

Porricondyla Rondani, 1840*neglecta* Mamaev, 1965

M

nigripennis (Meigen, 1830)

B

quadridens Spungis, 1981

S

Winnertzia Rondani, 1861*lugubris* (Winnertz, 1853)

B

nigripennis Kieffer, 1896

S

CECIDOMYIINAE

LASIOPTERIDI

LASIOPTERINI

Aplonyx Stefani-Perez, 1908*chenopodii* Stefani-Perez, 1908

M

Hybolasioptera Rübsaamen, 1915*cerealis* (Lindeman, 1881)

B M

Lasioptera Meigen, 1818*arundinis* Schiner, 1854

B M S

calamagrostidis Rübsaamen, 1893

B M S

carophila F. Löw, 1874

B M S

eryngii (Vallot, 1829)

B M S

hungarica Möhn, 1968

B M S

kosarzewskeella Marikovskij, 1958

B M

populnea Wachtl, 1883

B M S

rubi (Schrank, 1803)

B M S

tiliarum H. Mamaeva, 1964

B

Microlasioptera Skuhravá, 1981*flexuosa* (Winnertz, 1853)

M

Ozirkhincus Rondani, 1840*anthemidis* (Rübsaamen, 1915)

B M S

longicollis Rondani, 1840

B M S

millefolii (Wachtl, 1884)

B M S

tanacetii (Kieffer, 1889)

B M S

Stefaniella Kieffer, 1898*atriplicis* Kieffer, 1898

M

brevipalpis Kieffer, 1898

M

cecconii Kieffer, 1909

M

OLIGOTROPHINI

Amerhapha Rübsaamen, 1914*gracilis* Rübsaamen, 1914

M

Apiomyia Kieffer, 1912*bergenstammi* (Wachtl, 1882)

M S

Arnoldiola Strand, 1926*gemmae* (Rübsaamen, 1891)

B M S

libera (Kieffer, 1909)

B M S

quercus (Binnie, 1877)

M S

sambuci (Kieffer, 1901)

B M S

Bayeriola Gagné, 1991*= Bayeria* Rübsaamen, 1914 nec Fritsch, 1905*erysimi* (Rübsaamen, 1914)

B M S

salicariae (Kieffer, 1888)

B M S

thymicola (Kieffer, 1888)

B M S

Blastomyia Kieffer, 1913*origani* (Tavares, 1901)

S

Bremiola Rübsaamen, 1915*onobrychidis* (Bremi, 1847)

B M

Camptoneuromyia Felt, 1908*petioli* H. Mamaeva, 1964

B

Craneiobia Kieffer, 1913*corni* (Giraud, 1863)

B M S

Cystiphora Kieffer, 1892*leontodontis* (Bremi, 1847)

M

sanguinea (Bremi, 1847)

B M S

schmidtii (Rübsaamen, 1914)

M

sonchi (Bremi, 1847)

B M S

taraxaci (Kieffer, 1888)

B M S

Dasineura Rondani, 1840*abietiperda* (Henschel, 1880)

B M

acrophila (Winnertz, 1853)

B M S

acuminata (Rübsaamen, 1915)

B M S

affinis (Kieffer, 1886)

B M S

alopecuri (Reuter, 1895)

B M

alpestris (Kieffer, 1909)

B M S

alyssi (Kieffer, 1901)

M

angelicae Rübsaamen, 1915

B M

aparines (Kieffer, 1889)

B M S

armoraciae Vimmer, 1936

B M S

asperulae (F. Löw, 1875)

B M S

astragalorum (Kieffer, 1909)

B M

auritae Rübsaamen, 1915

B M S

axillaris Kieffer, 1896

B M

bayeri Rübsaamen, 1914

B M S

berberidis (Kieffer, 1909)

B M S

bistortae (Kieffer, 1909)

B M S

<i>brassicae</i> (Winnertz, 1853)	B M S	<i>populeti</i> (Rübsaamen, 1889)	B M S
<i>bupleuri</i> (Wachtl, 1883)	B M	<i>populnea</i> (Kieffer, 1909)	M
<i>campanulae</i> Rübsaamen, 1914	B M	<i>potentillae</i> (Wachtl, 1885)	B M S
<i>capsulae</i> Kieffer, 1901	B M S	<i>praticola</i> (Kieffer, 1892)	B M S
<i>cardaminis</i> (Winnertz, 1853)	B M S	<i>prunicola</i> (F. Löw, 1889)	B S
<i>ceconiana</i> (Kieffer, 1909)	M	<i>pteridicola</i> (Kieffer, 1901)	B M S
<i>centaureae</i> (Kieffer, 1909)	M S	<i>pulsatillae</i> (Kieffer, 1894)	B S
<i>cerastii</i> (Bremi, 1877)	B M	<i>pustulans</i> (Rübsaamen, 1889)	B M S
<i>clausilia</i> (Bremi, 1847)	B M S	<i>pyri</i> (Bouché, 1847)	B M S
<i>clematidina</i> (Kieffer, 1913)	S	<i>ranunculi</i> (Bremi, 1847)	B M S
<i>comosae</i> Rübsaamen, 1915	B	<i>rossi</i> (Rübsaamen, 1914)	B
<i>corniculata</i> (Kieffer, 1909)	B M	<i>rubella</i> Kieffer, 1896	B M S
<i>crataegi</i> (Winnertz, 1853)	B M S	<i>ruebsaameni</i> (Kieffer, 1909)	B M S
<i>dactylidis</i> Metcalf, 1933	M	<i>salviae</i> (Kieffer, 1909)	B M S
<i>daphnes</i> (Kieffer, 1901)	B M	<i>sampaina</i> (Tavares, 1902)	M
<i>dioicae</i> (Rübsaamen, 1895)	B M	<i>sanguisorbae</i> (Rübsaamen, 1890)	B M S
<i>engstfeldi</i> (Rübsaamen, 1889)	B S	<i>saxifragae</i> (Kieffer, 1891)	B M
<i>epilobii</i> (F. Löw, 1889)	B M S	<i>serotina</i> (Winnertz, 1853)	B M S
<i>erigerontis</i> Rübsaamen, 1912	M	<i>schulzei</i> Rübsaamen, 1917	M S
<i>excavans</i> (Kieffer, 1909)	B M S	<i>silvestris</i> (Kieffer, 1909)	B M
<i>fäurmairei</i> (Kieffer, 1896)	B M	<i>silvicola</i> (Kieffer, 1909)	B
<i>festucae</i> Barnes, 1939	M	<i>similis</i> (F. Löw, 1888)	B M
<i>filicina</i> (Kieffer, 1889)	B M S	<i>sisymbrii</i> (Schränk, 1803)	B M S
<i>filipendulae</i> (Kieffer, 1909)	M	<i>spadicea</i> Rübsaamen, 1917	B M
<i>foliumcrispans</i> (Rübsaamen, 1896)	M	<i>stellariae</i> Rübsaamen, 1915	M
<i>fraxinea</i> Kieffer, 1907	B M S	<i>strumosa</i> (Bremi, 1847)	B M S
<i>fraxini</i> (Bremi, 1847)	B M S	<i>subpatula</i> (Bremi, 1847)	B M
<i>fractum</i> (Rübsaamen, 1895)	M	<i>symphyti</i> (Rübsaamen, 1891)	B M S
<i>galiicola</i> (F. Löw, 1880)	B M S	<i>szepligetii</i> (Kieffer, 1909)	M
<i>genistarum</i> (Kieffer, 1909)	S	<i>tetensi</i> (Rübsaamen, 1891)	B M S
<i>gentheri</i> Pritchard, 1953	B M S	<i>tetrahit</i> (Kieffer, 1909)	B M
<i>geranii</i> (Kieffer, 1907)	B M S	<i>teucarii</i> (Tavares, 1903)	S
<i>glechomae</i> (Kieffer, 1889)	B M S	<i>thomasiana</i> (Kieffer, 1888)	B M S
<i>glyciphylli</i> Rübsaamen, 1912	B M S	<i>tiliae</i> (Schränk, 1803)	B M S
<i>harrisoni</i> Bagnall, 1922	B S	<i>tortilis</i> (Bremi, 1847)	B M S
<i>hygrophila</i> (Mik, 1883)	B M S	<i>tortrix</i> (F. Löw, 1877)	B M S
<i>hyperici</i> (Bremi, 1847)	B M S	<i>trali</i> (Kieffer, 1909)	B
<i>irregularis</i> (Bremi, 1847)	B M S	<i>trifolii</i> (F. Löw, 1874)	B M S
<i>kellneri</i> (Henschel, 1875)	B M S	<i>tubularis</i> (Kieffer, 1909)	S
<i>kiefferiana</i> (Rübsaamen, 1891)	B M S	<i>tympani</i> (Kieffer, 1909)	B M S
<i>lamii</i> (Kieffer, 1909)	B M S	<i>ulmaria</i> (Bremi, 1847)	B M S
<i>lamicola</i> (Mik, 1888)	B M	<i>urticae</i> (Perris, 1840)	B M S
<i>lathyri</i> (Kieffer, 1909)	B M S	<i>viciae</i> (Kieffer, 1888)	B M S
<i>lathyricola</i> (Rübsaamen, 1890)	B M S	<i>violae</i> (F. Löw, 1880)	B M S
<i>legumicola</i> (Lintner, 1879)	B M S	<i>virgaeaureae</i> (Liebel, 1889)	B M S
<i>lithospermi</i> (H. Loew, 1850)	M S	<i>vitisidaea</i> (Kieffer, 1909)	B S
<i>loewiana</i> Rübsaamen, 1917	B M	<i>xylostei</i> (Kieffer, 1909)	S
<i>loewii</i> (Mik, 1882)	B M	<i>Didymomyia</i> Rübsaamen, 1912	
<i>lotharingiae</i> (Kieffer, 1888)	B M	<i>tiliacea</i> (Bremi, 1847)	B M S
<i>lupulinae</i> (Kieffer, 1891)	B M S	<i>Dryomyia</i> Kieffer, 1898	
<i>mali</i> (Kieffer, 1904)	B M S	<i>circinans</i> (Giraud, 1861)	M S
<i>medicaginis</i> (Bremi, 1847)	B M S	<i>Geocrypta</i> Kieffer, 1913	
<i>myosotidis</i> (Kieffer, 1902)	B M S	<i>braueri</i> (Handlirsch, 1884)	B M
<i>papaveris</i> (Winnertz, 1853)	B M S	<i>galii</i> (H. Loew, 1850)	B M S
<i>periclymeni</i> (Rübsaamen, 1889)	M S	<i>trachelii</i> (Wachtl, 1885)	B M S
<i>phyteumatis</i> (F. Löw, 1885)	B M S	<i>Gephyraululus</i> Rübsaamen, 1915	
<i>plicatrix</i> (H. Loew, 1850)	B M S	<i>raphanistri</i> (Kieffer, 1886)	B M S

<i>Giraudiella</i> Rübsaamen, 1915				<i>lamii</i> Rübsaamen, 1915	B	M	S
<i>inclusa</i> (Frauenfeld, 1862)	B	M	S	<i>laserpitii</i> Rübsaamen, 1917	B		S
<i>Harrisomyia</i> Skuhravá, 1986				<i>loniceræ</i> Rübsaamen, 1912	B	M	
<i>vitrina</i> (Kieffer, 1909)	B	M	S	<i>luceti</i> Kieffer, 1898	B	M	S
<i>Hartigola</i> Rübsaamen, 1912				<i>marteli</i> Kieffer, 1892	B		
<i>annulipes</i> (Hartig, 1839)	B	M	S	<i>orobi</i> (F. Löw, 1877)	B	M	S
<i>Iteomyia</i> Kieffer, 1913				<i>pavida</i> (Winnertz, 1853)	B		
<i>capreae</i> (Winnertz, 1853)	B	M	S	<i>pilosellæ</i> (Binnie, 1877)	B	M	S
<i>Jaapiella</i> Rübsaamen, 1915				<i>podagrariæ</i> Stelter, 1962	B	M	S
<i>alpina</i> (F. Löw, 1885)			S	<i>ruebsaameni</i> Hedicke, 1938	B	M	S
<i>bryoniae</i> (Bouché, 1847)	B	M	S	<i>saliceti</i> (H. Loew, 1850)	B		
<i>cirsicola</i> (Rübsaamen, 1915)	B	M	S	<i>stellariæ</i> (Liebel, 1889)	B	M	S
<i>clethrophila</i> Rübsaamen, 1917	B	M	S	<i>Mayetiola</i> Kieffer, 1896			
<i>compositarum</i> (Kieffer, 1888)	B			<i>avenae</i> (Marchal, 1895)		M	
<i>cucubali</i> (Kieffer, 1909)		M	S	<i>baudysi</i> Ertel, 1975	B		
<i>dittrichi</i> (Rübsaamen, 1895)			S	<i>bromi</i> (Hammerschmidt, 1834)			S
<i>floriperda</i> (F. Löw, 1888)	B	M	S	<i>bromicola</i> Roberti, 1953		M	
<i>genistamtorquens</i> (Kieffer, 1888)		M	S	<i>dactylidis</i> Kieffer, 1896	B	M	S
<i>genisticola</i> (F. Löw, 1877)	B	M	S	<i>destructor</i> (Say, 1817)	B	M	
<i>hedickei</i> Rübsaamen, 1921	B	M	S	<i>hellwigi</i> (Rübsaamen, 1912)	B	M	S
<i>jaaptana</i> (Rübsaamen, 1914)	B	M	S	<i>holci</i> Kieffer, 1896	B	M	
<i>knautiae</i> Rübsaamen, 1917	B	M	S	<i>lanceolatae</i> (Rübsaamen, 1895)	B	M	
<i>loticola</i> (Rübsaamen, 1889)	B	M	S	<i>molinae</i> (Rübsaamen, 1895)		M	
<i>medicaginis</i> (Rübsaamen, 1912)	B	M	S	<i>phalaris</i> Barnes, 1928		M	
<i>moraviae</i> (Wachtl, 1883)		M		<i>poae</i> (Bosc, 1817)	B	M	S
<i>rubicundula</i> (Rübsaamen, 1891)	B	M	S	<i>Mikiola</i> Kieffer, 1896			
<i>scabiosae</i> (Kieffer, 1888)	B	M	S	<i>fagi</i> (Hartig, 1839)	B	M	S
<i>schmidtii</i> (Rübsaamen, 1912)	B	M	S	<i>Mikomya</i> Kieffer, 1912			
<i>thalictri</i> (Rübsaamen, 1895)			S	<i>coryli</i> (Kieffer, 1901)	B	M	S
<i>vaccinorum</i> (Kieffer, 1895)	B	M		<i>Neomikiella</i> Hedicke, 1938			
<i>veronicae</i> (Vallot, 1827)	B	M	S	<i>beckiana</i> (Mik, 1885)	B	M	S
<i>viscaria</i> (Kieffer, 1886)		M		<i>lychnidis</i> (Heyden, 1861)		M	S
<i>volvens</i> Rübsaamen, 1917	B	M	S	<i>Oligotrophus</i> Latreille, 1805			
<i>Janetia</i> Kieffer, 1896				<i>juniperinus</i> (Linnaeus, 1758)	B	M	S
<i>cerris</i> (Kollar, 1850)	B	M	S	<i>panteli</i> Kieffer, 1898	B	M	
<i>homocera</i> (F. Löw, 1877)		M	S	<i>schmidtii</i> Rübsaamen, 1914		M	
<i>nervicola</i> Kieffer, 1909	B	M	S	<i>szepligetii</i> Kieffer, 1909			S
<i>pustularis</i> (Kieffer, 1909)	B	M	S	<i>Pemphigocecis</i> Rübsaamen, 1915			
<i>szepligetii</i> (Kieffer, 1896)	B		S	<i>ventricola</i> (Rübsaamen, 1899)		M	
<i>Janetiella</i> Kieffer, 1898				<i>Phegomyia</i> Kieffer, 1913			
<i>lemeei</i> (Kieffer, 1904)	B	M	S	<i>fagicola</i> (Kieffer, 1901)	B	M	S
<i>stskyoi</i> Felt, 1917	B		S	<i>Physemocecis</i> Rübsaamen, 1914			
<i>thymi</i> (Kieffer, 1888)	B	M	S	<i>hartigi</i> (Liebel, 1892)	B	M	S
<i>Kaltenbachiola</i> Hedicke, 1938				<i>ulmi</i> (Kieffer, 1909)	B	M	S
<i>strobi</i> (Winnertz, 1853)	B	M	S	<i>Rabdophaga</i> Westwood, 1847			
<i>Lathyromyza</i> Rübsaamen, 1915				<i>albipennis</i> (H. Loew, 1850)	B	M	S
<i>florum</i> Rübsaamen, 1915		M		<i>clavifex</i> (Kieffer, 1891)	B	M	S
<i>schlechtendali</i> (Kieffer, 1886)	B	M		<i>dubiosa</i> (Kieffer, 1913)	B	M	S
<i>Macrolabis</i> Kieffer, 1892				<i>gemmicola</i> (Kieffer, 1896)	B	M	
<i>alnicola</i> Rübsaamen, 1914			S	<i>heterobia</i> (H. Loew, 1850)	B	M	S
<i>aquilegiae</i> (Kieffer, 1909)			S	<i>insignis</i> (Kieffer, 1906)	B		
<i>cirsii</i> (Rübsaamen, 1890)	B			<i>iteobia</i> (Kieffer, 1890)	B	M	S
<i>heraclei</i> (Kaltenbach, 1862)	B	M	S	<i>karschi</i> (Kieffer, 1892)	B	M	
<i>hieracii</i> Rübsaamen, 1917	B	M	S	<i>marginemtorquens</i> (Bremi, 1847)	B	M	S
<i>holosteae</i> Rübsaamen, 1917	B	M	S	<i>nielsenii</i> (Kieffer & J. C. Nielsen, 1906)		M	
<i>incolens</i> Rübsaamen, 1895	B			<i>pierreana</i> (Kieffer, 1909)	B	M	S
<i>jaapi</i> Rübsaamen, 1915	B	M		<i>pierrei</i> (Kieffer, 1896)	B	M	S

<i>pseudococcus</i> (Thomas, 1890)	B	<i>riparia</i> (Winnertz, 1853)	B M S
<i>pulvini</i> (Kieffer, 1891)	B M	<i>rosarum</i> (Hardy, 1850)	B M S
<i>purpureaperda</i> (Barnes, 1935)	M	<i>stachydis</i> (Bremi, 1847)	B M S
<i>ramicola</i> (Rübsaamen, 1915)	B M	Zygiobia Kieffer, 1913	
<i>repentis</i> (Skuhravá, 1986)	B M S	<i>carpini</i> (F. Löw, 1874)	B M S
<i>rosaria</i> (H. Loew, 1850)	B M S		
<i>saliciperda</i> (Dufour, 1841)	B M S	STOMATOSEMIDI	
<i>salicis</i> (Schränk, 1803)	B M S	<i>Stomatosema</i> Kieffer, 1913	
<i>schwangeri</i> (Rübsaamen, 1915)	B M S	<i>obscura</i> (Mamaev, 1967)	S
<i>terminalis</i> (H. Loew, 1850)	B M S		
<i>triandraperda</i> (Barnes, 1935)	B	ASPHONDYLIIDI	
Rhizomyia Kieffer, 1898		<i>Asphondylia</i> H. Loew, 1850	
<i>ornata</i> Mamaev, 1967	B	<i>baudysi</i> Vimmer, 1937	B M S
Rhopalomyia Rübsaamen, 1892		<i>cytisi</i> Frauenfeld, 1873	B M S
<i>artemisiae</i> (Bouché, 1834)	B M S	<i>dorycnii</i> (Müller, 1870)	M S
<i>baccarum</i> (Wachtl, 1883)	B M S	<i>echii</i> (H. Loew, 1850)	B M S
<i>baudysi</i> Vimmer, 1928	M S	<i>ervi</i> Rübsaamen, 1896	M S
<i>cristaegalli</i> (Karsch, 1877)	B	<i>genistae</i> H. Loew, 1850	B M S
<i>flavipalpis</i> Vimmer, 1928	M	<i>hornigi</i> Wachtl, 1880	B
<i>florum</i> (Kieffer, 1890)	B	<i>lathyri</i> Rübsaamen, 1914	B M S
<i>foliorum</i> (H. Loew, 1850)	B M S	<i>lupulinae</i> Kieffer, 1909	B S
<i>hypogaea</i> (F. Löw, 1885)	M S	<i>massalongoi</i> Rübsaamen, 1893	S
<i>luetkemuellei</i> Thomas, 1893	M	<i>melanopus</i> Kieffer, 1890	B M S
<i>millefolii</i> (H. Loew, 1850)	B M S	<i>menthae</i> Kieffer, 1902	B M S
<i>pseudofoliorum</i> Vimmer, 1924	M	<i>miki</i> Wachtl, 1880	B M S
<i>ptarmicae</i> (Vallot, 1849)	B M	<i>ononidis</i> F. Löw, 1873	B M S
<i>simulans</i> Vimmer, 1924	M	<i>pruniperda</i> Rondani, 1867	B M S
<i>syngenesiae</i> (H. Loew, 1850)	B S	<i>sarothamni</i> H. Loew, 1850	B M S
<i>tanaceticola</i> (Karsch, 1879)	B M S	<i>scrophulariae</i> Schiner, 1856	M
<i>tubifex</i> (Bouché, 1847)	B M S	<i>serpylli</i> Kieffer, 1898	B M
Rondaniola Rübsaamen & Hedicke, 1938		<i>stachydis</i> Stelter, 1965	S
<i>bursaria</i> (Bremi, 1847)	B M S	<i>verbasci</i> (Vallot, 1827)	B M S
Sackenomyia Felt, 1908		Kiefferia Mik, 1895	
<i>recaumurii</i> (Bremi, 1847)	B M S	<i>pericarpiicola</i> (Bremi, 1847)	B M S
Semudobia Kieffer, 1913		Placochela Rübsaamen, 1916	
<i>betulae</i> (Winnertz, 1853)	B M S	<i>ligustri</i> (Rübsaamen, 1899)	B M S
<i>skuhravae</i> Roskam, 1977	B S	<i>nigripes</i> (F. Löw, 1877)	B M S
<i>tarda</i> Roskani, 1977	B	Polystepha Kieffer, 1897	
Schmidtella Rübsaamen, 1914		<i>malpighii</i> (Kieffer, 1909)	B M S
<i>gemmarum</i> Rübsaamen, 1914	B M S	<i>quercus</i> Kieffer, 1897	B S
Schueziella Möm, 1961		Schizomyia Kieffer, 1889	
<i>fagicola</i> (Barnes, 1939)	S	<i>galiorum</i> Kieffer, 1889	B M S
Spurgia Gagné, 1990			
<i>capitigena</i> (Bremi, 1847)	B M S	CECIDOMYIID	
<i>esulae</i> Gagné, 1990	B M	Acaroletes Kieffer, 1912	
Taxomyia Rübsaamen, 1912		<i>tetranychorum</i> Kieffer, 1913	M
<i>taxi</i> (Inchbald, 1861)	B M S	Acodiplosis Kieffer, 1895	
Trotteria Kieffer, 1901		<i>inulae</i> (H. Loew, 1847)	B M S
<i>galii</i> Rübsaamen, 1912	B M S	Ametrodiplosis Rübsaamen, 1910	
<i>ligustri</i> Barnes, 1954	S	<i>rudimentalis</i> (Kieffer, 1901)	B M
<i>obtusa</i> (H. Loew, 1850)	B M S	<i>thalicticola</i> (Rübsaamen, 1895)	B M S
<i>umbelliferarum</i> Kieffer, 1901	B M	Anabremia Kieffer, 1912	
Wachtliella Rübsaamen, 1915		<i>bellevoyei</i> (Kieffer, 1896)	B M S
<i>dalmatica</i> Rübsaamen, 1915	S	<i>medicaginis</i> Rübsaamen, 1917	M
<i>ericina</i> (F. Löw, 1885)	B	<i>trotteri</i> (Kieffer, 1909)	M
<i>niebleri</i> Rübsaamen, 1915	B M S	<i>viciae</i> Kieffer, 1913	B M
<i>persicariae</i> (Linnaeus, 1767)	B M S		

<i>Anisostephus</i> Rübsaamen, 1917			<i>festucae</i> Jones, 1940	B M
<i>betulinum</i> (Kieffer, 1889)	B M S		<i>floriperda</i> Rübsaamen, 1917	B M
<i>Antichiridium</i> Rübsaamen, 1911			<i>florum</i> Rübsaamen, 1917	M
<i>striatum</i> (Rübsaamen, 1911)	B M		<i>gei</i> Kieffer, 1909	B M S
<i>Aphidoletes</i> Kieffer, 1904			<i>helianthemi</i> (Hardy, 1850)	M S
<i>abietis</i> (Kieffer, 1896)	B		<i>heraclei</i> (Rübsaamen, 1889)	B
<i>aphidimyza</i> (Rondani, 1847)	B M S		<i>hyperici</i> Barnes, 1952	B
<i>thompsoni</i> Möhn, 1954	B S		<i>hypocheridis</i> (Rübsaamen, 1891)	B M S
<i>urticae</i> (Kieffer, 1895)	B S		<i>inquilina</i> Rübsaamen, 1917	B M
<i>Arthrocnodax</i> Rübsaamen, 1895			<i>inulicola</i> Stelter, 1965	S
<i>coryligallarum</i> (Targioni-Tozzetti, 1886)	B S		<i>jaapi</i> Rübsaamen, 1914	B M
<i>peregrina</i> (Winnertz, 1853)	B M		<i>jacobaeae</i> (H. Loew, 1850)	B M
<i>Aschistonyx</i> Rübsaamen, 1917			<i>lamiicola</i> Rübsaamen, 1915	B M S
<i>carpinicolus</i> Rübsaamen, 1917	B M S		<i>lepidii</i> Kieffer, 1909	M
<i>Atrichosema</i> Kieffer, 1904			<i>lentis</i> Aczél, 1942	M S
<i>aceris</i> Kieffer, 1904	M		<i>lilii</i> Kieffer, 1909	B M
<i>Blastodiplosis</i> Kieffer, 1912			<i>loti</i> (De Geer, 1776)	B M S
<i>artemisiae</i> (Kieffer, 1901)	B M		<i>lysimaehiae</i> (Rübsaamen, 1893)	B M S
<i>Brachydiplosis</i> Rübsaamen, 1910			<i>marchali</i> Kieffer, 1896	B M
<i>caricum</i> Rübsaamen, 1910	B M		<i>martagonis</i> Kieffer, 1909	S
<i>Bremia</i> Rondani, 1860			<i>medicaginis</i> Kieffer, 1895	B M S
<i>bifurcata</i> Kieffer, 1904		S	<i>melanocera</i> Kieffer, 1904	B M S
<i>Buhriella</i> Stelter, 1960			<i>merceri</i> Barnes, 1930	B M
<i>rubicola</i> Stelter, 1960	B		<i>molluginis</i> (Rübsaamen, 1889)	B M S
<i>Cecidomyia</i> Meigen, 1803			<i>nasturtii</i> (Kieffer, 1888)	B M S
<i>pini</i> (De Geer, 1782)	B		<i>nicolayi</i> (Rübsaamen, 1895)	B M
<i>Clinodiplosis</i> Kieffer, 1894			<i>onobrychidis</i> Kieffer, 1895	B M S
<i>bonularia</i> (Winnertz, 1853)	B		<i>ononidis</i> Kieffer, 1899	B
<i>citharus</i> (Kieffer, 1889)	B M S		<i>pastinacae</i> (Rübsaamen, 1891)	B M
<i>cirsii</i> Kieffer, 1909	B M		<i>petioli</i> (Kieffer, 1898)	B M S
<i>Contarinia</i> Rondani, 1860			<i>pilosellae</i> Kieffer, 1896	B M S
<i>acerplicans</i> (Kieffer, 1889)	B M S		<i>pisi</i> (Winnertz, 1854)	B M S
<i>acetosellae</i> (Rübsaamen, 1891)	B M		<i>poae</i> Barnes, 1946	B
<i>aconitifloris</i> Stelter, 1962	B M S		<i>populi</i> (Rübsaamen, 1917)	B M S
<i>acrocecis</i> Stelter, 1962		S	<i>pruniflorum</i> Coutin & Rambier, 1953	B M S
<i>aequalis</i> Kieffer, 1898	B M S		<i>pulchripes</i> (Kieffer, 1890)	B M S
<i>anthobia</i> (F. Löw, 1877)	B M S		<i>pyrivora</i> (Riley, 1886)	B M S
<i>anthophthora</i> (F. Löw, 1880)	B M S		<i>quercicola</i> (Rübsaamen, 1899)	M S
<i>arrhenatheri</i> Kieffer, 1901	B M		<i>quercina</i> (Rübsaamen, 1890)	B M S
<i>artemisiae</i> Rübsaamen, 1917	B S		<i>quinquenotata</i> (F. Löw, 1888)	B M S
<i>asclepiadis</i> (Giraud, 1863)		M	<i>rhamni</i> (Rübsaamen, 1892)	M
<i>baeri</i> (Prell, 1931)	B M S		<i>ribis</i> Kieffer, 1909	B M
<i>ballotae</i> Kieffer, 1898	B M S		<i>rubicola</i> Kieffer, 1909	M S
<i>barbichi</i> (Kieffer, 1890)	B M S		<i>rumicis</i> (H. Loew, 1850)	B M S
<i>brizae</i> Kieffer, 1896	B M S		<i>sambuci</i> (Kaltenbach, 1873)	B M S
<i>campanulae</i> (Kieffer, 1895)	B M S		<i>scoparii</i> (Rübsaamen, 1889)	B
<i>carpini</i> Kieffer, 1897	B M S		<i>scrophulariae</i> Kieffer, 1896	M S
<i>cerriperda</i> Skuhravá, 1991		S	<i>scutati</i> Rübsaamen, 1910	B
<i>chrysanthemi</i> (Kieffer, 1895)	B M		<i>silvestris</i> Kieffer, 1897	B M
<i>coryli</i> (Kaltenbach, 1859)	B M S		<i>solani</i> (Rübsaamen, 1891)	B M S
<i>colini</i> Kieffer, 1901	B		<i>sorbi</i> Kieffer, 1896	B M S
<i>cracca</i> Kieffer, 1897	B M S		<i>steini</i> (Karsch, 1881)	B M S
<i>crispans</i> Kieffer, 1909		M S	<i>subulifex</i> Kieffer, 1897	M S
<i>cybelae</i> Gagné, 1972	B M S		<i>tanacetii</i> Rübsaamen, 1921	B M
<i>dactylidis</i> (H. Loew, 1851)		M	<i>tiliarum</i> (Kieffer, 1890)	B M S
<i>echii</i> (Kieffer, 1895)		M	<i>tragopogonis</i> Kieffer, 1909	B M S
<i>fagi</i> Rübsaamen, 1921	B M S		<i>tremulae</i> Kieffer, 1909	B M S

<i>tritici</i> (Kirby, 1798)	B M S	<i>Paradiplosis</i> Felt, 1908	
<i>valerianae</i> (Rübsaamen, 1890)	B M S	<i>abietis</i> (Hubault, 1945)	B M S
<i>viburnorum</i> Kieffer, 1913	M	<i>Parallelodiplosis</i> Rübsaamen, 1910	
<i>vincetoxici</i> Kieffer, 1909	M S	<i>bupleuri</i> (Rübsaamen, 1895)	B
<i>viticola</i> Rübsaamen, 1906	S	<i>galliperda</i> (F. Löw, 1889)	B M S
<i>Coquillettomyia</i> Felt, 1908		<i>Planetella</i> Westwood, 1840	
<i>caricis</i> (Mölin, 1955)	S	<i>arenariae</i> (Rübsaamen, 1899)	B M
<i>Dichodiplosis</i> Rübsaamen, 1910		<i>billoti</i> (Kieffer, 1909)	B M
<i>langeni</i> Rübsaamen, 1910	B M S	<i>caricis</i> (Rübsaamen, 1911)	B M
<i>Diodaulus</i> Rübsaamen, 1917		<i>cornifex</i> (Kieffer, 1898)	B M
<i>linariae</i> (Winnertz, 1853)	B M S	<i>fischeri</i> (Frauenfeld, 1867)	B M
<i>traili</i> (Kieffer, 1889)	B M S	<i>frireni</i> (Kieffer, 1909)	B M S
<i>Drisina</i> Giard, 1893		<i>gallarum</i> (Rübsaamen, 1899)	B M
<i>glutinosa</i> Giard, 1893	B M S	<i>granifex</i> (Kieffer, 1898)	B M S
<i>Endopsylla</i> Meijere, 1907		<i>kneuckeri</i> Kieffer, 1909	B M
<i>agilis</i> Meijere, 1907	M	<i>rosenhaueri</i> (Rübsaamen, 1892)	B
<i>Feltiella</i> Rübsaamen, 1911		<i>subterranea</i> (Kieffer & Trotter, 1904)	B
<i>tetranychii</i> Rübsaamen, 1911	B	<i>tuberifica</i> (Rübsaamen, 1899)	B M S
<i>Hadrobremia</i> Kieffer, 1912		<i>tumorifica</i> (Rübsaamen, 1899)	B M
<i>ononidis</i> (Kieffer, 1898)	S	<i>Plemeliella</i> Seitner, 1908	
<i>Haplodiplosis</i> Rübsaamen, 1910		<i>abietina</i> Seitner, 1908	B M
<i>marginata</i> (von Roser, 1840)	B M S	<i>betulicola</i> (Kieffer, 1889)	B M S
= <i>equestris</i> (Wagner, 1871)		<i>Putoniella</i> Kieffer, 1896	
<i>Harmandia</i> Kieffer, 1896		<i>pruni</i> (Kaltenbach, 1872)	B M S
<i>cavernosa</i> (Rübsaamen, 1899)	B M S	<i>Resseliella</i> Seitner, 1906	
<i>globuli</i> (Rübsaamen, 1889)	B M S	<i>dizygomyzae</i> (Barnes, 1933)	M
<i>populi</i> Rübsaamen, 1917	B M S	<i>oculiperda</i> (Rübsaamen, 1893)	B M S
<i>pustulans</i> Kieffer, 1909	B M S	<i>piceae</i> Seitner, 1906	B M S
<i>tremulae</i> Winnertz, 1853	B M S	<i>quercivora</i> (Mamaev, 1965)	B
<i>Hygrodiplosis</i> Kieffer, 1912		<i>skuhravyorum</i> Skrzypczynska, 1975	B M
<i>vaccinii</i> (Kieffer, 1897)	B M	<i>theobaldi</i> (Barnes, 1927)	B
<i>Lestodiplosis</i> Kieffer, 1894		<i>Sitodiplosis</i> Kieffer, 1913	
<i>alternans</i> Kieffer, 1898	S	<i>mosellana</i> (Géhin, 1857)	B M S
<i>frireni</i> (Kieffer, 1888)	B	<i>Thecodiplosis</i> Kieffer, 1895	
<i>holstei</i> Kieffer, 1920	M	<i>brachyntera</i> (Schwägrichen, 1835)	B M S
<i>longifilis</i> (Kieffer, 1901)	M	<i>Therodiplosis</i> Kieffer, 1912	
<i>polypori</i> (H. Loew, 1850)	B	<i>persicae</i> Kieffer, 1912	B M
<i>raphani</i> Barnes, 1929	M	<i>Thurauia</i> Rübsaamen, 1899	
<i>trifolii</i> Barnes, 1928	M	<i>aquatica</i> Rübsaamen, 1899	B
<i>Loewiola</i> Kieffer, 1896		<i>Tricholaba</i> Rübsaamen, 1917	
<i>centaureae</i> (F. Löw, 1875)	B M S	<i>trifolii</i> Rübsaamen, 1917	B M S
<i>serratulae</i> Kieffer, 1905	B	<i>Trisopsis</i> Kieffer, 1912	
<i>Macro-diplosis</i> Kieffer, 1895		<i>acicularis</i> Mamaev, 1961	M
<i>dryobia</i> (F. Löw, 1877)	B M S	<i>Xenodiplosis</i> Felt, 1911	
<i>volvans</i> Kieffer, 1895	B M S	<i>laeviusculi</i> (Rübsaamen, 1910)	B M
<i>Mamaevia</i> Skuhravá, 1967		<i>Xylodiplosis</i> Kieffer, 1894	
<i>vysineki</i> Skuhravá, 1967	B	<i>aestivalis</i> Kieffer, 1904	B
<i>Massalongia</i> Kieffer, 1897		<i>nigritarsis</i> (Zetterstedt, 1850)	B
<i>rubra</i> (Kieffer, 1890)	B M S	<i>Zeuxidiplosis</i> Kieffer, 1904	
<i>Monarthropalpus</i> Rübsaamen, 1892		<i>giardi</i> (Kieffer, 1896)	S
<i>flavus</i> (Schrark, 1776)	B M S		
<i>Monodiplosis</i> Rübsaamen, 1910			
<i>liebeli</i> (Kieffer, 1889)	B		
<i>Mycodiplosis</i> Rübsaamen, 1895			
<i>kraussei</i> (Wolff, 1910)	S		
<i>melampsorae</i> Rübsaamen, 1889)	B M S		
<i>plasmoparae</i> Rübsaamen, 1906	B		
<i>saundersi</i> Barnes, 1927	B M S		
		PSYCHODIDAE	
		Jan Ježek	
		SYCORACINAE	
		<i>Sycorax</i> Curtis, 1839	
		<i>bicornua</i> Kreck, 1970	B

<i>silacea</i> Curtis, 1839	B M S	<i>verneysicus</i> Vaillant, 1972	B
<i>similis</i> (Müller, 1927)	B M	<i>Parajungiella</i> Vaillant, 1972	
<i>slovacus</i> Halgoš, 1975	S	<i>consors</i> (Eaton, 1893)	B M S
<i>tonnoiri</i> Jung, 1954	B S	= <i>interna</i> (Nielsen, 1964)	
		<i>ellisi</i> (Withers, 1987)	B
TRICHOMYIINAE		<i>longicornis</i> (Tonnoir, 1919)	B M S
<i>Trichomyia</i> Curtis, 1839		<i>pseudolongicornis</i> (Wagner, 1975)	B
<i>urbica</i> Curtis, 1839	B S	<i>serbica</i> (Krek, 1985)	B
		<i>Paramormia</i> Enderlein, 1935	
PSYCHODINAE		sg. <i>Duckhousiella</i> Vaillant, 1972	
MORMIINI		<i>ustulata</i> (Walker, 1856)	B M S
<i>Jovamormia</i> Ježek, 1983		sg. <i>Paramormia</i> s. str.	
<i>caliginosa</i> (Eaton, 1893)	M	<i>polyascoidea</i> (Krek, 1971)	B M S
<i>Lepimormia</i> Enderlein, 1936		<i>Peripsychoda</i> Enderlein, 1935	
<i>palposa</i> (Tonnoir, 1919)	S	<i>auriculata</i> (Curtis, 1839)	B M S
<i>tatrica</i> (Enderlein, 1936)		= <i>zangerii</i> Sarà, 1952	
<i>Oomormia</i> Ježek, 1984		<i>fusca</i> (Macquart, 1826)	M
<i>andrenipes</i> (Strobl, 1910)	B S	<i>Philosepedon</i> Eaton, 1904	
= <i>nigripennis</i> (Krek, 1972)		<i>austriacus</i> Vaillant, 1974	B S
<i>Promormia</i> Ježek, 1983		<i>balkanicus</i> Krek, 1971	B S
<i>eatonj</i> (Tonnoir, 1940)	B M	<i>humeralis</i> (Meigen, 1818)	B M S
<i>silesiensis</i> Ježek, 1983	M	<i>kowarzi</i> Ježek, 1997	B
<i>Psychomormia</i> Ježek, 1983		<i>nickerli</i> Ježek, 1997	B
<i>incerta</i> (Eaton, 1893)	B M	<i>pragensis</i> Ježek, 1997	B
<i>vaillanti</i> (Wagner, 1977)	B	<i>Psycmera</i> Ježek, 1984	
<i>Taramormia</i> Ježek, 1984		<i>integella</i> (Jung, 1956)	B M S
<i>pulcherrima</i> (Wagner, 1979)	B	<i>Telmatoscopus</i> Eaton, 1904	
<i>tatrica</i> Ježek, 1984	S	<i>britteni</i> Tonnoir, 1940	B
		<i>carpathicus</i> Ježek, 1988	S
PARAMORMIINI		<i>carthusianus</i> (Vaillant, 1972)	B
<i>Feuerhorniella</i> Vaillant, 1971		<i>gressicus</i> (Vaillant, 1972)	B M
<i>obscura</i> (Tonnoir, 1919)	B M	<i>Threticus</i> Eaton, 1904	
<i>Jungiella</i> Vaillant, 1972		<i>balkaneoalpinus</i> Krek, 1972	S
sg. <i>Jungiella</i> s. str.		<i>incurvus</i> Krek, 1972	M S
<i>hygrophila</i> Ježek, 1987	B M	<i>lucifugus</i> (Walker, 1856)	B M S
<i>soleata</i> (Walker, 1856)	B M S	<i>negrobovi</i> Vaillant, 1972	S
<i>valachica</i> (Vaillant, 1963)	B M	<i>silvaticus</i> Ježek, 1985	B M S
sg. <i>Psychocha</i> Ježek, 1983		<i>Trichopsychoda</i> Tonnoir, 1922	
<i>acuminata</i> (Szabó, 1960)	B M S	<i>hirtella</i> (Tonnoir, 1919)	B M S
= <i>rozkošnyi</i> Vaillant, 1972			
<i>aquatica</i> Ježek, 1983	B	PSYCHODINI	
<i>laminata</i> (Szabó, 1960)	B	<i>Chodopsycha</i> Ježek, 1984	
<i>procera</i> Krek, 1971	M	<i>lobata</i> (Tonnoir, 1940)	B M S
<i>ripicola</i> (Bellier, 1967)	B M	<i>Copropsychoda</i> Vaillant, 1971	
sg. <i>Psychogella</i> Ježek, 1984		<i>brevicornis</i> (Tonnoir, 1940)	B M S
<i>bohémica</i> Ježek, 1979	B	<i>Logima</i> Eaton, 1904	
<i>inundationum</i> Ježek, 1997	B	<i>albipennis</i> (Zetterstedt, 1850)	B M S
<i>Lepiseodina</i> Enderlein, 1936		= <i>severini severini</i> (Tonnoir, 1922)	
<i>tristis</i> (Meigen, 1830)	B	= <i>severini parthenogenetica</i> (Tonnoir, 1920)	
<i>Panimerus</i> Eaton, 1913		<i>erminea</i> (Eaton, 1893)	B M S
<i>albifacies</i> (Tonnoir, 1919)	B	<i>satchelli</i> (Quate, 1955)	B M S
<i>denticulatus</i> Krek, 1971	B M	<i>zetterstedti</i> Ježek, 1983	B M S
<i>goetghebueri</i> (Tonnoir, 1919)	M	<i>Psycha</i> Ježek, 1984	
<i>kreki</i> Vaillant, 1972	B	<i>griseszens</i> (Tonnoir, 1922)	B M S
<i>maynei</i> (Tonnoir, 1920)	B	<i>Psychoda</i> Latreille, 1796	
<i>notabilis</i> (Eaton, 1893)	B M S	<i>phalaenoides</i> (Linnaeus, 1758)	B M S

<i>uniformata</i> Haseman, 1907	B M	<i>Tonnoiriella</i> Vaillant, 1972	
= <i>moravica</i> Vaillant, 1966		<i>nigricauda</i> (Tonnoir, 1919)	B M
<i>Psychodocha</i> Ježek, 1984		<i>pulchra</i> (Eaton, 1893)	B M S
<i>cinerea</i> (Banks, 1894)	B M S	<i>Ulomyia</i> Walker, 1856	
<i>gemina</i> (Eaton, 1904)	B M S	<i>annulata</i> (Tonnoir, 1919)	B M S
<i>Psychodula</i> Ježek, 1984		<i>cognata</i> (Eaton, 1893)	B M S
<i>minuta</i> (Banks, 1894)	B M S	<i>fuliginosa</i> (Meigen, 1804)	B M S
<i>Psychomora</i> Ježek, 1984		<i>plumata</i> (Tonnoir, 1919)	B M
<i>trinodulosa</i> (Tonnoir, 1922)	B M S	= <i>basaltica</i> Vaillant, 1983	
<i>vanharai</i> Ježek, 1995	M		
<i>Tinearia</i> Schellenberg, 1803		ANISOPODIDAE	
<i>alternata</i> (Say, 1824)	B M S	Ivo Kovář	
<i>lativentris</i> (Berdén, 1952)	B M S		
<i>Ypsidocha</i> Ježek, 1984		<i>Sylvicola</i> Harris, 1776	
<i>setigera</i> (Tonnoir, 1922)	B S	<i>cincta</i> (Fabricius, 1787)	B M
PERICOMINI		<i>fenestralis</i> (Scopoli, 1763)	B M S
<i>Bazarella</i> Vaillant, 1960		<i>fuscata</i> (Fabricius, 1775)	B S
sg. <i>Parabazarella</i> Vaillant, 1983		<i>limpida</i> (Edwards, 1923)	M S
<i>subneglecta</i> (Tonnoir, 1922)	B M S	<i>punctata</i> (Fabricius, 1787)	B M S
<i>Berdeniella</i> Vaillant, 1976			
<i>helvetica</i> (Sarà, 1957)	M		
<i>illiesi</i> Wagner, 1973	S		
<i>manicata</i> (Tonnoir, 1920)	B M S	MYCETOBIIIDAE	
<i>unispinosa</i> (Tonnoir, 1919)	B M S	Jaroslav Martinovský	
<i>vimmeri</i> Ježek, 1997	B M		
<i>Clytocerus</i> Eaton, 1904		MYCETOBIIINAE	
sg. <i>Boreoclytocerus</i> Duckhouse, 1978		<i>Mycetobia</i> Meigen, 1818	
<i>dali</i> (Eaton, 1893)	B S	<i>obscura</i> Mamaev, 1968	B
<i>ocellaris</i> (Meigen, 1804)	B M S	<i>pallipes</i> Meigen, 1818	B M
<i>rivosus</i> (Tonnoir, 1919)	B S		
<i>Pericoma</i> Walker, 1856		SCATOPSIDAE	
sg. <i>Pachypericoma</i> Vaillant, 1978		Jaroslav Martinovský	
<i>blandula</i> Eaton, 1893	B M S		
<i>fallax</i> Eaton, 1893	B M	ASPISTINAE	
sg. <i>Pericoma</i> s. str.		<i>Aspistes</i> Meigen, 1818	
<i>calcilega</i> Feuerborn, 1923	B S	<i>berolinensis</i> Meigen, 1818	B M S
<i>pseudoexquisita</i> Tonnoir, 1940	B M S		
<i>tatrica</i> Szabó, 1960	S		
<i>tonnoiri</i> Vaillant, 1978	S		
<i>trifasciata</i> (Meigen, 1804)	M S	ECTAETIINAE	
<i>Pneumia</i> Enderlein, 1935		<i>Ectaetia</i> Enderlein, 1912	
= <i>Saichelliella</i> Vaillant, 1979		<i>clavipes</i> (Loew, 1846)	B M S
<i>canescens</i> (Meigen, 1804)	B M S		
<i>crispi</i> (Freeman, 1953)	B M	PSECTROSCIARINAE	
<i>gracilis gracilis</i> (Eaton, 1893)	B M	<i>Anapausis</i> Enderlein, 1912	
<i>mutua</i> (Eaton, 1893)	B M S	<i>albohalterata</i> Duda, 1928	B M S
<i>nubila</i> (Meigen, 1818)	B M S	<i>chufourella</i> Haenni, 1984	M S
<i>palustris</i> (Meigen, 1804)	B	<i>haemorrhoidalis</i> Duda, 1928	M S
<i>pilularia</i> (Tonnoir, 1940)	B M S	<i>helvetica</i> Haenni, 1984	M S
<i>plumicornis</i> (Tonnoir, 1922)	B M S	<i>inermis</i> (Ruthé, 1831)	B M S
<i>stammeri</i> (Jung, 1956)	M S	<i>nigripes</i> (Zetterstedt, 1860)	B M
<i>trivialis</i> (Eaton, 1893)	B M S	<i>rectinervis</i> Duda, 1928	B M S
<i>Saraiella</i> Vaillant, 1973		<i>rohaceki</i> Martinovský, 1997	S
<i>rotunda</i> (Krek, 1970)	B M	<i>soluta</i> (Loew, 1846)	M S
<i>Szaboiella</i> Vaillant, 1979		<i>talpae</i> (Verrall, 1912)	B M S
<i>hibernica</i> (Tonnoir, 1940)	B M		

SCATOPSINAE

SCATOPSINI

Apiloscatopse Cook, 1974*flavicollis* (Meigen, 1818) B M S*flavocincta* (Duda, 1928) B M*gracilis* (Duda, 1928) M S*hennigi* Cook, 1974 S*mattheyi* Haenni, 1981 M*picea* (Meigen, 1818) B M S*scutellata* (Loew, 1846) B M S*styriaca* (Enderlein, 1926) S*Colobostema* Enderlein, 1926*nigripennis* (Meigen, 1830) B M S*triste* (Zetterstedt, 1850) B M S*Cookella* Freeman, 1985*albitarsis* (Zetterstedt, 1850) B M S*Ferneiella* Cook, 1974*brevifurca* (Enderlein, 1912) B*incompleta* (Verrall, 1886) B M S*Holoplagia* Enderlein, 1912*bullata* (Edwards, 1925) M S*lucifuga* (Loew, 1870) M S*richardsi* (Edwards, 1934) S*transversalis* (Loew, 1846) B S*Reichertella* Enderlein, 1912sg. *Pharsoreichertella* Cook, 1974*hamifera* (Strobl, 1909) B*simplicinervis* (Duda, 1928) Ssg. *Reichertella* s. str.*geniculata* (Zetterstedt, 1850) B M S*nigra* (Meigen, 1804) B M S*pulicaria* (Loew, 1846) B M S*Scatopse* Geoffroy, 1762*notata* (Linné, 1758) B M S

SWAMMERDAMELLINI

Coboldia Melander, 1916*fuscipes* (Meigen, 1830) B M S*Rhexosa* Enderlein, 1936*subnitens* (Verrall, 1886) M S*Swammerdamella* Enderlein, 1912*acuta* Cook, 1956 B S*adecotris* Cook, 1972 B S*brevicornis* (Meigen, 1830) B M S*genypodis* Cook, 1972 S*pediculata* (Duda, 1928) S

THRIPOMORPHINI

Thripomorpha Enderlein, 1905*bifida* (Zilahy-Sebess, 1956) B M S*collini* (Cook, 1969)*freyi* (Cook, 1969) B M*haenni* Martinovský, 1997 S*halteratum* (Meigen, 1838) B M S*paludicola* Enderlein, 1905 B M S*edwardsi* (Collin, 1954)*verralli* (Edwards, 1934) B M S

SYNNEURIDAE

Jaroslav Martinovský

Synneuron Lundström, 1910*annulipes* Lunström, 1910 S*= sylvestre* Mamaev & Krivosheina, 1969

CANTHYLOSCELIDAE

Jaroslav Martinovský

Hyperoscelis Hardy & Nagatomi, 1960*eximia* (Boheman, 1858) M*veternosa* Mamaev & Krivosheina, 1969 S

PTYCHOPTERIDAE

Jaroslav Starý

Ptychoptera Meigen, 1803*albimana* (Fabricius, 1787) B M S*contaminata* (Linnaeus, 1758) B M S*handlirschi* (Czížek, 1919) M S*lacustris* Meigen, 1830 B M S*longicauda* (Tonnoir, 1919) B M S*minuta* Tonnoir, 1919 B*paludosa* Meigen, 1804 B M S*scutellaris* Meigen, 1818 B M S*silvicola* Zwyrtek & Rozkošný, 1967 S

DIXIDAE

Jan Knoz

Dixa Meigen, 1818*dilatata* Strobl, 1984 M*maculata* Meigen, 1818 B M*nebulosa* Meigen, 1830 B M S*nubilipennis* Curtis, 1832 M*puberula* Loew, 1849 M S*submaculata* Edwards, 1920 M S*Dixella* Dyar & Shannon, 1924*aestivalis* (Meigen, 1938) B M*amphibia* (De Geer, 1776) B M*autumnalis* (Meigen, 1838) B M*hyperborea* (Bergroth, 1889) S*nigra* (Staeger, 1840) S*serotina* (Meigen, 1818) B

CHAOBORIDAE

Ladislav Jedlička & Ivan Országh

CHAOBORINAE

CHAOBORINI

Chaoborus Lichtenstein, 1800sg. *Chaoborus* s. str.*crystallinus* (De Geer, 1776) B M S

<i>flavicans</i> (Meigen, 1830)	B	M	S	<i>pulchritarsis</i> (Rondani, 1872)	B	S
<i>obscuripes</i> (van der Wulp, 1859)	B		S	rusticus group		
sg. <i>Peusomyia</i> Saether, 1970				<i>refiki</i> Medjid, 1928	M	S
<i>pallidus</i> (Fabricius, 1794)	B		S	<i>rusticus</i> (Rossi, 1790)	B	S
MOCHLONYXINI				<i>Culex</i> Linnaeus, 1758		
<i>Mochlonyx</i> Loew, 1844				sg. <i>Barraudius</i> Edwards, 1912		
<i>fuliginosus</i> (Felt, 1905)	B			<i>modestus</i> Ficalbi, 1890	B	M S
<i>velutinus</i> (Ruthe, 1831)	B			sg. <i>Culex</i> s. str.		
CULICIDAE				<i>pipiens pipiens</i> Linnaeus, 1758	B	M S
Jan Minář & Jozef Halgoš				<i>pipiens molestus</i> Forskål, 1775	B	M S
ANOPHELINAE				<i>theileri</i> Theobald, 1903		S
<i>Anopheles</i> Meigen, 1818				<i>torrentium</i> Martini, 1924	B	M S
sg. <i>Anopheles</i> s. str.				sg. <i>Neoculex</i> Dyar, 1905		
<i>atroparvus</i> Van Thiel, 1927		M	S	<i>martinii</i> Medjid, 1930		M
<i>claviger</i> (Meigen, 1804)	B	M	S	<i>terrilians</i> Walker, 1856	B	M S
<i>labbranchiae</i> Falleroni, 1926		M	S	sg. <i>Maillotia</i> Theobald, 1907		
<i>maculipennis</i> Meigen, 1818	B	M	S	<i>hortensis</i> Ficalbi, 1889	B	S
<i>messeae</i> Falleroni, 1926	B	M	S	<i>Culiseta</i> Felt, 1904		
<i>plumbeus</i> Stephens, 1828	B	M	S	sg. <i>Allotheobaldia</i> Broelemann, 1919		
CULICINAE				<i>longiareolata</i> (Macquart, 1838)		S
<i>Aedes</i> Meigen, 1818				sg. <i>Culicella</i> Felt, 1904		
sg. <i>Aedes</i> s. str.				<i>morsitans</i> (Theobald, 1901)	B	M S
<i>cinereus</i> Meigen, 1818	B	M	S	<i>ochroptera</i> (Peus, 1935)	B	S
<i>geminus</i> Peus, 1970			S	sg. <i>Culiseta</i> s. str.		
<i>rossicus</i> Dolbeskhin, Gorickaja & Mitrofanova, 1930	B	M	S	<i>alaskaensis</i> (Ludlow, 1906)	B	M S
sg. <i>Aedimorphus</i> Theobald, 1903				<i>annulata annulata</i> (Schränk, 1776)	B	M S
<i>vexans</i> Meigen, 1830	B	M	S	<i>annulata subochrea</i> (Edwards in Wesenberg-Lund, 1921)	B	M S
sg. <i>Finlaya</i> Theobald, 1903				<i>glaphyoptera</i> (Schiner, 1864)	B	M S
<i>geniculatus</i> (Olivier, 1791)	B	M	S	<i>Mansonia</i> Blanchard, 1901		
sg. <i>Ochlerotatus</i> Lynch Arribálzaga, 1891				sg. <i>Coquillettidia</i> Dyar, 1905		
cantans group				<i>richiardii</i> (Ficalbi, 1889)	B	M S
<i>annulipes</i> (Meigen, 1830)	B	M	S	<i>Uranotaenia</i> Lynch Arribálzaga, 1891		
<i>behningi</i> Martini, 1926			S	sg. <i>Pseudoficalbia</i> Theobald, 1912		
<i>cantans</i> (Meigen, 1818)	B	M	S	<i>unguiculata</i> Edwards, 1913		M S
<i>cyprinus</i> Ludlow, 1920			S	CHIRONOMIDAE		
<i>excrucians</i> (Walker, 1856)	B	M	S	Peter Bitušik & Bohumil Losos		
<i>flavescens</i> (Müller, 1764)	B	M	S	TANYPODINAE		
<i>ripartus</i> Dyar & Knab, 1907	B			COELOTANYPODINI		
communis group				<i>Clinotanypus</i> Kieffer, 1913		
<i>cataphylla</i> Dyar, 1916	B	M	S	<i>nervosus</i> (Meigen, 1818)	B	M
<i>communis</i> (De Geer, 1776)	B	M	S	TANYPODINI		
<i>diantaeus</i> Howard, Dyar & Knab, 1917	B	M	S	<i>Tanypus</i> Meigen, 1803		
<i>hungaricus</i> Mihályi, 1955			S	<i>kraatzii</i> (Kieffer, 1912)	B	M S
<i>intrudens</i> Dyar, 1919	B	M	S	<i>punctipennis</i> Meigen, 1818	B	M S
<i>leucomelas</i> (Meigen, 1804)	B	M	S	<i>vilipennis</i> (Kieffer, 1918)	B	S
<i>nigrinus</i> (Eckstein, 1918)			S	PROCLADIINI		
<i>pullatus</i> (Coquillett, 1904)	B		S	<i>Procladius</i> Skuse, 1889		
<i>puncator</i> (Kirby in Richardson, 1837)	B	M	S	sg. <i>Holotanypus</i> Roback, 1982		
<i>sticticus</i> (Meigen, 1838)	B	M	S	<i>choreus</i> (Meigen, 1804)	B	M S
dorsalis group				<i>culiciformis</i> (Linnaeus, 1767)	B	
<i>caspinus</i> (Pallas, 1771)	B	M	S	<i>sagittalis</i> (Kieffer, 1909)		S
<i>dorsalis</i> (Meigen, 1830)	B	M	S	<i>signatus</i> (Zetterstedt, 1850)	B	S

sg. <i>Psilotanypus</i> Kieffer, 1906			<i>Xenopelopia</i> Fittkau, 1962		
<i>rufovittatus</i> (van der Wulp, 1874)	S		<i>falcigera</i> (Kieffer, 1911)	B M S	
ANATOPYNIINI			<i>Zavrelimyia</i> Fittkau, 1962		
<i>Anatopynia</i> Johannsen, 1905			<i>barbatipes</i> (Kieffer, 1911)	M S	
<i>plumipes</i> (Fries, 1823)	B M		<i>hirtimana</i> (Kieffer, 1918)	B M	
MACROPELOPIINI			<i>melamura</i> (Meigen, 1804)	B M S	
<i>Apsectrotanypus</i> Fittkau, 1962			<i>nubila</i> (Meigen, 1830)	B M	
<i>trifasciipennis</i> (Zetterstedt, 1838)	B M S		<i>signatipennis</i> (Kieffer, 1924)	S	
<i>Macropelopia</i> Thienemann, 1916			BUCHONOMYIINAE		
<i>adaucta</i> Kieffer, 1916	B M S		<i>Buchonomyia</i> Fittkau, 1955		
<i>nebulosa</i> (Meigen, 1804)	B M S		<i>thienemanni</i> Fittkau, 1955	S	
<i>notata</i> (Meigen, 1818)	B M S		DIAMESINAE		
<i>Psectrotanypus</i> Kieffer, 1909			BOREOHEPTAGYINI		
<i>varius</i> (Fabricius, 1787)	B M S		<i>Boreoheptagyia</i> Brundin, 1966		
NATARSIINI			<i>legeri</i> (Goetghebuer, 1933)	M S	
<i>Natarsia</i> Fittkau, 1962			<i>monticola</i> (Serra-Tosio, 1964)	S	
<i>punctata</i> (Meigen, 1804)	B M S		DIAMESINI		
PENTANEURINI			<i>Diamesa</i> Meigen, 1835		
<i>Ablabesmyia</i> Johannsen, 1905			<i>bertrami</i> Edwards, 1935	S	
<i>monilis</i> (Linnaeus, 1758)	B M S		<i>cinerella</i> Meigen, 1835	S	
<i>Conchapelopia</i> Fittkau, 1957			<i>dampfi</i> (Kieffer, 1924)	S	
<i>melanops</i> (Meigen, 1818)	B M S		<i>incallida</i> (Walker, 1856)	S	
<i>pallidula</i> (Meigen, 1818)	B M S		<i>insignipes</i> Kieffer, 1908	M S	
<i>viator</i> (Kieffer, 1911)	S		<i>kasymovi</i> Kownacki & Kownacka, 1973	S	
<i>Guttipelopia</i> Fittkau, 1962			<i>permacra</i> (Walker, 1856)	S	
<i>guttipennis</i> (van der Wulp, 1861)	B M S		<i>starmachi</i> Kownacki & Kownacka, 1970	S	
<i>Krenopelopia</i> Fittkau, 1962			<i>steinboeckii</i> Goetghebuer, 1938	S	
<i>binotata</i> (Wiedemann, 1817)	B M S		<i>tonsa</i> (Walker, 1856)	B M S	
<i>nigropunctata</i> (Staeger, 1839)	B		<i>vallanti</i> Serra-Tosio, 1972	S	
<i>Larsia</i> Fittkau, 1962			<i>Potthastia</i> Kieffer, 1922		
<i>curticalcar</i> (Kieffer, 1918)	M S		<i>gaedii</i> (Meigen, 1838)	M S	
<i>Monopelopia</i> Fittkau, 1962			<i>longimana</i> (Kieffer, 1922)	M S	
<i>temicalcar</i> (Kieffer, 1918)	B M S		<i>Pseudodiamesa</i> Goetghebuer, 1939		
<i>Nilotanypus</i> Kieffer, 1923			<i>branickii</i> (Nowicki, 1873)	B M S	
<i>dubius</i> (Meigen, 1804)	S		<i>nivosa</i> (Goetghebuer, 1928)	S	
<i>Paramerina</i> Fittkau, 1962			<i>Pseudokiefferiella</i> Zavřel, 1941		
<i>cingulata</i> (Walker, 1856)	B M S		<i>parva</i> (Edwards, 1932)	S	
<i>divisa</i> (Walker, 1856)	S		<i>Sympotthastia</i> Pagast, 1947		
<i>Pentaneurella</i> Fittkau & Murray, 1983			<i>zavreli</i> Pagast, 1947	M	
<i>katterjoki</i> Fittkau & Murray, 1983	S		<i>Syndiamesa</i> Kieffer, 1918		
<i>Rheopelopia</i> Fittkau, 1962			<i>hygropetrica</i> (Kieffer, 1909)	S	
<i>ornata</i> (Meigen, 1838)	M S		PRODIAMESINAE		
<i>Telopelopia</i> Fittkau, 1962			<i>Monodiamesa</i> Kieffer, 1922		
<i>fascigera</i> (Verneaux, 1970)	S		<i>bathyphila</i> (Kieffer, 1918)	S	
<i>Telmatopelopia</i> Fittkau, 1962			<i>Odontomesa</i> Pagast, 1947		
<i>nemorum</i> (Goetghebuer, 1921)	M S		<i>fulva</i> (Kieffer, 1919)	M S	
<i>Thienemannimyia</i> Fittkau, 1957			<i>Prodiamesa</i> Kieffer, 1906		
<i>carnea</i> (Fabricius, 1805)	B M S		<i>olivacea</i> (Meigen, 1818)	B M S	
<i>geijeskesi</i> (Goetghebuer, 1934)	M S		<i>rufovittata</i> Goetghebuer, 1932	S	
<i>laeta</i> (Meigen, 1818)	B S		ORTHOCLADIINAE		
<i>lentiginosa</i> (Fries, 1832)	M		<i>Acricotopus</i> Kieffer, 1921		
<i>Trissopelopia</i> Kieffer, 1923			<i>lucens</i> (Zetterstedt, 1850)	B M S	
<i>flavida</i> Kieffer, 1923	B M S				

<i>Brillia</i> Kieffer, 1913			<i>coerulescens</i> (Kieffer, 1926)	M	S		
<i>longifurca</i> Kieffer, 1921	B	M	S	<i>cyanea</i> Thienemann, 1936	M	S	
<i>modesta</i> (Meigen, 1830)	B	M	S	<i>devonica</i> (Edwards, 1929)	M	S	
<i>Bryophaenocladus</i> Thienemann, 1934				<i>fuldensis</i> Lehmann, 1972		S	
<i>muscicola</i> (Kieffer, 1906)			S	<i>gracei</i> (Edwards, 1929)	B	M	S
<i>subvernalis</i> (Edwards, 1929)			S	<i>ilkleyensis</i> (Edwards, 1929)	M	S	
<i>Camptocladus</i> van der Wulp, 1874				<i>lobifera</i> Goetghebuer, 1934		S	
<i>stercorarius</i> (De Geer, 1776)			S	<i>minor</i> (Edwards, 1929)	M	S	
<i>Cardiocladus</i> Kieffer, 1912				<i>pseudomontana</i> Goetghebuer, 1935	M	S	
<i>capucinus</i> (Zetterstedt, 1850)			S	<i>similis</i> Goetghebuer, 1939	M		
<i>fuscus</i> Kieffer, 1924	M		S	<i>tirolensis</i> Goetghebuer, 1938		S	
<i>Chaetocladus</i> Kieffer, 1911				<i>Gymnometriocnemus</i> Goetghebuer, 1932			
<i>dentiforceps</i> (Edwards, 1929)			S	<i>subnudus</i> (Edwards, 1929)		S	
<i>perennis</i> (Meigen, 1830)			S	<i>Heleniella</i> Gouin, 1934			
<i>suecicus</i> (Kieffer, 1916)	M			<i>ornaticollis</i> (Edwards, 1929)	M	S	
<i>Corynoneura</i> Winnertz, 1846				<i>serratosioi</i> Ringe, 1976		S	
<i>carriana</i> Edwards, 1924			S	<i>Heterotanytarsus</i> Spärck, 1923			
<i>celeripes</i> Winnertz, 1852	B	M	S	<i>apicalis</i> (Kieffer, 1921)	B		
<i>celtica</i> Edwards, 1924			S	<i>Heterotrissocladus</i> Spärck, 1923			
<i>gratias</i> Schlee, 1968			S	<i>marcidus</i> (Walker, 1856)	B	M	S
<i>lobata</i> Edwards, 1924	M		S	<i>Hydrobaenus</i> Fries, 1830			
<i>scutellata</i> Winnertz, 1846			S	<i>distylus</i> (Kieffer, 1915)	M		
<i>Cricotopus</i> van der Wulp, 1874				<i>lugubris</i> Fries, 1830			S
sg. <i>Cricotopus</i> s. str.				<i>rufus</i> (Kieffer, 1923)	B		
<i>albiforceps</i> (Kieffer, 1916)			S	<i>Krenosmittia</i> Thienemann & Krüger, 1939			
<i>algarum</i> (Kieffer, 1911)	M			<i>boreoalpina</i> (Goetghebuer, 1944)			S
<i>annulator</i> Goetghebuer, 1927			S	<i>camptophleps</i> (Edwards, 1929)			S
<i>beckeri</i> Hirvenoja, 1973			S	<i>Limnophyes</i> Eaton, 1875			
<i>hicinctus</i> (Meigen, 1818)	M		S	<i>asquamatus</i> Andersen, 1937			S
<i>curtus</i> Hirvenoja, 1973			S	<i>minimus</i> (Meigen, 1818)	M		S
<i>ephippium</i> (Zetterstedt, 1838)			S	<i>ninae</i> Saether, 1975			S
<i>festivellus</i> (Kieffer, 1906)			S	<i>Metriocnemus</i> van der Wulp, 1874			
<i>fuscus</i> (Kieffer, 1909)			S	<i>atratus</i> (Zetterstedt, 1850)		M	
<i>similis</i> Goetghebuer, 1921			S	<i>fuscipes</i> (Meigen, 1818)	B	M	S
<i>tibialis</i> (Meigen, 1804)	M		S	<i>hirticollis</i> (Staeger, 1839)		M	S
<i>tremulus</i> (Linnaeus, 1758)			S	<i>hygropetricus</i> Kieffer, 1912	B		
<i>triannulatus</i> Macquart, 1826			S	<i>martinii</i> Thienemann, 1921	B	M	S
<i>trifascia</i> Edwards, 1929	M		S	<i>Nanocladus</i> Kieffer, 1913			
<i>vierriensis</i> Goetghebuer, 1935	M		S	<i>balticus</i> (Palmen, 1959)			S
sg. <i>Isocladus</i> Kieffer, 1909				<i>bicolor</i> (Zetterstedt, 1838)	B	M	S
<i>brevipalpis</i> Kieffer, 1909	B	M		<i>parvulus</i> (Kieffer, 1909)			S
<i>intersectus</i> (Staeger, 1839)			S	<i>rectinervis</i> (Kieffer, 1911)			S
<i>ornatus</i> (Meigen, 1818)			S	<i>Orthocladus</i> van der Wulp, 1874			
<i>sylvestris</i> (Fabricius, 1794)	B	M	S	sg. <i>Eudactylocladius</i> Thienemann, 1935			
<i>tricinctus</i> (Meigen, 1818)			S	<i>fuscimanus</i> (Kieffer, 1908)	B	M	S
<i>trifascianus</i> (Meigen, 1813)	M		S	<i>olivaceus</i> (Kieffer, 1911)			S
<i>Diplocladius</i> Kieffer, 1908				sg. <i>Euorthocladus</i> Thienemann, 1935			
<i>cultriger</i> Kieffer, 1908	B	M	S	<i>ashei</i> Sopenis, 1990			S
<i>Dratnalia</i> Saether & Halvorsen, 1981				<i>rivicola</i> Kieffer, 1921		M	S
<i>potamophylaxi</i> Fittkau & Lellák, 1971		M	S	<i>rivulorum</i> Kieffer, 1909		M	S
<i>Epoicocladus</i> Šulc & Zavřel, 1924				<i>saxosus</i> (Tokunaga, 1939)		M	S
<i>flavens</i> (Malloch, 1915)	B	M	S	<i>thienemanni</i> Kieffer, 1906		M	S
<i>Eukiefferiella</i> Thienemann, 1926				sg. <i>Symposiocladius</i> Cranston, 1982			
<i>ancyla</i> Svensson, 1986			S	<i>lignicola</i> (Kieffer, 1915)		M	S
<i>brevicalcar</i> (Kieffer, 1911)	B	M	S	sg. <i>Orthocladus</i> s. str.			
<i>claripennis</i> (Lundbeck, 1898)	B	M	S	<i>excavatus</i> Brundin, 1947			S
<i>clypeata</i> (Kieffer, 1923)	M		S	<i>frigidus</i> (Zetterstedt, 1838)	B	M	S

<i>glabripennis</i> (Goetghebuer, 1921)	B	S	<i>filiformis</i> (Kieffer, 1908)	S
<i>oblidens</i> (Walker, 1856)		S	<i>Pseudosmittia</i> Goetghebuer, 1932	
<i>obumbratus</i> Johannsen, 1905	M	S	<i>gracilis</i> (Goetghebuer, 1913)	M
<i>rivinus</i> Kieffer, 1915		S	<i>trilobata</i> (Edwards, 1929)	S
<i>rubicundus</i> (Meigen, 1818)	M	S	<i>Rheocricotopus</i> Thienemann	
– <i>saxicola</i> Kieffer, 1911			& Harnisch, 1932	
<i>ruffoi</i> Rossaro & Prato, 1991		S	<i>atripes</i> (Kieffer, 1913)	S
<i>wetterensis</i> Brundin, 1956	M	S	<i>chalybeatus</i> (Edwards, 1929)	S
<i>Paracladius</i> Hirvenoja, 1973			<i>effusus</i> (Walker, 1856)	M S
<i>alpicola</i> (Zetterstedt, 1850)		S	<i>fuscipes</i> (Kieffer, 1909)	M S
<i>conversus</i> (Walker, 1856)		S	<i>glabricollis</i> (Meigen, 1830)	S
<i>Paracricotopus</i> Thienemann & Harnisch, 1932			<i>tirolus</i> Lehmann, 1969	S
<i>niger</i> (Kieffer, 1913)	M	S	<i>Rheosmittia</i> Brundin, 1956	
<i>Parakiefferiella</i> Thienemann, 1936			<i>languida</i> (Brundin, 1956)	S
<i>bathophila</i> (Kieffer, 1912)		S	<i>spincornis</i> (Brundin, 1956)	S
<i>coronata</i> (Edwards, 1929)		S	<i>Smittia</i> Holmgren, 1869	
<i>pyrenaica</i> Moubayed, 1991		S	<i>contingens</i> (Walker, 1956)	S
<i>scandica</i> Brundin, 1956		S	<i>Stilocladius</i> Rossaro, 1979	
<i>smolandica</i> (Brundin, 1947)		S	<i>montanus</i> Rossaro, 1979	S
<i>wulkeri</i> Moubayed, 1994		S	<i>Symbiocladius</i> Kieffer, 1925	
<i>Parametriocnemus</i> Goetghebuer, 1932			<i>rhithrogenae</i> (Zavřel, 1924)	B M S
<i>arciger</i> (Kieffer, 1925)	B	M	<i>Synorthocladius</i> Thienemann, 1935	
<i>boreoalpinus</i> Gouin, 1942		S	<i>semivirens</i> (Kieffer, 1909)	B M S
<i>stylatus</i> (Kieffer, 1924)	B	M S	<i>Thienemannia</i> Kieffer, 1909	
<i>Paraphaenocladius</i> Thienemann, 1924			<i>fulvofasciata</i> (Kieffer, 1921)	S
<i>pseudirritus</i> Strenzke, 1950		S	<i>Thienemanniella</i> Kieffer, 1911	
<i>Paratrachocladius</i> Santos-Abreu, 1918			<i>clavicornis</i> (Kieffer, 1911)	M S
<i>rufiventris</i> (Meigen, 1830)		S	<i>vittata</i> (Edwards, 1924)	S
<i>skirwithensis</i> (Edwards, 1929)		S	<i>Tokunagaia</i> Saether, 1973	
<i>Paratrissocladius</i> Zavřel, 1937			<i>tonollii</i> (Rossaro, 1983)	S
<i>excerptus</i> (Walker, 1856)	M	S	<i>Tvetenia</i> Kieffer, 1922	
<i>Parorthocladius</i> Thienemann, 1935			<i>bavarica</i> (Goetghebuer, 1934)	M S
<i>nigrinus</i> (Goetghebuer, 1938)		S	<i>calvescens</i> (Edwards, 1929)	M S
<i>nudipennis</i> (Kieffer, 1908)	M	S	<i>discoloripes</i> (Goetghebuer, 1936)	M S
<i>Prosilocerus</i> Kieffer, 1923			<i>verralli</i> (Edwards, 1929)	M S
<i>lusatiensis</i> Barthelmes, 1964		S	<i>Zalutschia</i> Lipina, 1939	
<i>Psectrocladius</i> Kieffer, 1906			<i>tatrica</i> (Pagast in Zavřel & Pagast, 1935)	S
sg. <i>Allopsectrocladius</i> Wülker, 1956				
<i>obvius</i> (Walker, 1856)	B	S	CHIRONOMINAE	
– <i>dilatatus</i> (van der Wulp, 1858)			CHIRONOMINI	
<i>platypus</i> (Edwards, 1929)		S	<i>Chironomus</i> Meigen, 1803	
sg. <i>Mesopsectrocladius</i> Laville, 1971			sg. <i>Camptochironomus</i> Kieffer, 1918	
<i>barbatipes</i> Kieffer, 1923	B	M	<i>pallidivittatus</i> (Malloch, 1915)	S
sg. <i>Monopsectrocladius</i> Laville, 1971			<i>tentans</i> Fabricius, 1805	B M
<i>calcaratus</i> (Edwards, 1929)	M	S	sg. <i>Chironomus</i> s. str.	
sg. <i>Psectrocladius</i> s. str.			<i>annularius</i> Meigen, 1818	B S
<i>barbimanus</i> (Edwards, 1929)	B	M	<i>balatonicus</i> Dévai, Wülker & Schöll,	
<i>brehmi</i> Kieffer, 1923	B		1983	B S
<i>limbatellus</i> (Holmgren, 1869)		S	<i>bernensis</i> Klötzli, 1973	S
<i>octomaculatus</i> Wülker, 1956		S	<i>cingulatus</i> Meigen, 1830	B
<i>oligosenus</i> Wülker, 1956		S	<i>commutatus</i> Keyl, 1960	B
<i>oxyura</i> Langton, 1985		S	<i>dorsalis</i> Meigen, 1818	B M S
<i>psilopterus</i> (Kieffer, 1906)	B	M	<i>holomelas</i> Keyl, 1961	B
<i>sordidellus</i> (Zetterstedt, 1838)	B	M S	<i>lacunarius</i> Wülker, 1973	S
<i>Pseudorthocladius</i> Goetghebuer, 1932			<i>longipes</i> Staeger 1839	S
<i>berthelemyi</i> Moubayed, 1988		S	<i>longistylus</i> Goetghebuer, 1921	S
<i>curtistylus</i> (Goetghebuer, 1921)		S	<i>luridus</i> Strenzke, 1959	B S

<i>melanescens</i> Keyl, 1961	B		<i>Kiefferulus</i> Goetghebuer, 1922	
<i>melanotus</i> Keyl, 1961	B		<i>tendipediformis</i> (Goetghebuer, 1921)	S
<i>nuditarsis</i> Keyl, 1961	B	S	<i>Lauterborniella</i> Thienemann & Bause, 1913	
<i>obtusidens</i> Goetghebuer, 1921	B		<i>agrayloides</i> (Kieffer, 1911)	B M S
<i>parathummi</i> Keyl, 1961	B		<i>Microchironomus</i> Kieffer, 1918	
<i>piger</i> Strenzke, 1959	B		<i>tener</i> (Kieffer, 1918)	S
<i>plumosus</i> (Linnaeus, 1758)	B M S		<i>Microtendipes</i> Kieffer, 1915	
<i>pseudothummi</i> Strenzke, 1959	B	S	<i>chloris</i> (Meigen, 1818)	B M S
<i>riparius</i> Meigen, 1804	B M S		<i>pedellus</i> (De Geer, 1776)	B S
<i>sororius</i> Wülker, 1973	B		<i>tarsalis</i> (Walker, 1856)	M
<i>uliginosus</i> Keyl, 1960	B		<i>Omisus</i> Townes, 1945	
sg. <i>Lobochironomus</i> Ryser, Wülker & Scholl, 1990			<i>caledonicus</i> (Edwards, 1932)	S
<i>montuosus</i> Ryser, Wülker & Scholl, 1985		S	<i>Parachironomus</i> Lenz, 1921	
<i>Cladopelma</i> Kieffer, 1921			<i>arcuatus</i> (Goetghebuer, 1919)	S
= <i>Cryptocladopelma</i> Lenz, 1941			<i>frequens</i> (Johannsen, 1905)	M S
<i>lateralis</i> (Goetghebuer, 1934)		S	<i>parilis</i> (Walker, 1856)	M S
<i>virescens</i> (Meigen, 1818)		S	<i>varus</i> (Goetghebuer, 1921)	M S
<i>viridula</i> (Linnaeus, 1767)		S	<i>Paracladopelma</i> Harnisch, 1923	
<i>Cryptochironomus</i> Kieffer, 1918			<i>camptolabis</i> (Kieffer, 1913)	B M S
<i>albofasciatus</i> (Staeger, 1839)		S	<i>mikiana</i> (Goetghebuer, 1937)	S
<i>defectus</i> (Kieffer, 1913)	B M S		<i>nigritula</i> (Goetghebuer, 1942)	S
<i>denticulatus</i> (Goetghebuer, 1921)		S	<i>Paratendipes</i> Kieffer, 1911	
<i>obreptans</i> (Walker, 1856)		S	<i>albimanus</i> (Meigen, 1818)	B M S
<i>psittacinus</i> (Meigen, 1830)		S	<i>mudisquama</i> (Edwards, 1929)	S
<i>rostratus</i> Kieffer, 1921		S	<i>Polypedilum</i> Kieffer, 1912	
<i>supplicans</i> (Meigen, 1830)		S	sg. <i>Pentapedilum</i> Kieffer, 1913	
<i>Cryptotendipes</i> Lenz, 1941			<i>exsectum</i> (Kieffer, 1916)	M
<i>holsatus</i> Lenz, 1959		S	<i>sordens</i> (van der Wulp, 1874)	S
<i>pseudotener</i> (Goetghebuer, 1922)		S	<i>uncinatum</i> (Goetghebuer, 1921)	S
<i>Demeijerea</i> Kruseman, 1933			sg. <i>Polypedilum</i> s. str.	
<i>rufipes</i> (Linnaeus, 1761)	B M		<i>albicorne</i> (Meigen, 1838)	S
<i>Demicryptochironomus</i> Lenz, 1941			<i>arudinetum</i> (Goetghebuer, 1921)	S
<i>vulnerans</i> (Zetterstedt, 1838)	B	S	<i>convictum</i> (Walker, 1856)	M S
<i>Dicrotendipes</i> Kieffer, 1913			<i>cultellatum</i> Goetghebuer, 1931	S
<i>modestus</i> (Say, 1823)		S	<i>laetum</i> (Meigen, 1818)	M S
<i>nervosus</i> (Staeger, 1839)	B M S		<i>nubeculosum</i> (Meigen, 1804)	B M S
<i>Einfeldia</i> Kieffer, 1924			<i>pedestre</i> (Meigen, 1830)	M S
<i>dissidens</i> (Walker, 1856)		S	sg. <i>Tripodura</i> Townes, 1945	
<i>pagana</i> (Meigen, 1838)		M	<i>aegyptium</i> Kieffer, 1925	S
<i>Endochironomus</i> Kieffer, 1918			<i>bicrenatum</i> Kieffer, 1921	S
<i>albipennis</i> (Meigen, 1830)	B M S		<i>pullum</i> (Zetterstedt, 1838)	S
<i>dispar</i> (Meigen, 1830)	B M S		<i>scalaenum</i> (Schrank, 1803)	M
<i>tendens</i> (Fabricius, 1775)	B M S		<i>Saetheria</i> Jackson, 1977	
<i>Fleuria</i> Kieffer, 1924			<i>reissi</i> Jackson, 1977	S
<i>lacustris</i> Kieffer, 1924	B	S	<i>Sergentia</i> Kieffer, 1922	
<i>Glyptotendipes</i> Kieffer, 1913			<i>longiventris</i> Kieffer, 1924	S
<i>barbipes</i> (Staeger, 1839)	B M S		<i>Stenochironomus</i> Kieffer, 1919	
<i>follicola</i> Kieffer, 1918		S	<i>gibbus</i> (Fabricius, 1794)	S
<i>gripekoveni</i> (Kieffer, 1913)	B M S		<i>Stictochironomus</i> Kieffer, 1919	
<i>pallens</i> (Meigen, 1804)	B M S		<i>maculipennis</i> (Meigen, 1818)	S
<i>paripes</i> (Edwards, 1929)	B	S	<i>sticticus</i> (Fabricius, 1781)	B M S
<i>signatus</i> (Kieffer, 1909)		S	<i>Tribelos</i> Townes, 1945	
<i>Harnischia</i> Kieffer, 1921			<i>intextus</i> (Walker, 1856)	S
<i>curtilamellata</i> (Malloch, 1915)		S	<i>Xenochironomus</i> Kieffer, 1921	
<i>fuscimana</i> Kieffer, 1921	M S		<i>xenolabis</i> Kieffer, 1916	B M S
			<i>Zavreliella</i> Kieffer, 1920	
			<i>marmorata</i> (van der Wulp, 1858)	B M

PSEUDOCHIRONOMINI

Pseudochironomus Malloch, 1915*prasinatus* (Staeger, 1839) S

TANYTARSINI

Cladotanytarsus Kieffer, 1921*atridorsum* Kieffer, 1924 M S*lepidocalcar* Krüger, 1938 S*manus* (Walker, 1856) B M S*molesus* (Hirvenoja, 1962) S*pallidus* Kieffer, 1922 B S*Corynocera* Zetterstedt, 1838*ambigua* Zetterstedt, 1838 B*Micropsectra* Kieffer, 1909*aristata* Pinder, 1976 S*atrofasciata* (Kieffer, 1911) M S*bidentata* (Goetghebuer, 1921) S*fusca* (Meigen, 1804) S*junci* (Meigen, 1818) M S*lindrothi* Goetghebuer, 1931 S*notescens* (Walker, 1856) S*recurvata* Goetghebuer, 1928 M*Neostempellina* Reiss, 1984*thienemanni* Reiss, 1984 S*Neozavrelia* Goetghebuer, 1941*luteola* Goetghebuer, 1941 M S*Parapsectra* Reiss, 1969*styriaca* (Reiss, 1969) S*uliginosa* Reiss, 1969 S*Paratanytarsus* Thienemann & Bause, 1913*austriacus* (Kieffer, 1924) S*bituberculatus* (Edwards, 1929) S*boemicus* Kieffer, 1922 B M*confusus* Palmén, 1960 S*inopertus* (Walker, 1856) S*intricatus* (Goetghebuer, 1921) B S*laccophilus* (Edwards, 1929) S*laetipes* (Zetterstedt, 1850) S*lauderborni* (Kieffer, 1909) M S*penicillatus* (Goetghebuer, 1928) S*tenuis* (Meigen, 1830) M*Rheotanytarsus* Thienemann & Bause, 1913*curvistylus* (Goetghebuer, 1921) S*distinctissimus* (Brundin, 1947) S*musciicola* Thienemann, 1929 S*nigricauda* Fittkau, 1960 S*pentapoda* (Kieffer, 1909) S*photophilus* (Goetghebuer, 1921) S*reissi* Lehmann, 1970 S*rhenanus* Klink, 1983 S*Stempellina* Thienemann & Bause, 1913*bauseli* (Kieffer, 1911) M S*Stempellinella* Brundin, 1947*brevis* (Edwards, 1929) B S*minor* (Edwards, 1929) S*Tanytarsus* van der Wulp, 1874*aberrans* Lindeberg, 1970 S*bathophilus* Kieffer, 1911 S*brundini* Lindberg, 1963 S*chinyensis* Goetghebuer, 1934 S*curticornis* Kieffer, 1911 M S*ejuncidus* (Walker, 1856) S*eminulus* (Walker, 1856) S*gregarius* Kieffer, 1909 B M S*heusdensis* Goetghebuer, 1923 S*lestagei* Goetghebuer, 1922 S*medius* Reiss & Fittkau, 1971 S*mendax* Kieffer, 1925 S*= holochlorus* Edwards, 1929*palettaris* Verneaux, 1969 S*pallidicornis* (Walker, 1856) S*signatus* (van der Wulp, 1858) S*usmaensis* Pagast, 1931 M*Thienemanniola* Kieffer, 1929*ploenensis* Kieffer, 1921 B*Zavrelia* Kieffer, 1913*pentatoma* Kieffer, 1913 B

CERATOPOGONIDAE

Ivan Országh, Jan Knoz & Josef Chalupský

CERATOPOGONINAE

CULICOIDINI

Culicoides Latreille, 1809sg. *Avaritia* Fox, 1955*abchazicus* Dzhafarov, 1964 B S*chioterus* (Meigen, 1830) B M S*dewulfi* Goetghebuer, 1936 M S*obsoletus* (Meigen, 1818) B M S*scoticus* Downes & Kettle, 1952 B M Ssg. *Beltranmyia* Vargas, 1953*circumscripatus* Kieffer, 1918 B M S*manchuriensis* Tokunaga, 1941 B M S*salinarius* Kieffer, 1914 M S*sphagnumensis* Williams, 1955 Bsg. *Culicoides* s. str.*delta* Edwards, 1939 B M S*fagineus* Edwards, 1939 B M S*griseus* Edwards, 1939 B M S*impunctatus* Goetghebuer, 1920 B M S*newsteadi* Austen, 1921 S*= halophilus* Kieffer, 1924*pulicaris* (Linnaeus, 1758) B M S*punctatus* (Meigen, 1804) B M Ssg. *Monoculicoides* Khalaf, 1954*longicollis* Glukhova, 1971 S*mubeculosus* (Meigen, 1830) B M S*parroti* Kieffer, 1922 B S*puncticollis* (Becker, 1903) S*riethi* Kieffer, 1914 M S*stigma* (Meigen, 1818) B M Ssg. *Oecacta* Poey, 1851*achrayi* Kettle & Lawson, 1955 B M S

<i>albicans</i> (Winnertz, 1852)	B	M	S	<i>Ceratoculicoides</i> Wirth & Ratanaworabhan,	
<i>alazanicus</i> Dzhanfarov, 1961		M	S	1971	
= <i>musilator</i> Krömer & Callot, 1961				<i>moravicus</i> Knoz, 1987	M
<i>brunnicans</i> Edwards, 1939	B		S	<i>Alluaudomyia</i> Kieffer, 1913	
<i>cameroni</i> Campbell & Pelham-Clinton, 1960			S	<i>falcata</i> Knoz & Ratajský, 1987	M
<i>comosioculatus</i> Tokunaga, 1956	B	M	S	<i>needhami</i> Thomsen, 1935	M
<i>dispersus</i> Gutsevich & Smatov, 1966			S	<i>splendida</i> (Winnertz, 1852)	B M S
<i>duddingstoni</i> Kettle & Lawson, 1955		M	S	STILOBEZZIINI	
<i>dzhanfarovi</i> Remm, 1967		M	S	<i>Kolenohoelea</i> de Meillon & Wirth, 1981	
<i>fascipennis</i> (Staeger, 1839)	B	M	S	<i>calcarata</i> (Goetghebuer, 1920)	B M
<i>festivipennis</i> Kieffer, 1914	B	M	S	= <i>sharpi</i> (Edwards, 1929)	
= <i>odibilis</i> Austen, 1921				<i>Stilobezzia</i> Kieffer, 1911	
<i>furcillatus</i> Callot, Krömer & Paradis, 1962			S	sg. <i>Acanthohoelea</i> Kieffer, 1917	
<i>haranti</i> Rioux, Descous & Pech, 1959		M		<i>gracilis</i> (Haliday, 1833)	M S
<i>heliophilus</i> Edwards, 1921	B	M	S	<i>ochracea</i> (Winnertz, 1852)	M S
<i>jurensis</i> Callot, Krömer & Deduit, 1962		M	S	sg. <i>Stilobezzia</i> s. str.	
<i>kibunensis</i> Tokunaga, 1937	B	M	S	<i>antennalis</i> (Coquillett, 1901)	M
= <i>cubitalis</i> Edwards, 1939				<i>flavivirostris</i> (Winnertz, 1852)	B M
<i>maritimus</i> Kieffer, 1924	B	M	S	<i>Monohelea</i> Kieffer, 1917	
<i>minutissimus</i> (Zetterstedt, 1855)	B	M	S	sg. <i>Allohelea</i> Kieffer, 1917	
<i>pallidicornis</i> Kieffer, 1919	B	M	S	<i>tesselata</i> (Zetterstedt, 1850)	S
<i>pictipennis</i> (Staeger, 1939)	B	M	S	sg. <i>Monohelea</i> s. str.	
<i>pseudoheliophilus</i> Callot & Krömer, 1961			S	<i>estonica</i> Remm, 1965	B
<i>reconditus</i> Campbell & Pelham-Clinton, 1960	B	M	S	sg. <i>Schizohoelea</i> Kieffer, 1917	
<i>segnis</i> Campbell & Pelham-Clinton, 1960	B	M	S	<i>leucopeza</i> (Meigen, 1804)	B S
<i>semimaculatus</i> Clastrier, 1958		M	S	<i>Serromyia</i> Meigen, 1818	
<i>shaklawensis</i> Khalaf, 1957		M	S	<i>atra</i> (Meigen, 1818)	B S
<i>simulator</i> Edwards, 1939		M	S	= <i>spinosipes</i> Kieffer, 1919	
<i>subfascipennis</i> Kieffer, 1919	B	M	S	<i>femorata</i> (Meigen, 1804)	B M S
<i>truncorum</i> Edwards, 1939		M	S	<i>ledicola</i> Kieffer, 1925	M
= <i>sylvarum</i> Callot & Krömer, 1961				<i>morio</i> (Fabricius, 1775)	B S
<i>ustini</i> Shevchenko, 1962	B	M	S	<i>rufitarsis</i> (Meigen, 1818)	B S
<i>vexans</i> (Staeger, 1839)	B	M	S	= <i>dipetala</i> Remm, 1965	
<i>vidourensis</i> Krömer, Molet & Bach, 1968		M	S	SPHAEROMIINI	
<i>zhogolevi</i> Remm, 1968			S	<i>Macropeza</i> Meigen, 1818	
sg. <i>Pontoculicoides</i> Remm, 1968				<i>albitarsis</i> Meigen, 1818	B S
<i>saevus</i> Kieffer, 1922			S	<i>Probezzia</i> Kieffer, 1906	
<i>slovacus</i> Országh, 1969			S	<i>seminigra</i> (Panzer, 1796)	M S
<i>tauricus</i> Gutsevich, 1959		M	S	<i>Mallochohelea</i> Wirth, 1962	
				<i>inermis</i> (Kieffer, 1909)	B M
CERATOPOGONINI				<i>nitida</i> (Macquart, 1826)	M S
<i>Brachypogon</i> Kieffer, 1899				<i>remota</i> (Kieffer, 1919)	S
sg. <i>Brachypogon</i> s. str.				<i>setigera</i> (Loew, 1864)	B M
<i>vitiosus</i> (Winnertz, 1852)			S	<i>Sphaeromyia</i> Curtis, 1829	
sg. <i>Isohelea</i> Kieffer, 1917				<i>fasciatus</i> (Meigen, 1804)	B M S
<i>nitidulus</i> (Edwards, 1921)	B			<i>miricornis</i> (Kieffer, 1919)	B M S
<i>perpusillus</i> (Edwards, 1921)	B			<i>pictus</i> (Meigen, 1818)	B M S
<i>sociabilis</i> (Goetghebuer, 1920)	B			PALPOMYIINI	
<i>Ceratopogon</i> Meigen, 1803				<i>Bezzia</i> Kieffer, 1899	
<i>communis</i> Meigen, 1804			S	sg. <i>Bezzia</i> s. str.	
<i>crassinervis</i> (Goetghebuer, 1920)	B			<i>coracina</i> (Zetterstedt, 1850)	B
<i>grandiforceps</i> (Kieffer, 1913)	B	M		= <i>albipes</i> (Winnertz, 1852)	
<i>lacteipennis</i> Zetterstedt, 1838	B			<i>flavicornis</i> (Staeger, 1839)	B M S
<i>niveipennis</i> Meigen, 1804	B		S		

<i>flavipluma</i> Kieffer, 1919					
<i>gracilipes</i> (Winnertz, 1852)			S		
<i>nobilis</i> (Winnertz, 1852)	B	M			
<i>ornata</i> (Meigen, 1830)	B	M	S		
<i>winnertziana</i> Kieffer, 1919		M			
sg. <i>Homobezzia</i> Macfie, 1932					
<i>annulipes</i> (Meigen, 1830)	B	M			
<i>bicolor</i> (Meigen, 1804)	B	M			
<i>bohémica</i> Kieffer, 1919	B				
<i>circumdata</i> (Staeger, 1839)	B				
– <i>solstitialis</i> (Winnertz, 1852)					
<i>xanthocephala</i> Goetghebuer, 1911		M			
sg. <i>Pygobezzia</i> Remm, 1974					
<i>albicornis</i> (Meigen, 1818)	B	M	S		
<i>Phaenobezzia</i> Haeselbarth, 1965					
<i>rubiginosa</i> (Winnertz, 1852)		M			
<i>Palpomyia</i> Meigen, 1818					
sg. <i>Gluhhovia</i> Remm, 1976					
<i>miki</i> Goetghebuer, 1934			S		
sg. <i>Palpomyia</i> s. str.					
<i>armata</i> Kieffer, 1921		M			
<i>brachialis</i> (Haliday, 1833)		M	S		
– <i>longipennis</i> Kieffer, 1919					
<i>distincta</i> (Haliday, 1833)	B	M			
<i>flavipes</i> (Meigen, 1804)	B	M	S		
<i>fulva</i> (Macquart, 1826)	B	M			
<i>infuscata</i> Kieffer, 1919			S		
<i>lineata</i> (Meigen, 1804)	B		S		
<i>nigradorsum</i> Kieffer, 1919		M	S		
<i>nigripes</i> Kieffer, 1919			S		
<i>nigripes</i> (Meigen, 1830)	B				
<i>praeusta</i> (Loew, 1869)			S		
<i>rufipes</i> (Meigen, 1818)			S		
<i>semiermis</i> Goetghebuer, 1914		M			
<i>serripes</i> (Meigen, 1818)		M	S		
<i>spinipes</i> (Panzer, 1806)			S		
<i>tibialis</i> (Meigen, 1818)			S		
<i>untipennis</i> Kieffer, 1919		M			
DASYHELEINAE					
<i>Dasyhelea</i> Kieffer, 1911					
sg. <i>Dasyhelea</i> s. str.					
<i>dufouri</i> (Laboulbène, 1869)	B	M	S		
<i>lignicola</i> Kieffer, 1919		M			
<i>saxicola</i> (Edwards, 1929)	B	M			
<i>zavreli</i> Kieffer, 1918	B	M			
sg. <i>Dicryptoscena</i> Enderlein, 1936					
<i>inclusa</i> Kieffer, 1918	B				
<i>modesta</i> (Winnertz, 1852)		M			
<i>semistriata</i> Goetghebuer, 1921			S		
sg. <i>Prokempia</i> Kieffer, 1913					
<i>flaviventris</i> (Goetghebuer, 1910)		M			
sg. <i>Pseudoculicoides</i> Malloch, 1915					
<i>punctiventris</i> Goetghebuer, 1940	B	M			
<i>turficola</i> Kieffer, 1925		M			
sg. <i>Sebessia</i> Remm, 1979					
<i>acuminata</i> Kieffer, 1919		M			
FORCIPOMYIINAE					
<i>Atrichopogon</i> Kieffer, 1906					
sg. <i>Atrichopogon</i> s. str.					
<i>brunnipes</i> (Meigen, 1804)				B	M S
<i>fuscus</i> (Meigen, 1804)				B	M S
<i>griseolus</i> (Zetterstedt, 1855)				B	M
<i>hirtidorsum</i> Remm, 1961				B	M
<i>infuscus</i> Goetghebuer, 1929				B	M S
<i>minutus</i> (Meigen, 1830)				B	M S
<i>pavidus</i> (Winnertz, 1852)					M S
<i>tatricus</i> Remm, 1981					S
<i>thienemanni</i> Kieffer, 1921					S
<i>transversalis</i> Kieffer, 1918					S
<i>unilineatus</i> Remm, 1967					M
sg. <i>Lophomyidium</i> Cordero, 1929					
<i>rostratus</i> (Winnertz, 1852)				B	S
sg. <i>Meloehalea</i> Wirth, 1956					
<i>lucorum</i> (Meigen, 1818)				B	M S
<i>meloesugans</i> Kieffer, 1922					S
<i>oedemerarum</i> Storå, 1939				B	M S
<i>setosipennis</i> (Kieffer, 1911)				B	M
sg. <i>Psammopogon</i> Remm, 1979					
<i>albiscapula</i> Kieffer, 1918					S
<i>flaveolus</i> Zilahi-Sebess, 1936					S
<i>trifasciatus</i> Kieffer, 1918					M S
sg. <i>Psilokempia</i> Enderlein, 1936					
<i>appendiculatus</i> (Goetghebuer, 1920)					S
<i>maculatus</i> (Lundström, 1910)					S
– <i>avasiensis</i> Remm, 1959					
<i>forcipatus</i> (Winnertz, 1852)				B	S
<i>Forcipomyia</i> Meigen, 1818					
sg. <i>Caloforcipomyia</i> Saunders, 1956					
<i>glauca</i> Macfie, 1934				B	
sg. <i>Euprojoannisia</i> Brèthes, 1914					
<i>alacris</i> (Winnertz, 1852)				B	M S
<i>hirtipennis</i> (Malloch, 1915)				B	M
<i>titillans</i> (Winnertz, 1852)				B	M S
sg. <i>Forcipomyia</i> s. str.					
<i>altaica</i> Remm, 1972					M
<i>apricans</i> (Kieffer, 1919)					S
<i>bipunctata</i> (Linnaeus, 1767)				B	M S
<i>brevipennis</i> (Macquart, 1828)				B	M S
<i>ciliata</i> (Winnertz, 1852)				B	M S
<i>crassipes</i> (Winnertz, 1852)					M S
<i>hygrophila</i> Kieffer, 1925				B	M S
<i>kaltenbachii</i> (Winnertz, 1852)				B	M
<i>nigra</i> (Winnertz, 1852)				B	M
<i>nigrans</i> Remm, 1962					M
<i>pallida</i> (Winnertz, 1852)					M S
<i>picea</i> (Winnertz, 1852)					M S
<i>pulchrithorax</i> Edwards, 1924				B	M S
– <i>trilineata</i> Goetghebuer, 1934					
<i>sahariensis</i> Kieffer, 1923					M S
<i>squamigera</i> Kieffer, 1916					S
<i>temuis</i> (Winnertz, 1852)				B	M S
<i>temuisquama</i> Kieffer, 1924					M S
<i>tuzeti</i> W. & N. Huttel, 1952				B	M

sg. <i>Lasiohelea</i> Kieffer, 1921				<i>lundstromi</i> (Enderlein, 1921)	B	M	S
<i>sibirica</i> (Bujanova, 1962)		M		= <i>latigonium</i> (Rubtsov, 1956)			
<i>velox</i> (Winnertz, 1852)	B	M	S	<i>oligotuberculatum</i> (Knoz, 1965)		M	S
sg. <i>Microhelea</i> Kieffer, 1917				<i>vernum</i> (Macquart, 1838)	B	M	S
<i>albostylus</i> Remm, 1979		M		= <i>latipes</i> auct. nec Meigen, 1804			
<i>fuliginosa</i> (Meigen, 1818)	B	M	S	sg. <i>Obuchovia</i> Rubtsov, 1947			
<i>longisetosa</i> Krivosheina & Remm, 1973		M		<i>auricoma</i> (Meigen, 1818)		M	S
<i>rustica</i> (Kieffer, 1919)		M	S	sg. <i>Schoenbaueria</i> Enderlein, 1921			
<i>suberis</i> Clastrier, 1956		M		<i>pusillum</i> Fries, 1824		M	
sg. <i>Synthridomyia</i> Saunders, 1956				sg. <i>Simulium</i> s. str.			
<i>acidicola</i> (Tokunaga, 1937)	B			<i>argenteostriatum</i> Strobl, 1898	B	M	S
<i>murina</i> (Winnertz, 1852)		M	S	<i>argyreatum</i> Meigen, 1838	B	M	S
sg. <i>Thyridomyia</i> Saunders, 1925				= <i>rheophilum</i> (Knoz, 1961)			
<i>frutetorum</i> (Winnertz, 1852)			S	<i>haracorne</i> Smart, 1944			S
<i>microcera</i> (Kieffer, 1919)			S	<i>colombaschense</i> (Fabricius, 1787)			S
<i>monilicornis</i> (Coquillett, 1905)		M	S	<i>degrangei</i> Dorier & Grenier, 1960			S
sg. <i>Trichohalea</i> Goetghebuer, 1920				<i>maximum</i> (Knoz, 1961)	B	M	S
<i>eques</i> (Johannsen, 1908)	B		S	<i>monticola</i> Friederichs, 1920	B	M	S
<i>tonnoiri</i> (Goetghebuer, 1920)	B		S	<i>morsitans</i> Edwards, 1915	B	M	S

SIMULIIDAE

Jan Knoz & Ladislav Jedlička

SIMULIINAE

PROSIMULIINI

Prosimulium Roubaud, 1906

<i>fulvipes</i> (Edwards, 1921)			S	= <i>argyreatum</i> auct. nec Meigen, 1838			
<i>hirtipes</i> (Fries, 1824)	B	M	S	<i>ornatum</i> Meigen, 1818	B	M	S
<i>latimicro</i> (Enderlein, 1925)	B	M	S	<i>paramorsitans</i> Rubtsov, 1956			S
<i>rufipes</i> (Meigen, 1830)	B	M	S	<i>posticatum</i> Meigen, 1838	B	M	
<i>subrufipes</i> Knoz, 1980		M		= <i>austeni</i> Edwards, 1915			
<i>tomosvaryi</i> (Enderlein, 1921)	B	M	S	<i>reptans</i> (Linnaeus, 1758)	B	M	S
<i>Twinnia</i> Stone & Jamnback, 1955				<i>reptantoides</i> Carlsson, 1962	B	M	S
<i>hydroides</i> Novák, 1956	B	M	S	= <i>galeratum</i> auct. nec Edwards, 1920			

SIMULIINI

Simulium Latreille, 1802

sg. <i>Boophthora</i> Enderlein, 1921				<i>rostratum</i> (Lundström, 1911)	B	M	S
<i>erythrocephalum</i> (De Geer, 1776)	B	M	S	= <i>verecundum</i> auct. nec Stone & Jamnback, 1955			
sg. <i>Eusimulium</i> Roubaud, 1906				<i>trifasciatum</i> Curtis, 1839	B	M	S
<i>angustipes</i> Edwards, 1915		M	S	= <i>spinosum</i> Doby & Deblock, 1957			
= <i>securiforme</i> (Rubtsov, 1956)				<i>tuberosum</i> (Lundström, 1911)	B	M	S
<i>aureum</i> (Fries, 1824)	B	M	S	<i>variegatum</i> Meigen, 1818	B	M	S
<i>velutinum</i> (Santos Abreu, 1929)	B	M	S	<i>voilense</i> Serban, 1960			S
= <i>serbicum</i> Baranov, 1925				<i>vulgare</i> Dorogostaisky, Rubtsov			

sg. *Nevermannia* Enderlein, 1921

<i>angustatum</i> (Rubtsov, 1956)		M	S	& Vlasenko, 1935		M	S
<i>angustitarse</i> (Lundström, 1911)	B	M	S	sg. <i>Wilhelmia</i> Enderlein, 1921			
<i>bertrandi</i> (Grenier & Dorier, 1959)	B	M	S	<i>balcanicum</i> (Enderlein, 1924)			S
<i>brevicens</i> (Rubtsov, 1956)	B	M	S	<i>equinum</i> (Linnaeus, 1758)	B	M	S
<i>carpathicum</i> (Knoz, 1961)	B	M	S	<i>lineatum</i> (Meigen, 1804)	B	M	S
<i>carthusiense</i> (Grenier & Dorier, 1959)		M	S	<i>pseudequinum</i> Séguy, 1921			S
<i>codreanui</i> (Serban, 1958)		M	S	= <i>mediterraneum</i> Puri, 1925			
<i>costatum</i> (Friederichs, 1920)	B	M	S	THAUMALEIDAE			
<i>crenobium</i> (Knoz, 1961)	B	M	S	Jaroslav Martinovský			
<i>cryophilum</i> (Rubtsov, 1959)	B	M	S	<i>Androprosopa</i> Mik, 1898			

THAUMALEIDAE

Jaroslav Martinovský

<i>larvata</i> (Mik, 1888)	B	M	S	<i>nigra</i> (Loew, 1871)			S
<i>Orphnephilina</i> Enderlein, 1936				<i>Thaumalea</i> Ruthé, 1831			
<i>aperta</i> Martinovský & Rozkošný, 1976		M	S	<i>austriaca</i> Edwards, 1929		M	S
<i>bezzii</i> Edwards, 1929	B	M	S	<i>caudata</i> Bezzi, 1913		M	S

<i>decussiferens</i> Vaillant, 1969	S	<i>latipennis</i> (Loew, 1856)	B M S
<i>dentata</i> Edwards, 1929	M	<i>lineola</i> Fabricius, 1794	B M S
<i>miki</i> Edwards, 1929	M S	<i>maculatus</i> (De Geer, 1776)	B M S
<i>remota</i> Martinovský & Rozkošný, 1976	M S	<i>notatus</i> (Meigen, 1820)	B M S
<i>tatrica</i> Vaillant, 1969	S	<i>scolopaceus</i> (Linnaeus, 1758)	B M S
<i>testacea</i> Ruthé, 1831	B M S	<i>strigosus</i> Meigen, 1804	B M S
<i>truncata</i> Edwards, 1929	B M	<i>tringarius</i> (Linnaeus, 1758)	B M S
- <i>cebenica</i> Vaillant, 1977		<i>vitripennis</i> (Meigen, 1820)	B M S
<i>vaillanti</i> Martinovský & Rozkošný, 1976	M S	<i>Symphoromyia</i> Frauenfeld, 1867	
		<i>crassicornis</i> (Panzer, 1806)	B M S
		<i>immaculata</i> (Meigen, 1804)	B M
		<i>melaena</i> (Meigen, 1820)	B M S
XYLOPHAGIDAE			
Rudolf Rozkošný			
<i>Xylophagus</i> Meigen, 1803		SPANIINAE	
<i>ater</i> Meigen, 1803	B M S	<i>Ptiolina</i> Zetterstedt, 1842	
<i>cinctus</i> (De Geer, 1776)	B M S	<i>obscura</i> (Fallén, 1814)	B M S
<i>compeditus</i> Meigen, 1820	B M S	<i>paradoxa</i> Jaennicke, 1866	S
<i>junki</i> Szilády, 1932	B M	<i>Spania</i> Meigen, 1830	
		<i>nigra</i> Meigen, 1830	B M S
COENOMYIIDAE			
Rudolf Rozkošný			
<i>Coenomyia</i> Latreille, 1796		TABANIDAE	
<i>ferruginea</i> (Scopoli, 1763)	B M S	Milan Chvála	
		CHRYSOPSINAE	
		<i>Chrysops</i> Meigen, 1803	
		<i>caecutiens</i> (Linnaeus, 1758)	B M S
		<i>divaricatus</i> Loew, 1858	B S
		<i>flavipes</i> Meigen, 1804	M S
		<i>parallelogrammus</i> Zeller, 1842	B S
		<i>relictus</i> Meigen, 1820	B M S
		<i>rufipes</i> Meigen, 1820	B M S
		<i>sepulcralis</i> (Fabricius, 1794)	B
		<i>viduatus</i> (Fabricius, 1794)	B M S
		<i>Silvius</i> Meigen, 1820	
		<i>alpinus</i> (Scopoli, 1763)	B M S
ATHERICIDAE			
Rudolf Rozkošný & Karel Spitzer			
<i>Atherix</i> Meigen, 1803		TABANINAE	
<i>ibis</i> (Fabricius, 1798)	B M S	TABANINI	
<i>Atrichops</i> Verrall, 1909		<i>Atylotus</i> Osten-Sacken, 1876	
<i>crassipes</i> (Meigen, 1820)	S	<i>fulvus</i> (Meigen, 1820)	B M S
<i>Ihisia</i> Rondani, 1856		<i>latistriatus</i> (Brauer, 1880)	B
<i>marginata</i> (Fabricius, 1781)	B M S	<i>loewianus</i> (Villeneuve, 1920)	B M S
		<i>plebejus</i> (Fallén, 1817)	B M S
		<i>rusticus</i> (Linnaeus, 1767)	B M S
		<i>sublunaticornis</i> (Zetterstedt, 1842)	B
RHAGIONIDAE			
Rudolf Rozkošný & Karel Spitzer			
		<i>Glaucops</i> Szilády, 1923	
		<i>hirsutus</i> (Villers, 1789)	B M
		<i>Hybomitra</i> Enderlein, 1922	
		<i>acuminata</i> (Loew, 1858)	S
		<i>arpadi</i> (Szilády, 1923)	B M
		<i>auripila</i> (Meigen, 1820)	B M S
		<i>bimaculata</i> (Macquart, 1826)	B M S
		<i>borealis</i> (Fabricius, 1781)	B M
		<i>ciureai</i> (Séguy, 1937)	B M S
		<i>distinguenda</i> (Verrall, 1909)	B M S
		<i>kaurii</i> Chvála & Lyneborg, 1970	B M S
</			

<i>lundbecki</i> Lyneborg, 1959	B M S	<i>Xylomya</i> Rondani, 1861	
<i>lurida</i> (Fallén, 1817)	B M S	<i>maculata</i> (Meigen, 1804)	M S
<i>micans</i> (Meigen, 1804)	B M S	<i>varia</i> (Meigen, 1820)	B M
<i>montana</i> (Meigen, 1820)	B M S		
<i>muehlfeldi</i> (Brauer, 1880)	B M S	STRATIOMYIDAE	
<i>nigricornis</i> (Zetterstedt, 1842)	B	Rudolf Rozkošný	
<i>nitidifrons confiformis</i> Chvála & Moucha, 1971	B M S		
<i>pilosa</i> (Loew, 1858)	M S	BERIDINAE	
<i>tropica</i> (Linnaeus, 1758)	B M S	<i>Actina</i> Meigen, 1804	
Tabanus Linnaeus, 1758		<i>chalybea</i> Meigen, 1804	B M S
<i>autumnalis</i> Linnaeus, 1761	B M S	<i>Allognosta</i> Osten-Sacken, 1883	
<i>bifarius</i> Loew, 1858	S	<i>vagans</i> (Loew, 1873)	M S
<i>bovinus</i> Linnaeus, 1758	B M S	<i>Beris</i> Latreille, 1802	
<i>briani</i> Leclercq, 1962	S	<i>chalybata</i> (Forster, 1771)	B M S
<i>bromius</i> Linnaeus, 1758	B M S	<i>clavipes</i> (Linnaeus, 1767)	B M S
<i>cordiger</i> Meigen, 1820	B M S	<i>geniculata</i> Curtis, 1830	B M S
<i>exclusus</i> Pandellé, 1883	S	<i>fuscipes</i> Meigen, 1820	B M S
<i>glaucopis</i> Meigen, 1820	B M S	<i>morrisii</i> Dale, 1841	B M S
<i>maculicornis</i> Zetterstedt, 1842	B M S	<i>strobli</i> Dušek & Rozkošný, 1968	B M S
<i>miki</i> Brauer, 1880	B M S	<i>vallata</i> (Forster, 1771)	B M S
<i>paradoxus</i> Jaennicke, 1866	S	<i>Chorisops</i> Rondani, 1856	
<i>quatuornotatus</i> Meigen, 1820	B M S	<i>nagatomii</i> Rozkošný, 1979	M
<i>regularis</i> Jaennicke, 1866	S	<i>tibialis</i> (Meigen, 1820)	B M S
<i>spodopterus</i> Meigen, 1820	B M S	<i>Exodontha</i> Rondani, 1856	
<i>sudeticus</i> Zeller, 1842	B M S	<i>dubia</i> (Zetterstedt, 1838)	B S
<i>temicornis</i> (Enderlein, 1932)	M S		
<i>tergestinus</i> Egger, 1859	B M S	SARGINAE	
<i>unifasciatus</i> Loew, 1858	S	<i>Chloromyia</i> Duncan, 1837	
Therioplectes Zeller, 1842		<i>formosa</i> (Scopoli, 1763)	B M S
<i>gigas</i> (Herbst, 1787)	M S	<i>speciosa</i> (Macquart, 1834)	B M S
		<i>Microchrysa</i> Loew, 1855	
DIACHLORINI		<i>cyaneiventris</i> (Zetterstedt, 1842)	B M S
<i>Philipomyia</i> Olsufjev, 1964		<i>flavicornis</i> (Meigen, 1822)	B M S
<i>aprica</i> (Meigen, 1820)	B M S	<i>polita</i> (Linnaeus, 1758)	B M S
<i>graeca</i> (Fabricius, 1794)	B M S	<i>Sargus</i> Fabricius, 1798	
		<i>bipunctatus</i> (Scopoli, 1763)	B M S
HAEMATOPOTINI		<i>euprarius</i> (Linnaeus, 1758)	B M S
<i>Haematopota</i> Meigen, 1803		<i>flavipes</i> Meigen, 1822	B M S
<i>bigoti</i> Gobert, 1881	S	<i>iridatus</i> (Scopoli, 1763)	B M S
<i>crassicornis</i> Wahlberg, 1848	B M S		
<i>csikii</i> Szilády, 1922	S	STRATIOMYINAE	
<i>grandis</i> Meigen, 1820	M S	<i>Odontomyia</i> Meigen, 1803	
<i>italica</i> Meigen, 1804	B M S	<i>angulata</i> (Panzer, 1798)	B M S
<i>pluvialis</i> (Linnaeus, 1758)	B M S	<i>annulata</i> (Meigen, 1822)	M S
<i>scutellata</i> (Olsufjev, Moucha & Chvála, 1964)	B M S	<i>argentata</i> (Fabricius, 1794)	B M S
<i>subcylindrica</i> Pandellé, 1883	B M S	<i>flavissima</i> (Rossi, 1790)	B M S
Heptatoma Meigen, 1803		<i>hydroleon</i> (Linnaeus, 1758)	B M S
<i>pellucens</i> (Fabricius, 1776)	B M S	<i>microleon</i> (Linnaeus, 1758)	B
		<i>ornata</i> (Linnaeus, 1758)	B M S
XYLOMYIDAE		<i>tigrina</i> (Fabricius, 1775)	B M S
Rudolf Rozkošný		<i>Oplodontha</i> Rondani, 1864	
		<i>viridula</i> (Fabricius, 1775)	B M S
<i>Solva</i> Walker, 1859		<i>Stratiomys</i> Geoffroy, 1762	
<i>marginata</i> (Meigen, 1820)	B M S	<i>cenisia</i> Meigen, 1822	M S
		<i>chamaeleon</i> (Linnaeus, 1758)	B M S
		<i>equestris</i> Meigen, 1838	B M S

<i>longicornis</i> (Scopoli, 1763)	B M S	<i>gibbosus</i> (Linnaeus, 1778)	M S
<i>potamida</i> Meigen, 1822	B M S	<i>obscuripes</i> Chvála, 1980	B S
<i>ruficornis</i> (Macquart, 1838)	M S	<i>pallipes</i> Latreille, 1811	B M S
<i>singularior</i> (Harris, 1776)	B M S	<i>reginae</i> Trojan, 1956	S
		<i>varius</i> Latreille, 1811	S
		<i>zonatus</i> Erichson, 1840	B S
CLITELLARIINAE			
<i>Adoxomyia</i> Bezzi, 1908			
<i>lindneri</i> Dušek & Rozkošný, 1963	S		
<i>Clitellaria</i> Meigen, 1803		MYTHICOMYIIDAE	
<i>ephippium</i> (Fabricius, 1775)	B M S	Jaroslav Bosák	
<i>Lasiopa</i> Brullé, 1832			
<i>calva</i> (Meigen, 1822)	M S	<i>Glbellula</i> Bezzi, 1902	
<i>villosa</i> (Fabricius, 1794)	B M S	<i>arctica</i> (Zetterstedt, 1838)	B M S
<i>Nemotelus</i> Geoffroy, 1762			
<i>brevirostris</i> Meigen, 1822	B S		
<i>nigrinus</i> Fallén, 817	B M S	PHTHIRIIDAE	
<i>pantherinus</i> (Linnaeus, 1758)	B M S	Jaroslav Bosák	
<i>signatus</i> Frivaldsky in Schiner, 1855	M S		
<i>uliginosus</i> (Linnaeus, 1767)	B M S	<i>Phthiria</i> Meigen, 1803	
<i>Oxycera</i> Meigen, 1803		<i>canescens</i> Loew, 1846	M
<i>analís</i> Meigen, 1822	M S	<i>gaedei</i> Wiedemann in Meigen, 1820	M
<i>dives</i> Loew, 1845	B S	<i>minuta</i> (Fabricius, 1805)	B
<i>falleni</i> Staeger, 1844	B M S	<i>pulicaria</i> (Mikan, 1796)	B M S
<i>leonina</i> (Panzer, 1798)	B M S		
<i>meigenii</i> Staeger, 1844	B M S		
<i>morristi</i> Curtis, 1830	S	BOMBYLIIDAE	
<i>nigricornis</i> Olivier, 1812	M S	Jaroslav Bosák	
<i>pardalina</i> Meigen, 1822	B M S		
<i>pseudoamoena</i> Dušek & Rozkošný, 1974	S	LOMATIINAE	
<i>pygmaea</i> (Fallén, 1817)	B M S	<i>Lomatia</i> Meigen, 1822	
<i>rara</i> (Scopoli, 1763)	B M S	<i>erinnys</i> Loew, 1869	S
<i>terminata</i> Meigen, 1822	B M S	<i>lachesis</i> Egger, 1859	M
<i>trilineata</i> (Linnaeus, 1767)	B M S	<i>lateralis</i> (Meigen, 1820)	B M S
		<i>sabaea</i> (Fabricius, 1781)	M S
PACHYGASTERINAE			
<i>Berkshiria</i> Johnson, 1914		ANTHRACINAE	
<i>hungarica</i> (Kertész, 1921)	M	<i>Anthrax</i> Scopoli, 1763	
<i>Eupachygaster</i> Kertész, 1911		<i>anthrax</i> (Schrank, 1781)	B M S
<i>tarsalis</i> (Zetterstedt, 1842)	B M S	<i>binotatus</i> Wiedemann in Meigen, 1820	B M
<i>Neopachygaster</i> Austen, 1901		<i>trifasciatus</i> Meigen, 1804	B M S
<i>meromelaena</i> (Dufour, 1841)	B M S	<i>varius</i> Fabricius, 1794	B M S
<i>Pachygaster</i> Meigen, 1803		<i>Spogostylum</i> Macquart, 1840	
<i>atra</i> (Panzer, 1798)	B M S	<i>aethiops</i> (Fabricius, 1781)	M S
<i>leachi</i> (Curtis, 1824)	B M S		
<i>Zabrachia</i> Coquillett, 1901		EXOPROSOPINAE	
<i>minutissima</i> (Zetterstedt, 1838)	B M	VILLINI	
<i>tenella</i> (Jaenicke, 1866)	B M	<i>Hemipenthes</i> Loew, 1869	
		<i>maurus</i> (Linnaeus, 1758)	B M S
ACROCERIDAE		<i>morio</i> (Linnaeus, 1758)	B M S
Milan Chvála		<i>velutinus</i> (Meigen, 1820)	B S
		<i>Villa</i> Lioy, 1864	
<i>Acrocera</i> Meigen, 1803		<i>cingulata</i> (Meigen, 1804)	B M S
<i>orbicula</i> (Fabricius, 1787)	B M	<i>circumdata</i> (Meigen, 1820)	B S
<i>sanguinea</i> Meigen, 1804	B	<i>claripennis</i> (Kowarz, 1867)	S
<i>Ogcodes</i> Latreille, 1796		<i>deludens</i> François, 1966	B M S
<i>fumatus</i> Erichson, 1846	B	<i>halteralis</i> (Kowarz, 1883)	B M S
		<i>hottentotta</i> (Linnaeus, 1758)	B M S

<i>humilis</i> (Ruthe, 1831)	B	S	THEREVIDAE		
<i>ixion</i> (Fabricius, 1794)	B	S	Karel Spitzer		
<i>modesta</i> (Meigen, 1820)	B				
<i>occulta</i> (Wiedemann in Meigen, 1820)	B		<i>Acrosanthe</i> Irwin & Lyneborg, 1981		
<i>paniscus</i> (Rossi, 1790)	B	M S	<i>annulata</i> (Fabricius, 1805)	B	M S
<i>quinquefasciata</i> (Wiedemann in Meigen, 1820)	B		<i>Ammothereva</i> Lyneborg, 1984		
			<i>laticornis</i> (Loew, 1856)		M S
			<i>Clorismia</i> Enderlein, 1927		
THYRIDANTHRAXINI			<i>ardea</i> (Fabricius, 1794)	B	M S
<i>Exhyalanthrax</i> Becker, 1916			<i>rustica</i> (Panzer, 1804)	B	M
<i>afer</i> (Fabricius, 1794)	B	M S	<i>Dialineura</i> Rondani, 1856		
<i>Thyridanthrax</i> Osten-Sacken, 1886			<i>anilis</i> (Linnaeus, 1758)	B	M S
<i>fenestratus</i> (Fallén, 1814)	B	M S	<i>Pandivirilia</i> Irwin & Lyneborg		
			<i>caesia</i> (Meigen, 1838)	B	M S
EXOPROSOPINI			<i>eximia</i> (Meigen, 1820)	B	M S
<i>Exoprosopa</i> Macquart, 1840			<i>fuscipennis</i> (Meigen, 1820)	B	M
<i>capucina</i> (Fabricius, 1781)	B	S	<i>melaleuca</i> (Loew, 1847)	B	M
<i>cleomene</i> Egger, 1859	B	M S	<i>Thereva</i> Latreille, 1796		
<i>jaechus</i> (Fabricius, 1805)	B	S	<i>apicalis</i> Wiedemann, 1821		M S
<i>minos</i> (Meigen, 1804)		S	<i>aurata</i> Loew, 1854		S
			<i>bipunctata</i> Meigen, 1820	B	M S
BOMBYLIINAE			<i>brevicornis</i> Loew, 1847		S
CONOPHORINI			<i>cincta</i> Meigen, 1820	B	M S
<i>Conophorus</i> Meigen, 1803			<i>cinifera</i> Meigen, 1830	B	M
<i>virescens</i> (Fabricius, 1787)	B	M S	<i>fulva</i> Meigen, 1804	B	M S
			<i>handlirschi</i> Kröber, 1912	B	M S
DISCHISTINI			<i>innornata</i> Verrall, 1909	B	M S
<i>Bombylisoma</i> Rondani, 1856			<i>lanata</i> Zetterstedt, 1838	B	
<i>minimum</i> (Scopoli, 1772)		S	<i>marginula</i> Meigen, 1820	B	M S
<i>nigriceps</i> (Loew, 1862)		S	<i>microcephala</i> Loew, 1847	B	M S
			<i>nigripes</i> Loew, 1847		M S
BOMBYLIINI			<i>nobilitata</i> (Fabricius, 1775)	B	M S
<i>Anastoechus</i> Osten-Sacken, 1877			<i>oculata</i> Egger, 1859	B	M
<i>nitidulus</i> (Fabricius, 1794)	B	S	<i>plebeja</i> (Linnaeus, 1758)	B	M S
<i>Bombylius</i> Linnaeus, 1758			<i>praecox</i> Egger, 1859	B	M S
<i>ater</i> Scopoli, 1763	B	M S	<i>strigata</i> (Fabricius, 1794)	B	M S
<i>canescens</i> Mikan, 1796	B	M S	<i>valida</i> Loew, 1847	B	M S
<i>cinerascens</i> Mikan, 1796	B	M S			
<i>discolor</i> Mikan, 1796	B	M S			
<i>fimbriatus</i> Meigen, 1820	B	M S			
<i>fulvescens</i> Wiedemann in Meigen, 1820		M S			
<i>major</i> Linnaeus, 1758	B	M S	SCENOPINIDAE		
<i>medius</i> Linnaeus, 1758	B	M S	Jaroslav Bosák		
<i>minor</i> Linnaeus, 1758	B	M S			
<i>rufus</i> Mikan, 1796	B	M S	<i>Scenopinus</i> Latreille, 1802		
<i>pictus</i> Panzer, 1794	B	M S	<i>fenestralis</i> (Linnaeus, 1758)	B	M S
<i>quadrifarius</i> Loew, 1855		M S	<i>glabrifrons</i> Meigen, 1824	B	M S
<i>undatus</i> Mikan, 1796	B	M S	<i>niger</i> (De Geer, 1776)	B	M S
<i>venosus</i> Mikan, 1796	B	M S			
<i>vulpinus</i> Wiedemann in Meigen, 1820	B	M S			
<i>Systoechus</i> Loew, 1855			ASILIDAE		
<i>ctenopterus</i> (Mikan, 1796)	B	M S	Jaroslav Bosák		
HILARIMORPHIDAE			LAPHRIINAE		
Rudolf Rozkošný			LAPHRIINI		
			<i>Choerades</i> Walker, 1851		
<i>Hilarimorpha</i> Schiner, 1860			<i>castellani</i> (Hradský, 1962)		S
<i>singularis</i> Schiner, 1860		S	<i>dioctriaeformis</i> (Meigen, 1820)	B	M S
			<i>fimbriata</i> (Meigen, 1820)	B	M S

<i>fuliginosa</i> (Panzer, 1798)	M S	<i>lateralis</i> (Fallén, 1814)	B M S
<i>gilva</i> (Linnaeus, 1758)	B M S	<i>maculipennis</i> (Macquart, 1834)	B M S
<i>igneae</i> (Meigen, 1820)	B M S	<i>ruficornis</i> (Fabricius, 1794)	B M S
<i>marginata</i> (Linnaeus, 1758)	B M S	Holopogon Loew, 1847	
<i>mouchai</i> Hradský, 1985	S	<i>dimidiatus</i> (Meigen, 1820)	M S
<i>rufipes</i> (Fallén, 1814)	B S	<i>fumipennis</i> (Meigen, 1820)	M S
Laphria Meigen, 1803		<i>nigripennis</i> (Meigen, 1820)	B M S
<i>aurea</i> (Fabricius, 1794)	S	<i>priscus</i> (Meigen, 1820)	M S
<i>empyrea</i> Gerstaecker, 1861	S	Stenopogon Loew, 1847	
<i>ephippium</i> (Fabricius, 1781)	B M S	<i>callosus</i> (Pallas, 1818)	M S
<i>flava</i> (Linnaeus, 1761)	B M S	<i>sabaudus</i> (Fabricius, 1794)	M S
<i>gibbosus</i> (Linnaeus, 1758)	B M S		
		DASYPOGONINAE	
ANDRENOSOMATINI		ISOPOGONINI	
Andrenosoma Rondani, 1856		Leptarthrus Stephens, 1829	
<i>atra</i> (Linnaeus, 1758)	B M S	<i>brevirostris</i> (Meigen, 1804)	B M S
Pogonosoma Rondani, 1856		<i>vitripennis</i> (Meigen, 1820)	S
<i>maroccanum</i> (Fabricius, 1794)	S		
<i>minor</i> Loew, 1869	M S	DASYPOGONINI	
		Dasygogon Meigen, 1803	
LAPHYSTIINAE		<i>diadema</i> (Fabricius, 1781)	B M S
Laphystia Loew, 1847			
<i>sabulicola</i> Loew, 1847	S	LEPTOGASTRINAE	
		Leptogaster Meigen, 1803	
STENOPOGONINAE		<i>cylindrica</i> (De Geer, 1776)	B M S
DIOCTRIINI		<i>guttiventris</i> Zetterstedt, 1842	B M S
Dioctria Meigen, 1803		<i>pubicornis</i> Loew, 1847	S
<i>atricapilla</i> Meigen, 1804	B M S	<i>subtilis</i> Loew, 1847	S
<i>calceata</i> Meigen, 1820	M S		
<i>cothurnata</i> Meigen, 1820	B M S	ASILINAE	
<i>flavipennis</i> Meigen, 1820	B M S	Antipalus Loew, 1849	
<i>harcyniae</i> Loew, 1844	B	<i>similis</i> Moucha & Hradský 1966	S
<i>humeralis</i> Zeller, 1840	B M S	<i>varipes</i> (Meigen, 1820)	B M S
<i>hyalipennis</i> (Fabricius, 1794)	B M S	Antiphrisson Loew, 1849	
<i>lata</i> Loew, 1853	M	<i>trifarius</i> (Loew, 1849)	S
<i>lateralis</i> Meigen, 1804	B M S	Asilus Linnaeus, 1758	
<i>linearis</i> (Fabricius, 1787)	B M S	<i>crabroniformis</i> Linnaeus, 1758	B M S
<i>linurata</i> Loew, 1873	S	Cerdistus Loew, 1849	
<i>longicornis</i> Meigen, 1820	B M S	<i>graminicola</i> Lehr, 1967	S
<i>oelandica</i> (Linnaeus, 1758)	B M S	= <i>albicans oblitus</i> (Tsacas, 1968)	
<i>rufipes</i> (De Geer, 1776)	B M S	Dymachus Loew, 1860	
<i>sudetica</i> Duda, 1940	B M	<i>bifurcus</i> (Loew, 1848)	B M S
<i>wiedemanni</i> Meigen, 1820	M S	<i>bilobus</i> Loew, 1871	S
Molobratia Hull, 1958		<i>bimucronatus</i> (Loew, 1854)	B M S
<i>teutonius</i> (Linnaeus, 1767)	B M S	<i>cephalenus</i> Loew, 1871	S
		<i>cochleatus</i> (Loew, 1854)	B M S
STICHOPOGONINI		<i>fuscipennis</i> (Meigen, 1820)	B M S
Lasipogon Loew, 1847		<i>picipes</i> (Meigen, 1820)	B M S
<i>cinctus</i> (Fabricius, 1781)	B M S	<i>praemorsus</i> (Loew, 1854)	M S
<i>montanus</i> Schiner, 1862	S	<i>stylifer</i> (Loew, 1854)	M S
<i>lichtwardti</i> Oldenberg, 1924	S	<i>trigonus</i> (Meigen, 1804)	B M S
Stichopogon Loew, 1847		Echthistus Loew, 1849	
<i>elegantulus</i> (Wiedemann, 1820)	S	<i>rufinervis</i> (Meigen, 1820)	B M S
<i>scaliger</i> Loew, 1847	S	Erax Scopoli, 1763	
		<i>barbatus</i> Scopoli, 1763	B M S
STENOPOGONINI		Eutolmus Loew, 1848	
Cyrtopogon Loew, 1847		<i>rufibarbis</i> (Meigen, 1820)	M S
<i>flavimanus</i> (Meigen, 1820)	B S		

Machimus Loew, 1849

<i>arthriticus</i> (Zeller, 1840)	M S
<i>annulipes</i> (Brullé, 1832)	M S
<i>chrysitis</i> (Meigen, 1820)	B M S
<i>cyanopus</i> (Loew, 1849)	B M S
<i>gonatistes</i> (Zeller, 1840)	B M S
<i>rusticus</i> (Meigen, 1820)	B M S
<i>senharbis</i> (Loew, 1849)	B M S
Neopitriptus Lehr, 1992	
<i>ninae</i> Lehr, 1992	S
<i>setosulus</i> (Zeller, 1840)	B M S
Neoitamus Osten-Sacken, 1878	
<i>cothurnatus</i> (Meigen, 1820)	B M S
<i>cyanurus</i> (Loew, 1849)	B M S
<i>dasymallus</i> (Gerstaecker, 1861)	S
<i>socius</i> (Loew, 1871)	B M S
Neomochtherus Osten-Sacken, 1878	
<i>hungaricus</i> Engel, 1927	S
<i>flavicornis</i> (Ruthe, 1831)	B M S
<i>flavipes</i> (Meigen, 1820)	M S
<i>pallipes</i> (Meigen, 1820)	B M S
<i>perplexus</i> (Becker, 1923)	S
<i>schineri</i> (Egger, 1855)	S
Pamponerus Loew, 1849	
<i>germanicus</i> (Linnaeus, 1758)	B M S
Paritamus Verrall, 1909	
<i>geniculatus</i> (Meigen, 1820)	B M S
Philonicus Loew, 1849	
<i>albiceps</i> (Meigen, 1820)	B M S
Rhadiurgus Loew, 1849	
<i>variabilis</i> (Zetterstedt, 1838)	B M S
Stilpnogaster Loew, 1849	
<i>aemula</i> (Meigen, 1820)	B M S
Tolmerus Loew, 1849	
<i>atricapillus</i> (Fallén, 1814)	B M S
<i>atripes</i> Loew, 1854	B M S
<i>micans</i> (Meigen, 1820)	M
<i>cingulanus</i> (Fabricius, 1781)	B M S
<i>poecilogaster</i> (Loew, 1849)	B M
<i>pyragra</i> (Zeller, 1840)	B M S
<i>strandii</i> (Duda, 1940)	B S

EMPIDIDAE

Milan Chvála

OREOGETONINAE**Anthepiscopus** Becker, 1891

<i>caelebs</i> Becker, 1891	M S
<i>ribesii</i> Becker, 1891	S
<i>zontacki</i> (Nowicki, 1871)	S

Gloma Meigen, 1822

<i>fuscipennis</i> Meigen, 1822	B M S
---------------------------------	-------

Iteaphila Zetterstedt, 1838

<i>macquarti</i> Zetterstedt, 1838	S
<i>nitidula</i> Zetterstedt, 1838	B S

Oreogeton Schiner, 1860

<i>basalis</i> (Loew, 1856)	B S
-----------------------------	-----

EMPIDINAE**HILARINI****Hilara** Meigen, 1822

<i>aartseni</i> Chvála, 1997	B
<i>abdominalis</i> Zetterstedt, 1838	B
<i>aeronetha</i> Mik, 1892	B M S
<i>albipennis</i> von Roser, 1840	B M S
<i>albitarsis</i> von Roser, 1840	B M
<i>albiventris</i> von Roser, 1840	B M S
<i>apta</i> Collin, 1927	B S
<i>beckeri</i> Strobl, 1892	B M S
<i>biseta</i> Collin, 1927	B S
<i>bohémica</i> Straka, 1976	B M
= <i>baehrmanii</i> Straka, 1985	
<i>borealis</i> Oldenberg, 1916	B S
<i>brevistyla</i> Collin, 1927	B M S
<i>brevivittata</i> Macquart, 1827	B M S
<i>caerulescens</i> Oldenberg, 1916	S
= <i>chvalai</i> Straka, 1976	
<i>calinota</i> Collin, 1969	S
<i>canescens</i> Zetterstedt, 1849	B M S
<i>chorica</i> (Fallén, 1816)	B M S
<i>cilipes</i> Meigen, 1822	B M S
<i>clavipes</i> (Harris, 1776)	B M S
<i>clypeata</i> Meigen, 1822	B M S
<i>cornicula</i> Loew, 1873	B M S
<i>cuneata</i> Loew, 1873	B
= <i>strobliana</i> Bezzi, 1899	
<i>discalis</i> Chvála, 1997	B M S
<i>discoidalis</i> Lundbeck, 1910	B M S
<i>discolor</i> Strobl, 1892	B M
<i>diversipes</i> Strobl, 1892	B S
<i>eviana</i> Straka, 1976	B S
<i>femorella</i> Zetterstedt, 1842	B S
<i>flavidipes</i> Chvála, 1997	M
<i>flavipes</i> Meigen, 1822	B M S
= <i>cingulata</i> Dahlbom, 1850	
<i>flavocoxa</i> Straka, 1976	S
<i>fuscipes</i> (Fabricius, 1794)	B M S
<i>galactoptera</i> Strobl, 1910	S
<i>gallica</i> (Meigen, 1804)	B M S
<i>griseifrons</i> Collin, 1927	B S
<i>griseola</i> Zetterstedt, 1838	B S
= <i>terriphylla</i> Straka, 1976	
<i>heterogastra</i> Nowicki, 1868	M S
<i>hirta</i> Strobl, 1892	B
<i>hybrida</i> Collin, 1961	B S
<i>hyposeta</i> Straka, 1976	B S
<i>hystrix</i> Strobl, 1892	B
<i>intermedia</i> (Fallén, 1816)	B M S
<i>interstincta</i> (Fallén, 1816)	B M S
<i>lasiochira</i> Strobl, 1892	B
<i>lasiopa</i> Strobl, 1892	B M S
= <i>setimana</i> Straka, 1976	
= <i>matronella</i> Straka, 1976	
<i>litorea</i> (Fallén, 1816)	B M S
<i>longesetosa</i> Strobl, 1910	M

<i>longicornis</i> Strobl, 1894		S	<i>vitripennis</i> Meigen, 1822	B	M	S	
<i>longivittata</i> Zetterstedt, 1842	B	M	S	<i>volucris</i> Wiedemann, 1822	B	S	
<i>lurida</i> (Fallén, 1816)	B	M	S	sg. <i>Empis</i> s.str.			
<i>macquarti</i> Straka, 1984		S	<i>acinerea</i> Chvála, 1985	B	S		
<i>manicata</i> Meigen, 1822	B	M	S	<i>aestiva</i> Loew, 1867	B	M	S
<i>maura</i> (Fabricius, 1776)	B	M	S	<i>alampra</i> Loew, 1873		S	
<i>medeteriformis</i> Collin, 1961	B	M	S	<i>albopilosa</i> de Meijere, 1935	B	M	
<i>media</i> Collin, 1927	B		S	<i>alpicola</i> Strobl, 1893		S	
<i>merula</i> Collin, 1927	B		S	<i>anfractuosa</i> Mik, 1884	B	M	S
<i>monedula</i> Collin, 1927	B	M	S	<i>beckeriana</i> Engel, 1946	B	M	S
<i>morata</i> Collin, 1927	B		S	<i>bicuspidata</i> Collin, 1927	B	M	S
<i>nigrina</i> (Fallén, 1816)	B	M	S	<i>bohémica</i> Chvála & Syrovátka, 1989	B	M	S
<i>nigritarsis</i> Zetterstedt, 1838	B			<i>caudatula</i> Loew, 1867	B	M	S
– <i>infans</i> Zetterstedt, 1842				<i>chioptera</i> Meigen, 1804	B	M	S
<i>nigrocincta</i> de Meijere, 1935	B		S	<i>ciliatopennata</i> Strobl, 1893	B		
<i>nigrohirta</i> Collin, 1927	B		S	<i>dasyprocta</i> Loew, 1867	B	M	S
<i>nitidorella</i> Chvála, 1997	B	M	S	<i>fasciculata</i> Strobl, 1901		S	
<i>nitidula</i> Zetterstedt, 1838	B	M	S	<i>filata</i> Loew, 1873	B	S	
<i>obscura</i> Meigen, 1822	B			<i>florisomna</i> Loew, 1856	B	M	S
<i>pectinipes</i> Strobl, 1892	B	M		<i>gemialis</i> Strobl, 1893	B	M	S
<i>perversa</i> Oldenberg, 1916		S		<i>gooti</i> Chvála, 1994	B	S	
<i>pilosa</i> Zetterstedt, 1838	B	M	S	<i>gymnopoda</i> Bezzi, 1908		M	S
<i>platyura</i> Loew, 1873	B	M	S	<i>laminata</i> Collin, 1927	B	S	
<i>primula</i> Collin, 1927	B		S	<i>levis</i> Loew, 1873	B	S	
<i>pruinosa</i> Wiedemann, 1822	B	M	S	<i>mariae</i> Syrovátka, 1991	B		
<i>pseudochorica</i> Strobl, 1892	B		S	<i>melanotricha</i> Loew, 1873	B	M	S
– <i>woodi</i> Collin, 1927				<i>modesta</i> Meigen, 1838		S	
<i>pseudosartrix</i> Strobl, 1892	B	M	S	<i>nigricoma</i> Loew, 1867	B		
– <i>subcalinota</i> Straka, 1876				<i>nigripes</i> Fabricius, 1794	B	M	S
<i>quadrifaria</i> Strobl, 1892		S		<i>nigritibialis</i> Strobl, 1898	B	M	S
– <i>triglavensis</i> Straka, 1976				<i>nitidissima</i> Strobl, 1893	B	M	S
<i>quadriseta</i> Collin, 1927		S		<i>nitidiventris</i> Loew, 1873		S	
<i>quadrivittata</i> Meigen, 1822	B	M	S	<i>nuntia</i> Meigen, 1838	B	M	S
<i>recedens</i> Walker, 1851		S		<i>pennipes</i> Linnaeus, 1758	B	M	S
<i>rejecta</i> Collin, 1927	B			<i>pilosa</i> Loew, 1867	B	M	S
<i>scrobiculata</i> Loew, 1873		M	S	<i>procera</i> Loew, 1873		S	
<i>sturmi</i> Meigen, 1822	B	M		<i>prodromus</i> Loew, 1867	B	M	S
– <i>emgulata</i> auct. nec Dahlbom, 1850				<i>pseudomalleola</i> Strobl, 1893	B		
<i>subpollinosa</i> Collin, 1927	B	M	S	<i>pseudoprodromus</i> Collin, 1969	B	M	S
<i>tenella</i> (Fallén, 1816)		S		<i>pusio</i> Egger, 1860		M	S
<i>thoracica</i> Macquart, 1827	B	M	S	<i>rufiventris</i> Meigen, 1838	B	M	S
<i>vinnensis</i> Straka, 1976		S		<i>serotina</i> Loew, 1867	B	M	
				<i>simulium</i> Nowicki, 1868	B	S	
EMPIDINI				<i>staegeri</i> Collin, 1963	B	S	
<i>Empis</i> Linnaeus, 1758				<i>syrovatkai</i> Chvála, 1985	B	M	S
sg. <i>Anacrostichus</i> Bezzi, 1909				<i>tanyphyra</i> Loew, 1873	B	M	S
– <i>histortae</i> Meigen, 1822	B	M	S	<i>woodi</i> Collin, 1927		M	S
– <i>nitida</i> Meigen, 1804	B	M		sg. <i>Euempis</i> Frey, 1953			
sg. <i>Argyrandrus</i> Bezzi, 1909				– <i>ciliata</i> Fabricius, 1787	B		
– <i>dispar</i> Scholtz, 1851		S		– <i>kerteszi</i> Bezzi, 1900		S	
sg. <i>Coptophlebia</i> Bezzi, 1909				– <i>picipes</i> Meigen, 1804	B	S	
– <i>albinervis</i> Meigen, 1822	B	M	S	– <i>sericans</i> Brullé, 1832		S	
– <i>dasychira</i> Mik, 1878		M		– <i>tessellata</i> Fabricius, 1794	B	M	S
– <i>hyalipennis</i> Fallén, 1816	B	M	S	sg. <i>Kritempis</i> Collin, 1926			
– <i>impennis</i> Strobl, 1902	B	M	S	– <i>livida</i> Linnaeus, 1758	B	M	S
– <i>leptomorion</i> Bezzi, 1909	B	M	S	sg. <i>Leptempis</i> Collin, 1926			
– <i>pilimana</i> Loew, 1869	B		S	– <i>cognata</i> Egger, 1860	B	M	S

<i>confusa</i> Loew, 1865	B M S	<i>culicina</i> (Fallén, 1816)	B
<i>dimidiata</i> Meigen, 1835	M S	<i>flava</i> (Fallén, 1816)	B S
<i>flavitaris</i> von Roser, 1840	S	<i>flaviventris</i> Macquart, 1827	M S
<i>grisea</i> Fallén, 1816	B M S	<i>heterochroma</i> Bezzi, 1898	B M S
<i>maculata</i> Fabricius, 1781	M S	<i>lamellata</i> Collin, 1926	B M S
<i>mesogramma</i> Loew, 1867	B M S	<i>nigripennis</i> (Fabricius, 1794)	B M S
<i>nigricans</i> Meigen, 1804	B M S	<i>sciarina</i> (Fallén, 1816)	S
<i>rustica</i> Fallén, 1804	B M S	<i>trigemina</i> Oldenberg, 1927	B S
<i>variegata</i> Meigen, 1804	S	<i>umbripennis</i> Meigen, 1822	B M S
sg. <i>Lissemopsis</i> Bezzi, 1909		<i>variabilis</i> (Fallén, 1816)	B M
<i>nigritarsis</i> Meigen, 1804	B	sg. <i>Lundstroemiella</i> Frey, 1922	
sg. <i>Pachymeria</i> Stephens, 1829		<i>dudai</i> Oldenberg, 1927	S
<i>femorata</i> Fabricius, 1798	B M S	<i>hybotina</i> Zetterstedt, 1838	B M S
<i>scotica</i> Curtis, 1835	B M S	<i>kerteszi</i> Oldenberg, 1927	B S
sg. <i>Platyptera</i> Meigen, 1803		<i>magellensis</i> Frey, 1922	M S
<i>borealis</i> Linnaeus, 1758	B M S	sg. <i>Megacyttarus</i> Bigot, 1880	
sg. <i>Polyblepharis</i> Bezzi, 1909		<i>anomaliptennis</i> Meigen, 1822	B M S
<i>albicans</i> Meigen, 1822	S	<i>crassirostris</i> (Fallén, 1816)	B M S
<i>crassa</i> Nowicki, 1868	S	<i>nodipes</i> (Fallén, 1816)	B
<i>eversmanni</i> Loew, 1873	M	<i>poissoni</i> (Trehen, 1966)	B S
<i>fallax</i> Egger, 1860	B	sg. <i>Pararhamphomyia</i> Frey, 1922	
<i>gravipes</i> Loew, 1856	S	<i>aethiops</i> Zetterstedt, 1838	B S
<i>opaca</i> Meigen, 1804	B M S	<i>albidiventris</i> Strobl, 1898	S
<i>strigata</i> Loew, 1867	B	<i>albipennis</i> (Fallén, 1816)	S
sg. <i>Xanthempis</i> Bezzi, 1909		<i>albitarsis</i> Collin, 1926	B
<i>aemula</i> Loew, 1873	B M S	<i>amoena</i> Loew, 1840	M
<i>aequalis</i> Loew, 1867	B M	<i>anfractuosa</i> Bezzi, 1904	B M S
<i>albifrons</i> Bezzi, 1909	S	<i>atra</i> Meigen, 1822	B M S
<i>concolor</i> Verrall, 1872	S	<i>barbata</i> (Macquart, 1823)	B M
<i>digramma</i> Meigen, 1835	B M S	<i>caesia</i> Meigen, 1822	B M
<i>laeta</i> Loew, 1869	B	<i>curvula</i> Frey, 1913	B
<i>laetabilis</i> Collin, 1926	S	<i>genuculata</i> Meigen, 1830	B
<i>lutea</i> Meigen, 1804	B M S	<i>marginata</i> (Fabricius, 1787)	B M S
<i>punctata</i> Meigen, 1804	B M S	<i>micropyga</i> Collin, 1926	B
<i>rohaceki</i> Chvála, 1994	S	<i>modesta</i> Wahlberg, 1844	S
<i>scutellata</i> Curtis, 1835	B M S	<i>murina</i> Collin, 1926	B
<i>semicinerea</i> Loew, 1867	B S	<i>obscura</i> Zetterstedt, 1838	B
<i>stercorea</i> Linnaeus, 1761	B M S	<i>obscuripennis</i> Meigen, 1830	B M
<i>trigramma</i> Wiedemann, 1822	B M S	<i>physoprocta</i> Frey, 1913	B M
<i>univittata</i> Loew, 1867	B M S	<i>pilifer</i> Meigen, 1838	B M S
<i>Rhamphomyia</i> Meigen, 1822		<i>tarsata</i> Meigen, 1822	B M
sg. <i>Actonempis</i> Collin, 1926		<i>tibiella</i> Zetterstedt, 1842	B M S
<i>albohirta</i> Collin, 1926	B M S	<i>tipularia</i> (Fallén, 1816)	S
<i>euplerota</i> Loew, 1873	B M S	<i>unguiculata</i> Frey, 1913	B
<i>galactoptera</i> Strobl, 1893	B M S	sg. <i>Rhamphomyia</i> s.str.	
<i>longipes</i> (Meigen, 1804)	B M S	<i>albosegmentata</i> Zetterstedt, 1838	B S
<i>umbripes</i> Becker, 1887	B	<i>anthracina</i> Meigen, 1822	B M S
sg. <i>Amydroneura</i> Collin, 1926		<i>argentata</i> von Röder, 1887	B M S
<i>claripennis</i> Oldenberg, 1922	B M S	<i>cinerascens</i> (Meigen, 1804)	B M S
<i>crassicauda</i> Strobl, 1893	B M S	<i>crinita</i> Becker, 1887	S
<i>erythrophthalma</i> Meigen, 1830	B M S	<i>czizeki</i> Barták, 1982	M S
<i>gibba</i> (Fallén, 1816)	B M S	<i>janovensis</i> Barták, 1981	B
<i>hirsutipes</i> Collin, 1926	B	<i>laevipes</i> (Fallén, 1816)	B M S
<i>pseudogibba</i> Strobl, 1910	B	<i>lautereri</i> Barták, 1981	S
sg. <i>Holoclera</i> Schiner, 1860		<i>loewi</i> Nowicki, 1868	B S
<i>histriata</i> Strobl, 1910	B M S	<i>luridipennis</i> Nowicki, 1868	B S
<i>caliginosa</i> Collin, 1926	B M	<i>montana</i> Oldenberg, 1915	B M S

<i>morio</i> Zetterstedt, 1838	B		<i>bistigma</i> (Curtis, 1834)	B	S
<i>nitidula</i> Zetterstedt, 1842	B	M	<i>braueri</i> (Mik, 1880)		M
<i>plumipes</i> (Meigen, 1804)	B	M	<i>maxima</i> Wagner, 1984		S
<i>pokornyi</i> Bezzi, 1904	B		<i>rhynchops</i> (Nowicki, 1868)	B	M
<i>siebecki</i> Strobl, 1898		S	<i>stylifera</i> Mik, 1889		M
<i>spinipes</i> (Fallén, 1816)	B	M	<i>tricuspidata</i> (Bezzi, 1905)		S
<i>snigmosa</i> Macquart, 1827	B	M	<i>Clinocera</i> Meigen, 1803		
<i>sulcata</i> (Meigen, 1804)	B	M	sg. <i>Bergenstammia</i> Mik, 1881		
<i>sulcatella</i> Collin, 1926	B	M	<i>slovaca</i> (Wagner, 1984)		S
<i>sulcatina</i> Collin, 1926	B	M	sg. <i>Clinocera</i> s. str.		
<i>tibialis</i> Meigen, 1822	B	M	<i>appendiculata</i> (Zetterstedt, 1838)	B	M
<i>tristriolata</i> Nowicki, 1868		S	<i>fontinalis</i> (Haliday, 1833)	B	M
			<i>nigra</i> Meigen, 1804	B	
			<i>stagnalis</i> (Haliday, 1833)	B	S
			<i>wesmaeli</i> (Macquart, 1835)	B	M
HEMERODROMIINAE			sg. <i>Kowarzia</i> Mik, 1881		
<i>Chelifera</i> Macquart, 1823			<i>barbatula</i> (Mik, 1880)		M
<i>astigma</i> Collin, 1927		S	<i>bipunctata</i> (Haliday, 1833)	B	
<i>concinnicauda</i> Collin, 1927		M	<i>plectrum</i> (Mik, 1880)		M
<i>flavella</i> (Zetterstedt, 1838)	B	M	sg. <i>Phaeobalia</i> Mik, 1881		S
<i>monostigma</i> (Meigen, 1822)	B		<i>dimidiata</i> (Loew, 1869)	B	M
<i>pectinicauda</i> Collin, 1927		S	<i>elapha</i> (Vaillant, 1868)		S
<i>precatoria</i> (Fallén, 1815)	B	M	<i>inermis</i> (Loew, 1861)		S
<i>stigmatica</i> Schiner, 1862	B	M	<i>varipennis</i> (Nowicki, 1868)		S
<i>subangusta</i> Collin, 1961		M			
<i>trapezina</i> (Zetterstedt, 1838)		M			
<i>Chelipoda</i> Macquart, 1823					
<i>albiseta</i> (Zetterstedt, 1838)	B		TRICHOPEZINAE		
<i>vocatoria</i> (Fallén, 1815)	B	M	<i>Heleodromia</i> Haliday, 1833		
<i>Hemerodromia</i> Meigen, 1822			<i>immaculata</i> Haliday, 1833	B	S
<i>melangyna</i> Collin, 1927	B		<i>Trichopeza</i> Rondani, 1856		
<i>oratoria</i> (Fallén, 1815)		S	<i>longicornis</i> (Meigen, 1822)	B	M
<i>raptoria</i> Meigen, 1830	B				
<i>unilineata</i> Zetterstedt, 1842	B	M	GENUS incertae sedis		
<i>Phyllodromia</i> Zetterstedt, 1837			<i>Dryodromia</i> Rondani, 1856		
<i>melanocephala</i> (Fabricius, 1794)	B	M	<i>testacea</i> Rondani, 1856	B	M
		S			
CLINOCERINAE			BRACHYSTOMATINAE		
<i>Dolichocephala</i> Macquart, 1823			<i>Brachystoma</i> Meigen, 1822		
<i>cavatica</i> (Becker, 1889)		S	<i>vesiculosum</i> (Fabricius, 1794)		S
<i>engeli</i> Niesiolowski, 1992	B	S			
<i>gutata</i> (Haliday, 1833)		M	HYBOTIDAE		
<i>irrorata</i> (Fallén, 1815)	B	M	Milan Chvála		
<i>Wiedemannia</i> Zetterstedt, 1838					
sg. <i>Chamaedipsia</i> Mik, 1881			OCYDROMIINAE		
<i>beckeri</i> (Mik, 1889)		S	TRICHININI		
sg. <i>Eucelidia</i> Mik, 1882			<i>Bicellaria</i> Macquart, 1823		
<i>escheri</i> (Zetterstedt, 1838)		M	<i>albopilosa</i> Chvála, 1991	B	M
<i>pirata</i> (Mik, 1880)	B	S	<i>austriaca</i> Tuomikoski, 1955	B	M
<i>zetterstedti</i> (Fallén, 1826)	B	M	<i>intermedia</i> Lundbeck, 1910	B	M
sg. <i>Philolutra</i> Mik, 1881			<i>mera</i> Collin, 1961		S
<i>aquilex</i> (Loew, 1869)		M	<i>nigra</i> (Meigen, 1824)	B	M
<i>bohemani</i> (Zetterstedt, 1838)	B	M	<i>nigrita</i> Collin, 1926	B	S
<i>fallaciosa</i> (Loew, 1873)		S	<i>pilosa</i> Lundbeck, 1910	B	
<i>hygrobia</i> (Loew, 1858)	B	S	<i>simplicipes</i> (Zetterstedt, 1842)	B	S
<i>phantasma</i> (Mik, 1880)		S	<i>spuria</i> (Fallén, 1816)	B	M
sg. <i>Pseudowiedemannia</i> Engel, 1918			<i>subpilosa</i> Collin, 1926	B	M
<i>lamellata</i> (Loew, 1869)		M	<i>sulcata</i> (Zetterstedt, 1842)	B	S
sg. <i>Wiedemannia</i> s.str.		S	<i>vana</i> Collin, 1926	B	M

<i>Trichina</i> Meigen, 1830			<i>albifacies</i> (Collin, 1926)	B	M	S
<i>bilobata</i> Collin, 1926	B	M	<i>albiseta</i> (Panzer, 1806)	B	M	S
<i>clavipes</i> Meigen, 1830	B	M	<i>albistylus</i> Chvála, 1989			S
<i>elongata</i> Haliday, 1833	B	M	<i>alpigenus</i> (Strobl, 1893)			S
<i>opaca</i> Loew, 1864	B	M	<i>alpinus</i> Chvála, 1971	B		S
<i>pallipes</i> (Zetterstedt, 1838)		M	<i>analís</i> (Meigen, 1830)	B	M	
<i>Trichinomyia</i> Tuomikoski, 1959			<i>annulatus</i> (Fallén, 1815)	B	M	S
<i>flavipes</i> (Meigen, 1830)	B	M	<i>annulipes</i> (Meigen, 1822)	B	M	S
OEDALEINI			<i>aristatus</i> (Collin, 1926)	B	M	S
<i>Allanthalia</i> Melander, 1928			<i>articulatooides</i> (Frey, 1918)	B	M	S
<i>pallida</i> (Zetterstedt, 1838)	B		<i>articulatus</i> Macquart, 1827	B	M	S
<i>Anthalia</i> Zetterstedt, 1838			<i>aurantiacus</i> (Collin, 1926)		M	
<i>beatricella</i> Chandler, 1992	B		<i>australominutus</i> Grootaert, 1989	B		
<i>Euthyneura</i> Macquart, 1836			<i>balticus</i> Kovalev, 1971		M	S
<i>gyllenhali</i> (Zetterstedt, 1838)	B	M	<i>bartaki</i> Chvála, 1989	B		S
<i>halidayi</i> Collin, 1926		M	<i>biapicalis</i> Wéber, 1972	B		S
<i>myrtilli</i> Macquart, 1836	B	M	<i>bilobatus</i> Wéber, 1972	B	M	S
<i>Oedalea</i> Meigen, 1820			<i>boreoalpinus</i> Frey, 1943	B	M	
<i>austroholmgreni</i> Chvála, 1981	B		<i>brachystylus</i> (Bezzi, 1892)	B	M	S
<i>flavipes</i> Zetterstedt, 1842	B	M	<i>brevicornis</i> (Zetterstedt, 1842)	B	M	
<i>hybotina</i> (Fallén, 1816)	B	M	<i>calceatus</i> (Meigen, 1822)	B	M	S
<i>kowarzi</i> Chvála, 1981	B		<i>candicans</i> (Fallén, 1815)	B	M	S
<i>montana</i> Chvála, 1981	B		<i>celer</i> (Meigen, 1822)	B		
<i>stigmatella</i> Zetterstedt, 1842	B	M	<i>ciliaris</i> (Fallén, 1816)	B	M	S
<i>tibialis</i> Macquart, 1827	B	M	<i>cilitarsis</i> Frey, 1943		M	S
<i>tristis</i> Scholtz, 1851	B	M	<i>clarandus</i> (Collin, 1926)	B	M	S
<i>zetterstedti</i> Collin, 1926	B	M	<i>clypeatus</i> Kovalev, 1973	B		S
OCYDROMIINI			<i>collini</i> (Chvála, 1966)	B	M	S
<i>Leptopeza</i> Macquart, 1827			<i>commutatus</i> (Strobl, 1893)	B		S
<i>borealis</i> Zetterstedt, 1842	B	M	<i>confiformis</i> Chvála, 1971			S
<i>flavipes</i> (Meigen, 1820)	B	M	<i>cothurnatus</i> Macquart, 1827	B	M	S
<i>Ocydromia</i> Meigen, 1820			<i>cruralis</i> (Collin, 1961)	B	M	S
<i>glabricula</i> (Fallén, 1816)	B	M	<i>cryptospina</i> (Frey, 1909)	B	M	S
<i>melanopleura</i> Loew, 1840	B	M	<i>cursitans</i> (Fabricius, 1775)	B	M	S
<i>Oropezella</i> Collin, 1926			<i>dilatatorvittatus</i> (Strobl, 1910)			S
<i>sphenoptera</i> (Loew, 1873)	B	M	<i>divisus</i> Walker, 1851		M	S
HYBOTINAE			<i>ecalceatus</i> (Zetterstedt, 1838)	B	M	S
<i>Hybos</i> Meigen, 1803			<i>excisus</i> (Becker, 1907)	B	M	S
<i>culiciformis</i> (Fabricius, 1775)	B	M	<i>exilis</i> (Meigen, 1822)	B	M	S
<i>femoratus</i> (Müller, 1776)	B	M	<i>flavicornis</i> (Meigen, 1822)	B	M	
<i>grossipes</i> (Linnaeus, 1767)	B	M	<i>fuscicornis</i> (Zetterstedt, 1842)	B	M	S
<i>Syndyas</i> Loew, 1857			<i>infectus</i> (Collin, 1926)	B	M	S
<i>nigripes</i> (Zetterstedt, 1842)	B		<i>ingenuus</i> (Collin, 1926)		M	
<i>Syneches</i> Walker, 1852			<i>insperatus</i> Kovalev, 1971		M	S
<i>muscarinus</i> (Fabricius, 1794)		M	<i>interstinctus</i> (Collin, 1926)	B	M	
TACHYDROMIINAE			<i>kirtlingensis</i> Grootaert, 1986	B	M	S
TACHYDROMIINI			<i>laticinctus</i> Walker, 1851	B	M	S
<i>Dysaletria</i> Loew, 1864			<i>leucocephalus</i> (von Roser, 1840)	B		
<i>nigripennis</i> Chvála, 1975		S	<i>leucothrix</i> (Strobl, 1910)	B		
<i>Platypalpus</i> Macquart, 1827			<i>longicornis</i> (Meigen, 1822)	B	M	S
<i>aeneus</i> (Macquart, 1823)	B	M	<i>longimanus</i> (Corti, 1907)	B		
<i>agilis</i> (Meigen, 1822)	B	M	<i>longiseta</i> (Zetterstedt, 1842)	B	M	S
<i>albicornis</i> (Zetterstedt, 1842)	B	M	<i>luteicornis</i> (Meigen, 1838)	B	M	S
<i>albidifacies</i> Chvála, 1975		M	<i>luteipes</i> Zusková, 1966	B	M	S
			<i>luteoloides</i> Grootaert, 1983		M	
			<i>luteolus</i> (Collin, 1926)	B	M	S
			<i>luteus</i> (Meigen, 1804)	B	M	S

<i>macula</i> (Zetterstedt, 1842)	B M S	<i>Symballophthalmus</i> Becker, 1889	
<i>maculipes</i> (Meigen, 1822)	B M S	<i>dissimilis</i> (Fallén, 1815)	B
<i>major</i> (Zetterstedt, 1842)	B M S	<i>fuscitarsis</i> (Zetterstedt, 1859)	B M S
<i>melancholicus</i> (Collin, 1961)	B M S	<i>pictipes</i> (Becker, 1889)	B M S
<i>miki</i> (Becker, 1890)	B M S	<i>Tachydromia</i> Meigen, 1822	
<i>minutus</i> (Meigen, 1804)	B M S	<i>aemula</i> (Loew, 1864)	B M S
<i>namus</i> (Oldenberg, 1924)	M	<i>annulimana</i> Meigen, 1822	B M S
<i>nigellus</i> (Collin, 1969)	S	<i>arrogans</i> (Linnaeus, 1761)	B M S
<i>niger</i> (Meigen, 1804)	B M S	<i>calcaranea</i> (Meigen, 1838)	B M S
<i>nigricea</i> (Mik, 1884)	B S	<i>carpathica</i> Chvála, 1966	S
<i>nigrimanus</i> Strobl, 1880	B S	<i>connexa</i> Meigen, 1822	B M S
<i>nigritarsis</i> (Fallén, 1816)	B M S	<i>costalis</i> (von Roser, 1840)	B S
<i>nigrosetosus</i> (Strobl, 1893)	B	<i>interrupta</i> (Loew, 1864)	M S
<i>niveisetia</i> (Zetterstedt, 1842)	B M S	<i>microptera</i> (Loew, 1864)	S
<i>notatus</i> (Meigen, 1822)	B M S	<i>parva</i> Chvála, 1970	S
<i>nudithorax</i> Chvála, 1975	B S	<i>sabulosa</i> Meigen, 1830	S
<i>ochrocera</i> (Collin, 1961)	B M S	<i>smithi</i> Chvála, 1966	B S
<i>optimus</i> (Collin, 1926)	B M	<i>terricola</i> Zetterstedt, 1819	S
<i>palavensis</i> Chvála, 1989	B M S	<i>tuberculata</i> (Loew, 1864)	M
<i>pallidicornis</i> (Collin, 1926)	B M S	<i>umbrarum</i> Haliday, 1833	B S
<i>pallidicea</i> (Frey, 1913)	B S	<i>undulata</i> (Strobl, 1906)	S
<i>pallidiseta</i> Kovalev, 1978	S	<i>woodi</i> (Collin, 1926)	B S
<i>pallidiventris</i> (Meigen, 1822)	B M S	<i>Tachypeza</i> Meigen, 1830	
<i>pallipes</i> (Fallén, 1815)	B M S	<i>fennica</i> Tuomikoski, 1932	B M S
<i>parvicauda</i> (Collin, 1926)	B M S	<i>fuscipennis</i> (Fallén, 1815)	B M S
<i>pectoralis</i> (Fallén, 1815)	B M S	<i>heeri</i> Zetterstedt, 1838	B S
<i>pictitarsis</i> (Becker, 1902)	B M S	<i>nubila</i> (Meigen, 1804)	B M S
<i>pragensis</i> Chvála, 1989	B	<i>tanaisense</i> Kovalev, 1975	M
<i>pseudociliaris</i> (Strobl, 1910)	B	<i>truncorum</i> (Fallén, 1815)	B S
<i>pseudofulvipes</i> (Frey, 1909)	B M S		
<i>pseudorapidus</i> Kovalev, 1971	B M S	DRAPETINI	
<i>pulicarius</i> (Meigen, 1830)	B S	<i>Chersodromia</i> Walker, 1849	
<i>pygialis</i> Chvála, 1973	B M S	<i>cursitans</i> (Zetterstedt, 1819)	M
<i>rapidoides</i> Chvála, 1975	B M S	<i>Crossopalpus</i> Bigot, 1857	
<i>rapidus</i> (Meigen, 1822)	B M	<i>abditus</i> Kovalev, 1972	B S
<i>ringdahli</i> Chvála, 1975	B	<i>curvinervis</i> (Zetterstedt, 1842)	B
<i>ruficornis</i> (von Roser, 1840)	B S	<i>humilis</i> (Frey, 1913)	B M S
<i>scandinavicus</i> Chvála, 1972	S	<i>nigritellus</i> (Zetterstedt, 1842)	B M S
<i>sloveniensis</i> Bequaert, 1962	B M	<i>setiger</i> (Loew, 1859)	B M
<i>smirnovi</i> Kovalev, 1978	B	<i>Drapetis</i> Meigen, 1822	
<i>soosi</i> Chvála, 1989	B	sg. <i>Drapetis</i> s.str	
<i>stabilis</i> (Collin, 1961)	B M S	<i>arcuata</i> Loew, 1859	B M
<i>stackelbergi</i> Kovalev, 1971	B S	<i>assimilis</i> (Fallén, 1815)	B M S
<i>stigma</i> (Collin, 1926)	S	<i>convergens</i> Collin, 1926	B
<i>stigmatelloides</i> Grootaert & Chvála, 1988	S	<i>exilis</i> Meigen, 1822	B S
<i>stigmatellus</i> (Zetterstedt, 1842)	B M S	<i>flavipes</i> Macquart, 1834	B
<i>strakai</i> Chvála, 1989	S	<i>incompleta</i> Collin, 1926	B M S
<i>stroblii</i> (Mik, 1900)	B M S	<i>infitalis</i> Collin, 1961	B
<i>styloformis</i> Weber, 1972	M S	<i>parilis</i> Collin, 1926	B S
<i>subbrevis</i> (Frey, 1913)	B	<i>pusilla</i> Loew, 1859	B
<i>subtilis</i> (Collin, 1926)	B M	<i>simulans</i> Collin, 1961	B M
<i>sylvicola</i> (Collin, 1926)	M	sg. <i>Elaphropeza</i> Macquart, 1827	
<i>tonsus</i> (Collin, 1961)	B M S	<i>ephippiata</i> (Fallén, 1815)	B M S
<i>tuomikoskii</i> Chvála, 1972	B S	<i>Stilpon</i> Loew, 1859	
<i>vegetus</i> Frey, 1943	B	<i>graminum</i> (Fallén, 1815)	B M S
<i>vegrandis</i> Frey, 1943	B M S	<i>lunatus</i> (Walker, 1851)	M S
<i>verralli</i> (Collin, 1926)	B	<i>nubilus</i> Collin, 1926	B

DOLICHOPODIDAE

Jiří Olejníček

SCIAPODINAE

Sciapus Zeller, 1842

<i>albifrons</i> (Meigen, 1830)	B	M	S
<i>bellus</i> Loew, 1873		M	S
<i>contristans</i> (Wiedemann, 1817)	B	M	S
<i>flavicinctus</i> (Loew, 1857)			S
<i>frater</i> Parent, 1927			S
<i>gracilipes</i> (Loew, 1871)			S
<i>heteropygus</i> Parent, 1926		M	S
<i>lobipes</i> (Meigen, 1824)	B		S
<i>longulus</i> (Fallén, 1823)	B	M	S
<i>maritimus</i> Becker, 1918	B	M	
<i>nervosus</i> (Lehmann, 1822)	B		
<i>platypterus</i> (Fabricius, 1805)	B	M	S
<i>wiedemanni</i> (Fallén, 1823)	B		S

XANTHOCHLORINAE

Xanthochlorus Loew, 1857

<i>ornatus</i> (Haliday, 1832)	B	M	S
<i>tenellus</i> (Wiedemann, 1817)	B	M	S

RHAPHIINAE

Nematoproctus Loew, 1857

<i>distendens</i> (Meigen, 1824)	B	M	S
<i>longifilius</i> Loew, 1857	B	M	
<i>praeseclus</i> Loew, 1869	B	M	S

Rhaphium Meigen, 1803

<i>albifrons</i> Zetterstedt, 1843	B	M	
<i>albomaculatum</i> (Becker, 1901)	B		
<i>antennatum</i> (Carlier, 1835)	B	M	S
<i>auctum</i> Loew, 1857	B	M	S
<i>caliginosum</i> Meigen, 1824	B	M	S
<i>commune</i> (Meigen, 1824)	B	M	S
<i>crassipes</i> (Meigen, 1824)	B	M	S
<i>discigerum</i> Stenhammar, 1851	B		
<i>elegantulum</i> (Meigen, 1824)	B	M	
<i>ensicorne</i> Meigen, 1824	B	M	
<i>fasciatum</i> Meigen, 1824	B	M	S
<i>fascipes</i> (Meigen, 1824)	B	M	S
<i>fissum</i> Loew, 1850	B	M	S
<i>intermedium</i> (Becker, 1918)	B		
<i>laticorne</i> (Fallén, 1823)	B	M	S
<i>longicorne</i> (Fallén, 1823)	B	M	S
<i>macrocerum</i> Meigen, 1824	B	M	S
<i>micans</i> (Meigen, 1824)	B	M	S
<i>monotrichum</i> Loew, 1850	B	M	S
<i>nasutum</i> (Fallén, 1823)	B	M	S
<i>patulum</i> Raddatz, 1873	B		
<i>penicillatum</i> Loew, 1850	B	M	S
<i>praerosum</i> Loew, 1850	B	M	S
<i>rivale</i> (Loew, 1869)		M	S
<i>suave</i> (Loew, 1859)		M	S
<i>xiphias</i> Meigen, 1824	B		
<i>zetterstedti</i> (Parent, 1925)	B	M	S

PELOROPEODINAE

Acropsilus Mik, 1878

<i>niger</i> (Loew, 1869)	B		S
---------------------------	---	--	---

Anepsiomyia Bezzi, 1902

<i>flaviventris</i> (Meigen, 1824)	B		S
------------------------------------	---	--	---

Chrysotimus Loew, 1857

<i>flaviventris</i> (von Roser, 1840)	B	M	S
---------------------------------------	---	---	---

<i>mollicullus</i> (Fallén, 1823)		M	S
-----------------------------------	--	---	---

Micromorphus Mik, 1878

<i>albipes</i> (Zetterstedt, 1843)	B	M	S
------------------------------------	---	---	---

ENLININAE

Achalcus Loew, 1857

<i>cinereus</i> (Haliday, 1851)	B	M	S
---------------------------------	---	---	---

<i>flavicollis</i> (Meigen, 1824)	B	M	S
-----------------------------------	---	---	---

HYDROPHORINAE

THINOPHILINI

Peodes Loew, 1857

<i>forcipatus</i> Loew, 1857		M	S
------------------------------	--	---	---

<i>petsamoensis</i> Frey, 1930		M	
--------------------------------	--	---	--

Schoenophilus Mik, 1878

<i>versutus</i> (Haliday, 1851)		M	
---------------------------------	--	---	--

Thinophilus Wahlberg, 1844

<i>flavipalpis</i> (Zetterstedt, 1843)	B		
--	---	--	--

<i>ruficornis</i> (Haliday, 1838)	B		
-----------------------------------	---	--	--

HYDROPHORINI

Hydrophorus Fallén, 1823

<i>balticus</i> (Meigen, 1824)	B	M	S
--------------------------------	---	---	---

<i>bipunctatus</i> (Lehmann, 1822)	B	M	
------------------------------------	---	---	--

<i>borealis</i> Loew, 1857	B		
----------------------------	---	--	--

<i>litoreus</i> Fallén, 1823	B	M	S
------------------------------	---	---	---

<i>nebulosus</i> Fallén, 1823	B		
-------------------------------	---	--	--

<i>praecox</i> (Lehmann, 1822)	B	M	S
--------------------------------	---	---	---

<i>rufinasutus</i> Parent, 1925		M	
---------------------------------	--	---	--

<i>signiferus</i> Coquillett, 1899	B		
------------------------------------	---	--	--

<i>viridis</i> (Meigen, 1824)	B	M	S
-------------------------------	---	---	---

Liancalus Loew, 1857

<i>virens</i> (Scopoli, 1763)	B	M	S
-------------------------------	---	---	---

Scellus Loew, 1857

<i>notatus</i> (Fabricius, 1871)			S
----------------------------------	--	--	---

NEURIGONINAE

Neurigona Rondani, 1856

<i>abdominalis</i> Fallén, 1823	B		
---------------------------------	---	--	--

<i>cilipes</i> (Oldenberg, 1904)			S
----------------------------------	--	--	---

<i>erichsoni</i> (Zetterstedt, 1843)		M	S
--------------------------------------	--	---	---

<i>pallida</i> (Fallén, 1823)	B	M	S
-------------------------------	---	---	---

<i>quadrifasciata</i> (Fabricius, 1871)	B	M	S
---	---	---	---

<i>suturalis</i> (Fallén, 1823)	B	M	S
---------------------------------	---	---	---

Oncopygius Mik, 1866

<i>distans</i> (Loew, 1857)			S
-----------------------------	--	--	---

SYMPYCNINAE

SYMPYCNINI

Sympycnus Loew, 1857

<i>aeneicoxa</i> (Meigen, 1824)	B	M	S
---------------------------------	---	---	---

<i>annulipes</i> (Meigen, 1824)	B M S	DIAPHORINAE	
<i>brevimanus</i> Loew, 1857	S	ARGYRINI	
<i>cirripes</i> (Haliday in Walker, 1851)	B M S	<i>Argyra</i> Macquart, 1834	
<i>kowarzi</i> Parent, 1925	B	<i>argentina</i> Meigen, (1824)	B M S
<i>simplicipes</i> Becker, 1908	M	<i>argyria</i> (Meigen, 1824)	B M S
<i>spiculatus</i> Gerstaecker, 1864	S	<i>atriceps</i> Loew, 1857	B M S
		<i>auricollis</i> (Meigen, 1824)	B M S
CAMPSICNEMINI		<i>confinis</i> (Zetterstedt, 1849)	B M S
<i>Campsicnemus</i> Haliday, 1851		<i>diaphana</i> (Fabricius, 1775)	B M S
<i>armatus</i> (Zetterstedt, 1849)	B M S	<i>discedens</i> Becker, 1907	B M
<i>armoriacus</i> Parent, 1926	B	<i>elongata</i> (Zetterstedt, 1843)	B M
<i>compeditus</i> Loew, 1857	B M	<i>grata</i> Loew, 1857	B M S
<i>curvipes</i> (Fallén, 1823)	B M S	<i>hoffmeisteri</i> (Loew, 1850)	B M S
<i>filipes</i> Loew, 1859	S	<i>leucocephalla</i> (Meigen, 1824)	B M S
<i>loripes</i> (Haliday, 1832)	B M S	<i>loewi</i> Kowarz, 1879	B
<i>lunhatus</i> Loew, 1857	B M S	<i>miki</i> Kowarz, 1857	B
<i>magus</i> (Loew, 1845)	B	<i>spoliata</i> Kowarz, 1879	B
<i>mamillatus</i> Mik, 1869	B	<i>setimana</i> Loew, 1859	M S
<i>marginatus</i> Loew, 1857	B M S	<i>estita</i> (Wiedemann, 1817)	B M S
<i>pectinulatus</i> Loew, 1864	B		
<i>picicornis</i> (Zetterstedt, 1843)	B M S	DIAPHORINI	
<i>pumilio</i> (Zetterstedt, 1843)	B	<i>Asyndetus</i> Loew, 1869	
<i>pusillus</i> (Meigen, 1824)	B S	<i>latifrons</i> (Loew, 1857)	B M
<i>scambus</i> (Fallén, 1823)	B M S	<i>Chrysotus</i> Meigen, 1824	
<i>umbripennis</i> Loew, 1856	B M S	<i>angulicornis</i> Kowarz, 1874	B M S
<i>varipes</i> Loew, 1857	S	<i>cilipes</i> Meigen, 1824	B M S
<i>Lamprochromus</i> Mik, 1878		<i>cupreus</i> (Macquart, 1827)	B M S
<i>elegans</i> (Meigen, 1830)	B M S	<i>femoratus</i> Zetterstedt, 1843	B M S
<i>kowarzi</i> Negrobov & Tshalaja, 1988	S	<i>gramineus</i> (Fallén, 1823)	B M S
<i>moraviensis</i> Negrobov & Tshalaja, 1988	M	<i>laesus</i> (Wiedemann, 1817)	B M S
<i>strobli</i> Parent, 1925	M	<i>neglectus</i> (Wiedemann, 1817)	B M
<i>Syntormon</i> Loew, 1857		<i>obscuripes</i> Zetterstedt, 1838	B M S
– <i>Bathycranium</i> Strobl, 1892		<i>palustris</i> Verrall, 1876	B M
<i>aulicus</i> (Meigen, 1824)	B M	<i>pulchellus</i> Kowarz, 1874	B M S
<i>bicolorellum</i> (Zetterstedt, 1843)	B M S	<i>suavis</i> Loew, 1857	B M S
<i>fuscipes</i> (von Roser, 1840)	B M S	<i>viridifemoratus</i> (von Roser, 1840)	B M S
<i>metathesis</i> (Loew, 1850)	B S	<i>Diaphorus</i> Meigen, 1824	
<i>miki</i> Strobl, 1869	M	<i>disjunctus</i> Loew, 1857	S
<i>monilis</i> (Haliday, 1851)	B M S	<i>halteralis</i> Loew, 1869	S
<i>pallipes</i> (Fabricius, 1794)	B M S	<i>hoffmannseggii</i> Meigen, 1830	B M S
<i>pumilus</i> (Meigen, 1824)	B M S	<i>nigricans</i> Meigen, 1824	B M
<i>punctatus</i> (Zetterstedt, 1843)	B M	<i>oculatus</i> (Fallén, 1923)	B M S
<i>rufipes</i> (Meigen, 1824)	B M S	<i>winthemi</i> Meigen, 1824	B
<i>subinermis</i> Loew, 1869	M S	<i>Melanostolus</i> Kowarz, 1884	
<i>sulcipes</i> (Meigen, 1824)	B M	<i>melancholicus</i> (Loew, 1869)	B
<i>tarsatus</i> (Fallén, 1823)	B M S		
<i>zelleri</i> (Loew, 1850)	B	DOLICHOPODINAE	
<i>Telmaturgus</i> Mik, 1874		TACHYTRECHINI	
<i>tumichulus</i> (Raddatz, 1873)	M	<i>Tachytrechus</i> Haliday, 1851	
<i>Teuchophorus</i> Loew, 1857		<i>consobrinus</i> Haliday, 1851	B
<i>calcaratus</i> (Macquart, 1827)	B M S	<i>genualis</i> Loew, 1857	B S
<i>monacanthus</i> Loew, 1859	B M	<i>insignis</i> (Stannius, 1831)	B M S
<i>simplex</i> Mik, 1880	M	<i>kowarzii</i> Mik, 1865	S
<i>spinigerellus</i> (Zetterstedt, 1833)	B M	<i>notatus</i> (Stannius, 1831)	B S
<i>nigricosta</i> (von Roser, 1840)	B M S	<i>ocior</i> Loew, 1869	B S
		<i>ripicola</i> Loew, 1857	S

DOLICHOPODINI

Dolichopus Latreille, 1796sg. *Dolichopus* s. str.

<i>acuticornis</i> Wiedemann, 1817	B M S	<i>salictorum</i> Loew, 1871	B S
<i>agilis</i> Meigen, 1824	B M S	<i>signatus</i> Meigen, 1824	B M S
<i>albifrons</i> Loew, 1859	M	<i>signifer</i> Haliday, 1838	B M S
<i>annulipes</i> Zetterstedt, 1849	M	<i>simplex</i> Meigen, 1824	B M S
<i>apicalis</i> Zetterstedt, 1849	B	<i>subpennatus</i> D'Assis-Fonseca, 1976	B M S
<i>arbustorum</i> Stannius, 1831	B M S	<i>tanythrix</i> Loew, 1869	B M
<i>argyrotarsis</i> Wahlberg, 1850	B M S	<i>trivialis</i> Haliday, 1831	B M S
<i>armillatus</i> Wahlberg, 1850	B	<i>ungulatus</i> (Linnaeus, 1758)	B M S
<i>atratus</i> Meigen, 1824	B M S	<i>urbanus</i> Meigen, 1824	B S
<i>atripes</i> Meigen, 1824	B M S	<i>vitripennis</i> Meigen, 1824	B S
<i>brevipennis</i> Meigen, 1824	B M	<i>wahlbergi</i> Zetterstedt, 1843	B M S
<i>campestris</i> Meigen, 1824	B M S	sg. <i>Hygroceleuthus</i> Loew, 1857	
<i>cilifemoratus</i> Macquart, 1827	B M S	<i>latipennis</i> Fallén, 1823	B
<i>claviger</i> Stannius, 1831	B M S	<i>Hercostomus</i> Loew, 1857	
<i>clavipes</i> Haliday, 1832	B	sg. <i>Hercostomus</i> s. str.	
<i>consobrinus</i> Zetterstedt, 1859	S	<i>aerosus</i> (Fallén, 1823)	B M S
<i>cruralis</i> Wahlberg, 1850	B M S	<i>angustifrons</i> (Staeger, 1842)	B
<i>discimanus</i> Wahlberg, 1850	B M S	<i>assimilis</i> (Staeger, 1842)	B M
<i>eurypterus</i> Gerstaecker, 1864	B	<i>bicolor</i> (Macquart, 1827)	B M
<i>excisus</i> Loew, 1859	B M S	<i>blankaartensis</i> Pollet, 1990	M
<i>falcatus</i> Becker, 1917	M	<i>brevicornis</i> (Staeger, 1842)	B M
<i>festivus</i> Haliday, 1832	B M S	<i>caudatus</i> (Loew, 1859)	B
<i>flavipes</i> Stannius, 1831	B	<i>celer</i> (Meigen, 1824)	B M S
<i>geniculatus</i> Stannius, 1831	B	<i>chaerophyli</i> (Meigen, 1824)	B M
<i>griseipennis</i> Stannius, 1831	B M S	<i>chalybeus</i> (Wiedemann, 1817)	B M S
<i>hilaris</i> Loew, 1862	B M	<i>chetifer</i> (Walker, 1849)	B M S
<i>immaculatus</i> Becker, 1909	B	<i>cupreus</i> (Fallén, 1823)	B S
<i>kowarzianus</i> Stackelberg, 1928	B	<i>fugax</i> (Loew, 1857)	B M S
<i>lanlimbanus</i> Macquart, 1827	B M S	<i>fulvicaudis</i> (Haliday, 1851)	M S
<i>latipennis</i> Fallén, 1823	B	<i>fuscipennis</i> (Meigen, 1824)	B M S
<i>lepidus</i> Staeger, 1842	B M S	<i>germanus</i> (Wiedemann, 1817)	B M S
<i>linearis</i> Meigen, 1824	B M S	<i>inornatus</i> (Loew, 1857)	S
<i>lineaticornis</i> Zetterstedt, 1843	B M	<i>longiventris</i> (Loew, 1857)	B M
<i>litorellus</i> Zetterstedt, 1852	B S	<i>metallicus</i> (Stannius, 1831)	B M S
<i>longicornis</i> Stannius, 1831	B M S	<i>nanus</i> (Macquart, 1827)	B M S
<i>meigenii</i> Loew, 1857	M	<i>nigrilamellatus</i> (Macquart, 1827)	M S
<i>melanopus</i> Meigen, 1824	B S	<i>nigripennis</i> (Fallén, 1823)	B
<i>migrans</i> Zetterstedt, 1849	B S	<i>nigriplantis</i> (Stannius, 1831)	B M S
<i>nigricornis</i> Meigen, 1824	B M S	<i>plagiatus</i> (Loew, 1857)	B M
<i>nigripes</i> Fallén, 1823	B M	<i>praeceps</i> Loew, 1869	B M
<i>nitidus</i> Fallén, 1823	B M S	<i>rusticus</i> (Meigen, 1824)	B M S
<i>notatus</i> Staeger, 1842	B M S	<i>sahlbergi</i> (Zetterstedt, 1838)	B S
<i>nubilus</i> Meigen, 1824	B M S	<i>silvestris</i> Pollet, 1990	M
<i>occulus</i> Becker, 1917	B	<i>vivax</i> (Loew, 1857)	B M S
<i>parvicaudatus</i> Zetterstedt, 1843	B	<i>vockerothi</i> d'Assis-Fonseca, 1976	S
<i>pennatus</i> Meigen, 1824	B M S	sg. <i>Poecilobothrus</i> Mik, 1878	
<i>phaeopus</i> Haliday, 1851	B	<i>chrysozygos</i> (Wiedemann, 1817)	B M S
<i>picipes</i> Meigen, 1824	B M S	<i>comitalis</i> (Kowarz, 1867)	M S
<i>plantarsis</i> Fallén, 1823	B S	<i>fumipennis</i> (Stannius, 1831)	B M
<i>plumipes</i> (Scopoli, 1763)	B M S	<i>infuscatus</i> (Stannius, 1831)	B M S
<i>plumitarsis</i> Fallén, 1823	B M S	<i>nobilatus</i> (Linnaeus, 1767)	B M S
<i>popularis</i> Wiedemann, 1817	B M S	<i>regalis</i> (Meigen, 1824)	M S
<i>rupestris</i> Haliday, 1838	B M	<i>Hypophyllus</i> Haliday, 1832	
<i>sabinus</i> Haliday, 1838	B M S	<i>crinipes</i> (Staeger, 1842)	B M S
		<i>discipes</i> (Germar, 1817)	B M
		<i>obscurus</i> (Fallén, 1823)	B M S
		<i>sciophilus</i> Loew, 1869	M S

<i>sphenopterus</i> Loew, 1859	B M	<i>bellus</i> Loew, 1869	B M S
<i>Sybistroma</i> Meigen, 1824		<i>cuneatus</i> (Becker, 1917)	M
<i>nodicornis</i> Meigen, 1824	B M S	<i>laetus</i> Verrall, 1912	M
<i>setosa</i> Schiner, 1862	S	<i>smaragdinus</i> Gerstaecker, 1864	M
MEDETERINAE		MICROPHORINAE	
SYSTEMINI		MICROPHORINI	
<i>Systemus</i> Loew, 1857		<i>Microphor</i> Macquart, 1827	
<i>bipartitus</i> (Loew, 1850)	B	<i>anomalus</i> (Meigen, 1824)	B M S
<i>pallipes</i> (von Roser, 1840)	B M	<i>crassipes</i> Macquart, 1827	B M S
MEDETERINI		<i>holosericeus</i> (Meigen, 1804)	B M S
<i>Dolichophorus</i> Lichtward, 1902		<i>intermedius</i> Collin, 1961	B M S
<i>kerteszi</i> Lichtward, 1902	B M S	<i>strobli</i> Chvála, 1986	B S
<i>Medetera</i> Fisher von Waldheim, 1819		PARATHALASSIINI	
<i>abstrusa</i> Thunberg, 1955	B S	<i>Microphorella</i> Becker, 1909	
<i>ambigua</i> (Zetterstedt, 1843)	B S	<i>praecox</i> (Loew, 1864)	S
<i>annulitarsus</i> von Roser, 1840	B S	ATELESTIDAE	
<i>apicalis</i> (Zetterstedt, 1843)	B S	Milan Chvála	
<i>bispinosa</i> Negrobov, 1967	M	<i>Atelestus</i> Walker, 1837	
<i>borealis</i> Thunberg, 1955	B	<i>dissonans</i> Collin, 1961	B
<i>chrysotiformis</i> Kowarz, 1877	B M S	<i>pulicarius</i> (Fallén, 1816)	B M S
<i>dendrobaena</i> Kowarz, 1877	B	<i>Meghyperus</i> Loew, 1850	
<i>diadema</i> (Linnaeus, 1767)	B M S	<i>sudeticus</i> Loew, 1850	S
<i>dichrocera</i> Kowarz, 1878	B	OPETIIDAE	
<i>excellens</i> Frey, 1909	B	Jaromír Vaňhara	
<i>feminina</i> Negrobov, 1967	M	<i>Opetia</i> Meigen, 1830	
<i>flavipes</i> Meigen, 1824	B M	<i>nigra</i> Meigen, 1830	B M S
<i>glauca</i> Loew, 1869	M S	PLATYPEZIDAE	
<i>glauca</i> Kowarz, 1877	M S	Jaromír Vaňhara	
<i>impigra</i> Collin, 1941	B M	MICROSANIINAE	
<i>infumata</i> Loew, 1857	B	<i>Microsania</i> Zetterstedt, 1837	
<i>inspissata</i> Collin, 1952	S	<i>collarti</i> Chandler in Vaňhara, 1982	M
<i>jacula</i> (Fallén, 1823)	B M S	<i>pectipennis</i> (Meigen, 1830)	B M S
<i>jugalis</i> Collin, 1940	M S	<i>straeleni</i> Collart, 1954	M
<i>longicauda</i> Becker, 1917	S	CALLOMYIINAE	
<i>micacea</i> Loew, 1857	B M S	<i>Agathomyia</i> Verrall, 1901	
<i>mixta</i> Negrobov, 1967	B M S	<i>antennata</i> (Zetterstedt, 1819)	B M S
<i>muralis</i> Meigen, 1824	M S	<i>cinerea</i> (Zetterstedt, 1852)	S
<i>murina</i> Becker, 1917	B	<i>collini</i> Verrall, 1901	M
<i>nitida</i> Macquart, 1843	M	<i>elegantula</i> (Fallén, 1815)	B M S
<i>obscura</i> (Zetterstedt, 1838)	B S	<i>falleni</i> (Zetterstedt, 1819)	B M S
<i>pallipes</i> (Zetterstedt, 1843)	B S	<i>sexmaculata</i> v. Roser, 1840	M
<i>petrophiloides</i> Parent, 1925	M S	<i>setipes</i> Oldenberg, 1916	B S
<i>pinicola</i> Kowarz, 1879	B	<i>unicolor</i> Oldenberg, 1928	B M S
<i>plumbella</i> Meigen, 1824	B M S	<i>vernalis</i> Shatalkin, 1981	B M S
<i>signaticornis</i> Loew, 1857	B M	<i>viduella</i> (Zetterstedt, 1838)	B M S
<i>stackelbergiana</i> Negrobov, 1967	M	<i>wankowiczii</i> (Schnabl, 1884)	B M S
<i>tenuicauda</i> Loew, 1857	B M S		
<i>tertia</i> Becker, 1917	M		
<i>tristis</i> (Zetterstedt, 1838)	B M S		
<i>truncorum</i> Meigen, 1824	B M S		
<i>unisetosa</i> Collin, 1941	S		
<i>vagans</i> Becker, 1917	B S		
THRYPTICINI			
<i>Thrypticus</i> Gerstaecker, 1864			
<i>atomus</i> Frey, 1915	B M		

<i>woodella</i> Chandler in Hackman, 1980	M S	<i>subsultans</i> (Linnaeus, 1767)	B M S
Callomyia Meigen, 1804		Chaetopleurophora (Schmitz, 1922)	
<i>amoena</i> Meigen, 1824	B M S	<i>bohemani</i> (Becker, 1901)	B M S
<i>elegans</i> Meigen, 1804	B M S	<i>erythronota</i> (Strobl, 1892)	B M S
<i>speciosa</i> Meigen, 1824	B M S	<i>spinosior</i> Schmitz, 1938	M S
Platypezina Wahlgren, 1910		<i>spinosissima</i> (Strobl, 1892)	B S
<i>connexa</i> (Boheman, 1858)	B M	Conicera Meigen, 1830	
PLATYPEZINAE		<i>dauci</i> (Meigen, 1830)	B M S
Bolopus Enderlein, 1932		<i>floricola</i> Schmitz, 1938	B M S
<i>furcatus</i> (Fallén, 1826)	B M S	<i>schnittmanni</i> Schmitz, 1926	B M S
Kesselimyia Váňhara, 1981		<i>similis</i> (Haliday, 1833)	B M
<i>chandleri</i> Váňhara, 1981	M	<i>tarsalis</i> Schmitz, 1920	B M S
Lindneromyia Kessel, 1965		<i>tibialis</i> Schmitz, 1925	B M S
<i>dorsalis</i> (Meigen, 1804)	B M S	= <i>sorbis</i> Schmitz, 1936	
Paraplatypeza Kessel & Maggioncalda, 1968		Diplonevra Lioy, 1864	
<i>atra</i> (Meigen, 1804)	B M S	<i>abbreviata</i> (von Roser, 1840)	B M S
<i>bicincta</i> (Szilády, 1941)	B M S	<i>concinna</i> (Meigen, 1830)	B M S
Platypeza Meigen, 1803		<i>crassicornis</i> (Meigen, 1830)	B M S
<i>aterrima</i> Walker, 1836	M	<i>florea</i> (Fabricius, 1794)	B M S
<i>consobrina</i> Zetterstedt, 1844	B M S	<i>funnebris</i> (Meigen, 1830)	B M S
<i>fasciata</i> Meigen, 1804	B M S	<i>glabra</i> (Schmitz, 1927)	B M S
<i>hirticeps</i> Verrall, 1901	B M S	<i>lophochaeta</i> (Schmitz, 1927)	S
Polyporivora Kessel & Maggioncalda, 1968		<i>nitidula</i> (Meigen, 1830)	B M S
<i>boletina</i> (Fallén, 1815)	S	<i>pilosella</i> (Schmitz, 1927)	B M S
<i>ornata</i> (Meigen, 1838)	B M S	<i>sesquicornis</i> (Schmitz, 1927)	B M S
<i>picta</i> (Meigen, 1830)	S	<i>unisetalis</i> (Schmitz, 1935)	S
Protoclythia Kessel, 1949		Dohrniphora Dahl, 1898	
<i>modesta</i> (Zetterstedt, 1844)	B M S	<i>cornuta</i> (Bigot, 1856)	B S
<i>rufa</i> (Meigen, 1830)	B M S	Gymnophora Macquart, 1835	
Seri Kessel & Kessel, 1966		<i>arcuata</i> (Meigen, 1830)	B M S
<i>obscuripennis</i> (Oldenberg, 1916)	M	<i>healeyae</i> Disney, 1980	B M
PHORIDAE		<i>integralis</i> Schmitz, 1920	B M S
Bohuslav Mocek		<i>nigripennis</i> Schmitz, 1926	B M S
Aenigmatias Meinert, 1890		<i>quartomollis</i> Schmitz, 1920	B M S
<i>brevifrons</i> (Schmitz, 1955)	B	Gymnoptera Lioy, 1864	
<i>franzi</i> Schmitz, 1950	M	<i>longicostalis</i> Schmitz, 1933	B M
<i>lubbocki</i> (Verrall, 1877)	B	<i>vitripennis</i> (Meigen, 1830)	B M
Anevrina Lioy, 1864		Hypocera Lioy, 1864	
<i>curvinervis</i> (Becker, 1901)	B M S	<i>mordellaria</i> (Fallén, 1823)	B M S
<i>kozaneki</i> Brown, 1994	S	Megaselia Rondani, 1856	
<i>thoracica</i> (Meigen, 1804)	B M S	<i>aequalis</i> (Wood, 1909)	B M S
<i>unispinosa</i> (Zetterstedt, 1860)	B M S	<i>affinis</i> (Wood, 1909)	B M S
<i>urbana</i> (Meigen, 1830)	B M S	<i>albocingulata</i> (Strobl, 1906)	S
Beckerina Malloch, 1910		<i>alpina</i> Schmitz, 1965	S
<i>umbrimargo</i> (Becker, 1901)	B M S	<i>altifons</i> (Wood, 1909)	B M S
Billotia Schmitz, 1944		<i>analís</i> (Lundbeck, 1920)	B M
<i>inermis</i> Schmitz, 1944	B S	<i>angularis</i> (Schmitz, 1924)	B M S
Borophaga Enderlein, 1924		<i>aristica</i> (Schmitz, 1920)	M
<i>agilis</i> (Meigen, 1830)	B M S	<i>beckeri</i> (Wood, 1909)	B
<i>carinifrons</i> (Zetterstedt, 1848)	B M S	<i>berndseni</i> (Schmitz, 1919)	S
<i>erythrocerá</i> (Meigen, 1830)	B	<i>bovista</i> (Gimmerthal, 1848)	B M
<i>femorata</i> (Meigen, 1830)	B M S	<i>brevicostalis</i> (Wood, 1910)	B M S
<i>germanica</i> (Schmitz, 1918)	B M S	<i>brevisetá</i> (Wood, 1912)	S
<i>incrassata</i> (Meigen, 1830)	B M S	<i>breviterga</i> (Lundbeck, 1920)	B M
		<i>brunneicornis</i> (Schmitz, 1920)	M
		<i>brunneipenis</i> Costa, 1857	S
		<i>campestris</i> (Wood, 1908)	B M S

<i>ciliata</i> (Zetterstedt, 1848)	B	M	S	<i>mallochi</i> (Wood, 1909)	S
<i>cinereifrons</i> (Strobl, 1910)			S	<i>maura</i> (Wood, 1910)	S
<i>clara</i> (Schmitz, 1921)	B			<i>meconicera</i> (Speiser, 1925)	B M S
<i>coacta</i> (Lundbeck, 1920)		M	S	<i>meigeni</i> (Becker, 1901)	M S
<i>collini</i> (Wood, 1909)			S	<i>melanocephala</i> (von Roser, 1840)	B
<i>communifrons</i> (Schmitz, 1918)			S	<i>minor</i> (Zetterstedt, 1848)	B M S
<i>conformis</i> (Wood, 1909)	B	M	S	<i>minuta</i> (Aldrich, 1892)	M S
<i>consetigera</i> Schmitz, 1925			S	<i>mixta</i> (Schmitz, 1918)	M
<i>costalis</i> (von Roser, 1840)		M		<i>montana</i> Schmitz, 1935	S
<i>crassicosta</i> (Strobl, 1892)		M	S	<i>mortenseni</i> (Lundbeck, 1920)	S
<i>crassipes</i> (Wood, 1909)	B	M		<i>nasoni</i> (Malloch, 1914)	B M S
<i>dahli</i> (Becker, 1901)		M		<i>nigra</i> (Meigen, 1830)	B M S
<i>discreta</i> (Wood, 1909)			S	<i>nigriceps</i> (Loew, 1866)	B M S
<i>diversa</i> (Wood, 1909)		M		<i>obscuripennis</i> (Wood, 1909)	M S
<i>dubitalis</i> (Wood, 1908)	B		S	<i>paludosa</i> (Wood, 1908)	B S
<i>eccoptomera</i> Schmitz, 1927		M		<i>parva</i> (Wood, 1909)	B M S
<i>elongata</i> (Wood, 1914)		M	S	<i>pectoralis</i> (Wood, 1910)	S
<i>emarginata</i> (Wood, 1908)	B	M	S	<i>perfusca</i> Schmitz, 1935	M
<i>errata</i> (Wood, 1912)	B	M	S	<i>picta</i> (Lehmann, 1822)	B M S
<i>flava</i> (Fallén, 1823)	B	M	S	<i>pleuralis</i> (Wood, 1909)	B M S
<i>flavescens</i> (Wood, 1909)		M		<i>plurispinulosa</i> (Zetterstedt, 1860)	B S
<i>flavicans</i> Schmitz, 1935			S	<i>posticata</i> (Strobl, 1898)	S
<i>flavicoxa</i> (Zetterstedt, 1848)	B		S	<i>propinqua</i> (Wood, 1909)	B S
<i>frontalis</i> (Wood, 1909)		M	S	<i>protarsalis</i> Schmitz, 1927	M
<i>furva</i> Schmitz, 1929	B			<i>pulicaria</i> (Fallén, 1823)	B M S
<i>fusca</i> (Wood, 1909)			S	<i>pumila</i> (Meigen, 1830)	B M S
<i>fuscinervis</i> (Wood, 1908)	B	M	S	<i>pusilla</i> (Meigen, 1830)	B M S
<i>fuscopleuralis</i> Schmitz, 1929	B			<i>pygmaea</i> (Zetterstedt, 1848)	B M S
<i>fuscovariana</i> Schmitz, 1933			S	<i>quadriseta</i> (Schmitz, 1918)	B M S
<i>giraudi</i> (Egger, 1862)	B	M	S	<i>rivalis</i> (Wood, 1909)	S
<i>glabrifrons</i> (Wood, 1909)		M	S	<i>rubella</i> (Schmitz, 1920)	B S
<i>gregaria</i> (Wood, 1910)		M	S	<i>rudis</i> (Wood, 1909)	M
<i>groenlandica</i> (Lundbeck, 1901)			S	<i>rufa</i> (Wood, 1908)	M S
<i>halterata</i> (Wood, 1910)	B		S	<i>ruficornis</i> (Meigen, 1830)	B M S
<i>hirticus</i> (Schmitz, 1918)	B	M	S	<i>rufipes</i> (Meigen, 1804)	B M S
<i>hirtiventris</i> (Wood, 1909)		M	S	<i>rupestris</i> Schmitz, 1934	B
<i>hortensis</i> (Wood, 1909)			S	<i>scutellaris</i> (Wood, 1909)	M S
<i>humeralis</i> (Zetterstedt, 1838)	B	M	S	<i>sepulchralis</i> (Lundbeck, 1920)	B M S
<i>hyalipennis</i> (Wood, 1912)	B			<i>setulipalpis</i> Schmitz, 1938	S
<i>hypopygialis</i> (Lundbeck, 1920)			S	<i>simulans</i> (Wood, 1912)	B M S
<i>incrassata</i> (Schmitz, 1920)			S	<i>sordida</i> (Zetterstedt, 1838)	M S
<i>indifferens</i> (Lundbeck, 1920)		M	S	<i>spinicincta</i> (Wood, 1910)	S
<i>infraposa</i> (Wood, 1909)			S	<i>spinigera</i> (Wood, 1908)	B M
<i>intonsa</i> Schmitz, 1948	B	M		<i>stichata</i> (Lundbeck, 1920)	M S
<i>involuta</i> (Wood, 1910)	B	M	S	<i>stigmatica</i> (Schmitz, 1920)	M
<i>lata</i> (Wood, 1910)	B		S	<i>striolata</i> Schmitz, 1940	S
<i>latericia</i> Schmitz, 1935			S	<i>styloprocta</i> (Schmitz, 1921)	B M S
<i>latifemorata</i> (Becker, 1901)			S	<i>subnudipennis</i> (Schmitz, 1919)	S
<i>latifrons</i> (Wood, 1910)		M		<i>subpleuralis</i> (Wood, 1909)	M S
<i>limburgensis</i> (Schmitz, 1918)			S	<i>subtumida</i> (Wood, 1909)	B M S
<i>longicostalis</i> (Wood, 1912)			S	<i>sulphuripes</i> (Meigen, 1830)	S
<i>longipalpis</i> (Wood, 1910)	B	M	S	<i>superciliata</i> (Wood, 1910)	S
<i>longiseta</i> (Wood, 1909)		M		<i>tama</i> (Schmitz, 1926)	B M S
<i>lucifrons</i> (Schmitz, 1918)			S	<i>tarsalis</i> (Wood, 1910)	M
<i>lutea</i> (Meigen, 1830)	B	M	S	<i>tarsicia</i> Schmitz, 1926	M S
<i>luteipes</i> (Schmitz, 1918)	B	M	S	<i>tumida</i> (Wood, 1909)	B M S
<i>major</i> (Wood, 1912)	B	M	S	<i>uliginosa</i> (Wood, 1909)	S

<i>unguicularis</i> (Wood, 1909)	M S	<i>Stichilus</i> Enderlein, 1924	
<i>valvata</i> Schmitz, 1935	B S	<i>coronatus</i> (Becker, 1901)	B
<i>variana</i> Schmitz, 1926	B	<i>Triphleba</i> Rondani, 1856	
<i>verralli</i> (Wood, 1910)	S	<i>admirabilis</i> Schmitz, 1927	S
<i>vestita</i> (Wood, 1914)	S	<i>aequalis</i> (Schmitz, 1919)	B M
<i>virilis</i> (Schmitz, 1919)	S	<i>antricola</i> (Schmitz, 1918)	S
<i>woodi</i> (Lundbeck, 1922)	B M S	<i>aptina</i> (Schiner, 1853)	M
<i>xanthozona</i> (Strobl, 1892)	B M	<i>autumnalis</i> (Becker, 1901)	B M S
Menoziola Schmitz, 1927		<i>bicornuta</i> (Strobl, 1910)	B M S
<i>obscuripes</i> (Schmitz, 1927)	B	<i>bifidata</i> Schmitz, 1949	B
<i>schmitzi</i> (Menozi, 1921)	B M	<i>citrefiformis</i> (Becker, 1901)	B M S
Metopina Macquart, 1835		<i>crassinervis</i> (Strobl, 1910)	M
<i>braueri</i> (Strobl, 1880)	B M S	<i>distiguenda</i> (Strobl, 1892)	B M S
<i>crassinervis</i> Schmitz, 1920	S	<i>dudai</i> (Schmitz, 1918)	B M S
<i>formicomendicola</i> Schmitz, 1927	B	<i>excisa</i> (Lundbeck, 1921)	B
<i>galeata</i> (Haliday, 1833)7	B M S	<i>forfex</i> Schmitz, 1943	S
<i>heselhausi</i> Schmitz, 1914	B	<i>gracilis</i> (Wood, 1907)	S
<i>oligoneura</i> (Mik, 1967)	B M S	<i>hentrichi</i> Schmitz, 1934	S
<i>perpusilla</i> (Six, 1878)	B M S	<i>hyalinata</i> (Meigen, 1830)	B S
<i>pileata</i> Schmitz, 1936	B M S	<i>inaequalis</i> Schmitz, 1943	B S
<i>ulrichi</i> Disney, 1979	B M	<i>intermedia</i> (Malloch, 1908)	B M S
Phalacrotophora Enderlein, 1912		<i>lugubris</i> (Meigen, 1830)	B M S
<i>berolinensis</i> Schmitz, 1920	B M S	<i>luteifemorata</i> (Wood, 1906)	B
<i>delageae</i> Disney, 1979	B	<i>minuta</i> (Fabricius, 1787)	M S
<i>fasciata</i> (Fallén, 1823)	B M S	<i>nudipalpis</i> (Becker, 1901)	B M S
<i>spectabilis</i> Schmitz, 1925	S	<i>opaca</i> (Meigen, 1830)	B M S
Phora Latreille, 1796		<i>pachyneurella</i> (Schmitz, 1919)	S
<i>adducta</i> Schmitz, 1955	B	<i>palposa</i> (Zetterstedt, 1848)	S
<i>artifrons</i> Schmitz, 1920	M	<i>papillata</i> (Wingate, 1906)	B M S
<i>atra</i> (Meigen, 1804)	B M S	<i>smithi</i> Disney, 1982	M
<i>convallium</i> Schmitz, 1928	B M	<i>subcompleta</i> Schmitz, 1927	B M
<i>convergens</i> Schmitz, 1920	B M	<i>trinervis</i> (Becker, 1901)	B M
<i>dubia</i> (Zetterstedt, 1848)	B M S	<i>tumidula</i> (Schmitz, 1918)	S
<i>edentata</i> Schmitz, 1920	B M S	<i>vitrea</i> (Wood, 1906)	S
<i>hamata</i> Schmitz, 1927	B M S		
<i>holosericea</i> Schmitz, 1920	B M S		
<i>horrida</i> Schmitz, 1920	B M S		
<i>hyperborea</i> Schmitz, 1927	B M		
<i>indivisa</i> Schmitz, 1948	B M		
<i>limpida</i> Schmitz, 1935	B		
<i>obscura</i> (Zetterstedt, 1848)	B M S		
<i>penicillata</i> Schmitz, 1920	B M S		
<i>pubipes</i> Schmitz, 1920	B M		
<i>stictica</i> Meigen, 1830	B M S		
<i>tincta</i> Schmitz, 1920	B M S		
Plectanocnema Schmitz, 1926			
<i>nudipes</i> (Becker, 1901)	B M S		
Pseudacteon Coquillett, 1907			
<i>formicarium</i> (Verrall, 1877)	B M S		
<i>lundbecki</i> Schmitz, 1924	S		
Spiniphora Malloch, 1909			
<i>bergenstammi</i> (Mik, 1864)	B M S		
<i>dorsalis</i> (Becker, 1901)	B M		
<i>excisa</i> (Becker, 1901)	B M S		
<i>jugorum</i> (Schmitz, 1924)	B S		
<i>maculata</i> (Meigen, 1830)	B M		
<i>strobli</i> (Becker, 1901)	M S		
		SYRPHIDAE	
		Jiří Holinka & Libor Mazánek	
		SYRPHINAE	
		SYRPHINI	
		<i>Dasysyrphus</i> Enderlein, 1938	
		<i>albostrigatus</i> (Fallén, 1817)	B M S

<i>friuliensis</i> (van der Goot, 1960)	B M S	<i>erraticus</i> (Linnaeus, 1758)	B M S
<i>hilaris</i> (Zetterstedt, 1843)	B M S	= <i>annulipes</i> (Zetterstedt, 1838)	
<i>lenensis</i> Bagatshanova, 1980	B	Melangyna Verrall, 1901	
<i>pauxillus</i> (Williston, 1887)	B M S	<i>arctica</i> (Zetterstedt, 1838)	S
= <i>nigricornis</i> auct. nec Verrall, 1873		<i>barbifrons</i> (Fallén, 1817)	B M S
<i>pinastri</i> (De Geer, 1776)	B M S	<i>compositarum</i> (Verrall, 1873)	B M S
= <i>humidatus</i> (Meigen, 1822)		<i>ericarum</i> (Collin, 1946)	M S
<i>tricinctus</i> (Fallén, 1817)	B M S	<i>labiatarum</i> (Verrall, 1901)	B M S
<i>venustus</i> (Meigen, 1822)	B M S	<i>lasiophthalma</i> (Zetterstedt, 1843)	B M S
Didea Macquart, 1834		<i>lucifera</i> Nielsen, 1980	M
<i>alneri</i> (Fallén, 1817)	B M S	<i>quadrifasciata</i> (Verrall, 1873)	B M S
<i>fasciata</i> Macquart, 1834	B M S	<i>umbellatarum</i> (Fabricius, 1794)	B M S
<i>intermedia</i> Loew, 1854	B M S	Meligramma Frey, 1946	
Doros Meigen, 1803		<i>cingulata</i> (Egger, 1860)	S
<i>profuges</i> (Harris, 1780)	B M S	<i>gutkata</i> (Fallén, 1817)	B M S
= <i>conopseus</i> auct. nec Linnaeus, 1758		<i>triangulifera</i> (Zetterstedt, 1843)	B M S
Epistrophe Walker, 1852		Meliscaeva Frey, 1946	
<i>cryptea</i> Doczkal & Schmid, 1994	M	<i>auricollis</i> (Meigen, 1822)	B M S
<i>diaphana</i> (Zetterstedt, 1843)	B M S	<i>cinctella</i> (Zetterstedt, 1843)	B M S
<i>eligans</i> (Harris, 1780)	B M S	Parasyrphus Matsumura, 1917	
<i>flava</i> Doczkal & Schmid, 1994	B M S	<i>annulatus</i> (Zetterstedt, 1838)	B M S
= <i>melanostomoides</i> auct. nec Strobl, 1880		<i>lineola</i> (Zetterstedt, 1843)	B M S
= <i>ochrostoma</i> auct. nec Zetterstedt, 1849		<i>macularis</i> (Zetterstedt, 1843)	B M S
<i>grossulariae</i> (Meigen, 1822)	B M S	<i>malinellus</i> (Collin, 1952)	B M S
<i>leiophthalma</i> (Schiner & Egger, 1853)	B M S	<i>nigritarsis</i> (Zetterstedt, 1843)	B M S
<i>melanostoma</i> (Zetterstedt, 1843)	B M S	<i>punctulatus</i> (Verrall, 1873)	B M S
= <i>melanostomoides</i> (Strobl, 1880)		= <i>monticolus</i> (Becker, 1921)	
<i>nitidicollis</i> (Meigen, 1822)	B M S	<i>tarsatus</i> (Zetterstedt, 1838)	S
<i>ochrostoma</i> (Zetterstedt, 1849)	B M S	<i>vittiger</i> (Zetterstedt, 1843)	B M S
<i>similis</i> Doczkal & Schmid, 1994	B M	Scaeva Fabricius, 1805	
Epistrophella Dušek & Láska, 1967		<i>dignota</i> (Rondani, 1857)	B S
<i>euchroma</i> (Kowarz, 1885)	B M S	<i>pyrastris</i> (Linnaeus, 1758)	B M S
Episyrphus Matsumura & Adachi, 1917		<i>selenitica</i> (Meigen, 1822)	B M S
<i>balteatus</i> (De Geer, 1776)	B M S	Sphaerophoria Lepeletier & Serville, 1828	
Eriozona Schiner, 1860		<i>bankowskiae</i> Goeldlin, 1989	M
<i>syrphoides</i> (Fallén, 1817)	B M S	<i>batava</i> Goeldlin, 1974	B M S
Eupeodes Osten-Sacken, 1877		<i>estebani</i> Goeldlin, 1991	S
= <i>Metasyrphus</i> Matsumura, 1917		<i>fatarum</i> Goeldlin, 1989	B M
sg. Eupeodes s. str.		= <i>abbreviata</i> auct. nec Zetterstedt, 1859	
<i>corollae</i> (Fabricius, 1794)	B M S	<i>infuscata</i> Goeldlin, 1974	M
<i>flaviceps</i> (Rondani, 1857)	S	<i>menthastris</i> (Linnaeus, 1758)	B M S
<i>latifascianus</i> (Macquart, 1829)	B M S	<i>philanthus</i> (Meigen, 1822)	B M S
<i>latihumilatus</i> (Collin, 1931)	B M S	<i>rueppelli</i> (Wiedemann, 1820)	B M S
<i>lundbecki</i> (Soot-Ryen, 1946)	M	<i>scripta</i> (Linnaeus, 1758)	B M S
<i>luniger</i> (Meigen, 1822)	B M S	<i>shirchan</i> Violovitsh, 1957	M S
<i>nielsenii</i> Dušek & Láska, 1976	B M S	<i>taeniata</i> (Meigen, 1822)	B M S
<i>nitens</i> (Zetterstedt, 1843)	B M S	<i>virgata</i> Goeldlin, 1974	B M
sg. Lapposyrphus Dušek & Láska, 1967		Syrphus Fabricius, 1775	
<i>lapponicus</i> (Zetterstedt, 1838)	B M S	<i>ribesii</i> (Linnaeus, 1758)	B M S
Fagisyrphus Dušek & Láska, 1967		<i>torvus</i> Osten Sacken, 1875	B M S
<i>cinctus</i> (Fallén, 1817)	B M S	<i>vitripennis</i> Meigen, 1822	B M S
Ischyrosyrphus Bigot, 1882		Xanthogramma Schiner, 1860	
<i>glaucus</i> (Linnaeus, 1758)	B M S	<i>festivum</i> (Linnaeus, 1758)	B M S
<i>laternarius</i> (Müller, 1776)	B M S	= <i>citrofasciatum</i> (De Geer, 1776)	
Leucozona Schiner, 1860		<i>laetum</i> (Fabricius, 1794)	M S
<i>lucorum</i> (Linnaeus, 1758)	B M S	<i>pedissequum</i> (Harris, 1776)	B M S
Megasyrphus Dušek & Láska, 1967			

CHRYOTOSOXINI

Chrysotoxum Meigen, 1803

<i>arcuatum</i> (Linnaeus, 1758)	B M S
<i>festivum</i> auct. nec Linnaeus, 1758	
<i>hicinctum</i> (Linnaeus, 1758)	B M S
<i>cautum</i> (Harris, 1776)	B M S
<i>elegans</i> Loew, 1841	B M S
<i>fasciatum</i> (Müller, 1764)	B M S
<i>arcuatum</i> auct. nec Linnaeus, 1758	
<i>fasciolatum</i> (De Geer, 1776)	B M S
<i>intermedium</i> Meigen, 1822	B M S
<i>lineare</i> (Zetterstedt, 1819)	B S
<i>octomaculatum</i> Curtis, 1837	B M S
<i>vernale</i> Loew, 1841	B M S
<i>verralli</i> Colla, 1940	B M S

BACCHINI

Baccha Fabricius, 1805

<i>elongata</i> (Fabricius, 1775)	B M S
<i>obscuripennis</i> Meigen, 1822	B M S

MELANOSTOMATINI

Melanostoma Schiner, 1860

<i>chubium</i> (Zetterstedt, 1837)	B M S
<i>mellinum</i> (Linnaeus, 1758)	B M S
<i>scalare</i> (Fabricius, 1794)	B M S

Platycheirus Lepeletier & Serville, 1828

sg. <i>Pachysphyria</i> Enderlein, 1938	
<i>ambiguus</i> (Fallén, 1817)	B M S

sg. *Platycheirus* s. str.

<i>albimanus</i> (Fabricius, 1781)	B M S
<i>cyaneus</i> auct. nec Müller, 1764	
<i>angustatus</i> (Zetterstedt, 1843)	B M S
<i>clypeatus</i> (Meigen, 1822)	B M S
<i>complicatus</i> Becker, 1889	B M S
<i>discimanus</i> Loew, 1871	B M S
<i>europaeus</i> Goeldlin, Maibach & Speight, 1990	B M S
<i>fasciculatus</i> Loew, 1856	S
<i>fulviventris</i> (Macquart, 1829)	B M S
<i>immarginatus</i> (Zetterstedt, 1849)	M S
<i>manicatus</i> (Meigen, 1822)	B M S
<i>melanops</i> Loew, 1856	B M S
<i>occultus</i> Goeldlin, Maibach & Speight, 1990	B M S
<i>parvatus</i> Rondani, 1857	B M S
<i>ovalis</i> Becker, 1921	
<i>peltatus</i> (Meigen, 1822)	B M S
<i>perpallidus</i> Verrall, 1901	M S
<i>podagratus</i> (Zetterstedt, 1838)	B M S
<i>scambus</i> (Staecker, 1843)	B M S
<i>scutatus</i> (Meigen, 1822)	B M S
<i>sticticus</i> (Meigen, 1822)	B M S
<i>tarsalis</i> (Schummel, 1837)	B M S
<i>tatricus</i> Dušek & Láška, 1982	S
sg. <i>Pyrophæna</i> Schiner, 1860	
<i>granditarsis</i> (Förster, 1771)	B M S

rosarum (Fabricius, 1787)

B M S

Spazigaster Rondani, 1843*ambulans* (Fabricius, 1798)

B M S

Xanthandrus Verrall, 1901*comtus* (Harris, 1780)

B M S

PARAGINI

Paragus Latreille, 1804sg. *Pandasyophthalmus* Stuckenberg, 1954*haemorrhous* Meigen, 1822

B M S

sg. *Paragus* s. str.*albifrons* (Fallén, 1817)

B M S

bicolor (Fabricius, 1794)

B M S

finitimus Goeldlin, 1971

B M S

hermonensis Kaplan, 1981

M

majoranae Rondani, 1857

B M

quadrifasciatus Meigen, 1822

B M S

strigatus Meigen, 1822

B S

MILESIINAE

PIPIZINI

Heringia Rondani, 1856sg. *Heringia* s. str.*heringi* (Zetterstedt, 1843)

B M S

sg. *Neocnemodon* Goffe, 1944*brevidens* (Egger, 1865)

B

fulvimanus (Zetterstedt, 1843)

S

latitarsis (Egger, 1865)

B M S

pubescens (Delucchi & Pschorn-Walcher, 1955)

B M

vitripennis (Meigen, 1822)

B M S

Pipiza Fallén, 1810*austriaca* Meigen, 1822

B M S

bimaculata Meigen, 1822

B M S

fasciata Meigen, 1822

B M S

fenestrata Meigen, 1822

B S

festiva Meigen, 1822

B M S

lugubris (Fabricius, 1775)

B M S

huteitarsis Zetterstedt, 1843

M S

noctiluca (Linnaeus, 1758)

B M S

notata Meigen, 1822

M

quadrimaculata (Panzer, 1804)

B M S

signata Meigen, 1822

B S

Pipizella Rondani, 1856*annulata* (Macquart, 1829)

B M S

divicci (Goeldlin, 1974)

B M S

maculipennis (Meigen, 1822)

M S

viduata (Linnaeus, 1758)

B M S

= varipes (Meigen, 1822)*virens* (Fabricius, 1805)

B M S

Trichopsomyia Williston, 1888*carbonaria* (Meigen, 1822)

B M S

flavitaris (Meigen, 1822)

B M S

Triglyphus Loew, 1840*primus* Loew, 1840

B M S

CHEILOSIINI

Chamaesyrrhus Mik, 1895*scaevoldes* (Fallén, 1817) B M S**Cheilosia** Meigen, 1822*aerea* Dufour, 1848 M S*correcta* Becker, 1894*= zetterstedti* Becker, 1894*aestivalis* Becker, 1892 S*ahenea* (Roser, 1840) B M S*albipila* Meigen, 1838 B M S*albitarsis* (Meigen, 1822) B M S*antiqua* (Meigen, 1822) B M S*barbata* Loew, 1857 B M S*honesta* Rondani, 1868*bergenstammi* Becker, 1894 B M S*brachysoma* Egger, 1860 M S*caerulescens* (Meigen, 1822) B S*canicularis* (Panzer, 1801) B M S*similis* (Mühl, 1911)*carbonaria* Egger, 1860 B M S*chloris* (Meigen, 1822) B M S*chrysocoma* (Meigen, 1822) B M S*cynocephala* Loew, 1840 B M S*derasa* Loew, 1857 S*fasciata* Schiner & Egger, 1853 B M S*flavipes* (Panzer, 1798) B M S*fraterna* (Meigen, 1830) B M S*frontalis* Loew, 1857 B M S*gagatea* Loew, 1857 S*gigantea* (Zetterstedt, 1838) B M S*grisella* Becker, 1894 B M S*grossa* (Fallén, 1817) B M S*hercyniae* Loew, 1857 M S*illustrata* (Harris, 1780) B M S*impressa* Loew, 1840 B M S*impudens* Becker, 1894 M S*insignis* Loew, 1857 M S*laeviventris* Loew, 1857 S*lasiopa* Kowarz, 1885 B M S*= honesta* auct. nec Rondani, 1868*latifacies* Loew, 1857 M S*latifrons* (Zetterstedt, 1843) B M S*= intonsa* Loew, 1857*lens* Becker, 1894 B M S*= omissa* Becker, 1894*loewi* Becker, 1894 B M S*longifila* Becker, 1894 S*longula* (Zetterstedt, 1838) B M S*marginata* Becker, 1894 M S*melanopa* (Zetterstedt, 1843) B S*melanura* Becker, 1894 B M S*mixta* Becker, 1894 M S*montana* Egger, 1860 B M S*morio* (Zetterstedt, 1838) B M S*mutabilis* (Fallén, 1817) B M S*nebulosa* (Verrall, 1871) B M S*= langhofferi* Becker, 1894*nigripes* (Meigen, 1822) B M S*orthotricha* Vujić & Claussen, 1994 B M*pagana* (Meigen, 1822) B M S*pascuorum* Becker, 1894 M S*personata* Loew, 1758 M S*pictipennis* Egger, 1860 M S*pini* Becker, 1894 M S*polita* Becker, 1894 S*praecox* (Zetterstedt, 1843) B M S*= ruralis* auct. nec Meigen, 1822*proxima* (Zetterstedt, 1843) B M S*psilophthalma* Becker, 1894 S*pubera* (Zetterstedt, 1838) B M S*rhynchops* Egger, 1860 B M S*rotundiventris* Becker, 1894 M S*ruficollis* Becker, 1894 M S*rufimana* Becker, 1894 M S*ruralis* (Meigen, 1822) M*= globulipes* Becker, 1894*sahlbergi* Becker, 1894 M S*scanica* (Ringdahl, 1937) M S*scutellata* (Fallén, 1817) B M S*semifasciata* Becker, 1894 B M S*soror* (Zetterstedt, 1843) B M S*variabilis* (Panzer, 1798) B M S*velutina* Loew, 1840 B M S*vernalis* (Fallén, 1817) B M S*vicina* (Zetterstedt, 1849) B M S*= nasutula* Becker, 1894*vulpina* (Meigen, 1822) B M S*= conops* Becker, 1894**Ferdinandea** Rondani, 1844*cuprea* (Scopoli, 1763) B M S*nigrifrons* (Egger, 1860) B M S*ruficornis* (Fabricius, 1775) B M S**Pelecocera** Meigen, 1822*latifrons* Loew, 1856 M S*tricincta* Meigen, 1822 B S**Portevinia** Goffe, 1944*maculata* (Fallén, 1817) M S**Rhingia** Scopoli, 1763*borealis* Ringdahl, 1928 B M S*= austriaca* auct. nec Meigen, 1830*campestris* Meigen, 1822 B M S*rostrata* (Linnaeus, 1758) B M S

CALLICERINI

Callicera Panzer, 1809*aenea* (Fabricius, 1781) B M S

VOLUCCELLINI

Volucella Geoffroy, 1762*bombylans* (Linnaeus, 1758) B M S*inanis* (Linnaeus, 1758) B M S*inflata* (Fabricius, 1794) B M S

<i>pellucens</i> (Linnaeus, 1758)	B M S	<i>clavata</i> (Scopoli, 1763)	S
<i>zonaria</i> (Poda, 1761)	B M S	<i>chunipes</i> (Fallén, 1816)	B M S
CHRYSOGASTERINI		<i>elegans</i> Schummel, 1843	B M S
<i>Brachyopa</i> Meigen, 1822		= <i>kimakowiczi</i> Strobl, 1897	
<i>bicolor</i> (Fallén, 1817)	B M S	<i>latifrons</i> Egger, 1865	B M S
<i>dorsata</i> Zetterstedt, 1837	B M S	<i>montana</i> Becker, 1921	B M S
<i>insensilis</i> Collin, 1939	M S	<i>sibirica</i> Stackelberg, 1953	M S
<i>maculipennis</i> Thompson, 1980	M S	<i>spheginea</i> (Zetterstedt, 1838)	B
<i>panzeri</i> Goffe, 1945	M S	<i>verecunda</i> Collin, 1937	B M S
<i>pilosa</i> Collin, 1939	B M		
<i>plena</i> Collin, 1939	B	SERICOMYIINI	
<i>scutellaris</i> Robineau-Desvoidy, 1843	B M	<i>Arctophila</i> Schiner, 1860	
<i>testacea</i> (Fallén, 1817)	M S	<i>bombiformis</i> (Fallén, 1810)	B M S
<i>vittata</i> Zetterstedt, 1843	M S	<i>superbiens</i> Müller, 1776	B M S
Chrysogaster Meigen, 1803		= <i>fulva</i> (Harris, 1780)	
<i>cemiteriorum</i> (Linnaeus, 1758)	B M S	= <i>mussitans</i> (Fabricius, 1777)	
= <i>chalybeata</i> Meigen, 1822		Sericomyia Meigen, 1803	
<i>solstitialis</i> (Fallén, 1817)	B M S	<i>lappona</i> (Linnaeus, 1758)	B M S
<i>virescens</i> Loew, 1854	M	<i>silentis</i> (Harris, 1776)	B M S
Hammerschmidtia Schummel, 1834			
<i>ferruginea</i> (Fallén, 1817)	S	EUMERINI	
Lejogaster Roncani, 1857		<i>Eumerus</i> Meigen, 1822	
<i>metallina</i> (Fabricius, 1781)	B M S	<i>flavitaris</i> Zetterstedt, 1843	B M S
<i>tarsata</i> (Meigen, 1822)	B M S	<i>grandis</i> Meigen, 1822	S
= <i>splendida</i> (Meigen, 1822)		<i>longicornis</i> Loew, 1855	S
Melanogaster Rondani, 1857		<i>ornatus</i> Meigen, 1822	B S
<i>aerosa</i> (Loew, 1843)	M	<i>ovatus</i> Loew, 1848	B S
<i>hirtella</i> (Loew, 1843)	B M	<i>ruficornis</i> Meigen, 1822	B
<i>nuda</i> (Macquart, 1829)	B M S	<i>sabulorum</i> (Fallén, 1817)	B M S
= <i>viduata</i> auct. nec Linnaeus, 1758		<i>sinuatus</i> Loew, 1855	M S
<i>parumplicata</i> (Loew, 1840)	B M S	<i>sogdianus</i> Stackelberg, 1952	B M S
= <i>macquarti</i> Loew, 1843		<i>strigatus</i> (Fallén, 1817)	B M S
Myolepta Newman, 1838		<i>tarsalis</i> Loew, 1848	S
<i>luteola</i> (Gmelin, 1790)	B M S	<i>tricolor</i> (Fabricius, 1798)	B M S
<i>obscura</i> Becher, 1882	S	<i>tuberculatus</i> Rondani, 1857	B M S
<i>vara</i> (Panzer, 1798)	S	Merodon Meigen, 1803	
Orthonevra Macquart, 1829		<i>aberrans</i> Egger, 1860	S
<i>brevicornis</i> (Loew, 1843)	B M S	<i>aeneus</i> Meigen, 1822	S
<i>elegans</i> (Meigen, 1822)	B	<i>armipes</i> Rondani, 1843	B M S
<i>frontalis</i> (Loew, 1843)	B S	<i>auripes</i> Sack, 1913	S
<i>geniculata</i> (Meigen, 1830)	B M S	<i>avidus</i> (Rossi, 1790)	B M S
<i>nobilis</i> (Fallén, 1817)	B M S	= <i>spinipes</i> (Fabricius, 1794)	
<i>plumbago</i> (Loew, 1840)	M S	<i>cinereus</i> (Fabricius, 1794)	S
<i>splendens</i> (Meigen, 1822)	M	<i>clavipes</i> (Fabricius, 1781)	B M S
Neoascia Williston, 1886		<i>constans</i> (Rossi, 1794)	S
<i>annexa</i> (Müller, 1776)	B M S	<i>crymensis</i> Paramanov, 1925	S
= <i>floralis</i> auct. nec Meigen, 1822		<i>equestris</i> (Fabricius, 1794)	B M S
<i>geniculata</i> (Meigen, 1822)	B M	<i>haemorrhoidalis</i> Sack, 1913	S
<i>interrupta</i> (Meigen, 1822)	B M S	<i>ruficornis</i> Meigen, 1822	S
<i>meticulosa</i> (Scopoli, 1763)	B	<i>rufus</i> Meigen, 1838	B M S
= <i>aenea</i> (Meigen, 1822)		Psilota Meigen, 1822	
<i>obliqua</i> Coe, 1940	B M S	<i>anthracina</i> Meigen, 1822	M S
<i>podagrica</i> (Fabricius, 1775)	B M S	<i>innupta</i> Rondani, 1857	S
<i>tenur</i> (Harris, 1780)	B M S		
= <i>dispar</i> auct. nec Meigen, 1822		PSARINI	
Sphegina Meigen, 1822		<i>Psarus</i> Latreille, 1804	
		<i>abdominalis</i> (Fabricius, 1794)	B M S

<i>segnis</i> (Linnaeus, 1758)	B M S	<i>Cephalosphaera</i> Enderlein, 1936	
<i>semulatra</i> (Harris, 1790)	B M S	<i>furcata</i> (Egger, 1860)	B M S
= <i>abiens</i> Meigen, 1822		<i>germanica</i> Aczél, 1940	M
<i>sylvarium</i> (Linnaeus, 1758)	B M S	<i>Cephalops</i> Fallén, 1810	
<i>tarda</i> Meigen, 1822	B M S	<i>aeneus</i> Fallén, 1810	B M S
<i>xanthocnema</i> Collin, 1939	B M S	<i>carinatus</i> (Verrall, 1901)	B M S
		<i>obtusinervis</i> (Zetterstedt, 1844)	B M S
MICRODONTINAE		<i>perspicuus</i> (de Meijere, 1905)	M S
<i>Microdon</i> Meigen, 1803		<i>semifumosus</i> (Kowarz, 1887)	B M S
<i>devius</i> (Linnaeus, 1761)	B M S	<i>signatus</i> (Becker, 1900)	B M S
<i>eggeri</i> Mik, 1897	B M S	<i>subultimus</i> Collin, 1956	B M S
<i>latifrons</i> Loew, 1856	M S	<i>ultimus</i> (Becker, 1900)	B M S
<i>mutabilis</i> (Linnaeus, 1758)	B M S	<i>vittipes</i> (Zetterstedt, 1844)	B M S
		<i>Beckerias</i> Aczél, 1939	
PIPUNCULIDAE		<i>pannonicus</i> Aczél, 1939	B M S
Pavel Lauterér		<i>Microcephalops</i> de Meyer, 1989	
		<i>vestitus</i> (Becker, 1900)	B M
CHALARINAE		<i>Eudorylas</i> Aczél, 1940	
<i>Chalarus</i> Walker, 1834		<i>bartaki</i> Kozánek, 1993	S
<i>argentatus</i> Coe, 1966	B	<i>demeyeri</i> Kozánek, 1993	B
<i>basalis</i> Loew, 1873	B M	<i>electus</i> (Hardy, 1947)	B
<i>brevicaudis</i> Jervis, 1992	B M	<i>elephas</i> (Becker, 1898)	M S
<i>decorus</i> Jervis, 1992	M	<i>fascipes</i> (Zetterstedt, 1844)	B M S
<i>fimbriatus</i> Coe, 1966	B M S	<i>furvulus</i> Collin, 1956	M
<i>gynocephalus</i> Jervis, 1992	M	<i>fuscipes</i> (Zetterstedt, 1844)	B M S
<i>latifrons</i> Hardy, 1943	B M S	<i>fuscus</i> (Zetterstedt, 1844)	B M
<i>parmenteri</i> Coe, 1966	M	<i>horridus</i> (Becker, 1898)	M S
<i>pughi</i> Coe, 1966	B M S	<i>imperfectus</i> (Becker, 1921)	S
<i>spurius</i> (Fallén, 1818)	B M S	<i>inferus</i> Collin, 1956	B S
<i>Verrallia</i> Mik, 1899		<i>jenkinsoni</i> Coe, 1966	M S
<i>aucta</i> (Fallén, 1817)	B M S	<i>kowarzi</i> (Becker, 1898)	S
<i>Jassidophaga</i> Aczél, 1939		<i>longifrons</i> Coe, 1966	B S
<i>beatricis</i> (Coe, 1966)	B M S	<i>melanostolus</i> (Becker, 1898)	M
<i>pilosa</i> (Zetterstedt, 1838)	B S	<i>montium</i> (Becker, 1898)	B M S
<i>setosa</i> (Verrall, 1901)	B M S	<i>nemoralis</i> Kozánek, 1993	S
<i>villosa</i> (von Roser, 1840)	B M S	<i>obliquus</i> Coe, 1966	B M S
		<i>obscurus</i> Coe, 1966	B M S
NEPHROCERINAE		<i>opacus</i> (Fallén, 1816)	B M S
<i>Nephrocerus</i> Zetterstedt, 1838		<i>restriclus</i> Coe, 1966	B M S
<i>flavicornis</i> Zetterstedt, 1844		<i>ruralis</i> (Meigen, 1824)	B M S
<i>lapponicus</i> Zetterstedt, 1838	B M	<i>slovacus</i> Kozánek, 1993	M S
<i>scutellatus</i> (Macquart, 1834)	B S	<i>subfascipes</i> Collin, 1956	B M S
		<i>subterminalis</i> Collin, 1956	B M S
PIPUNCULINAE		<i>terminalis</i> (Thomson, 1869)	B M S
<i>Pipunculus</i> Latreille, 1802		<i>zermattensis</i> (Becker, 1898)	B M S
<i>calceatus</i> von Roser, 1840	B M S	<i>zonatus</i> (Zetterstedt, 1849)	M
<i>campestris</i> Latreille, 1804	B M S	<i>zonellus</i> Collin, 1956	B M S
<i>fonsecar</i> Coe, 1966	S	<i>Dorylomorpha</i> Aczél, 1939	
<i>lichtwardti</i> Kozánek, 1981	S	sg. <i>Dorylomyia</i> Albrecht, 1990	
<i>oldenbergi</i> Collin, 1956	B M S	<i>beckeri</i> (Aczél, 1935)	B M S
<i>omissinervis</i> Becker, 1889	S	<i>incognita</i> (Verrall, 1901)	B M S
<i>spinipes</i> Meigen, 1830	B M S	<i>xanthocera</i> (Kowarz, 1887)	B M S
<i>tenuirostris</i> Kozánek, 1981	B M S	sg. <i>Dorylomyza</i> Albrecht, 1990	
<i>thomsoni</i> Becker, 1898	B M S	<i>albitarsis</i> (Zetterstedt, 1844)	B M S
<i>varipes</i> Meigen, 1824	B M S	<i>clavifemora</i> Coe, 1966	M
<i>zugmayeriae</i> Kowarz, 1887	B M S	<i>fennica</i> Albrecht, 1979	B M S
		<i>hungarica</i> (Aczél, 1939)	B M S

<i>infirmata</i> (Collin, 1937)	
<i>lautereri</i> Albrecht, 1990	M S
<i>platystylis</i> Albrecht, 1979	M S
<i>praetermissa</i> Albrecht, 1979	S
<i>xanthopus</i> (Thomson, 1870)	B M S
sg. <i>Dorylomorpha</i> s. str.	
<i>aczei</i> Hardy, 1947	B M S
<i>confusa</i> (Verrall, 1901)	B M S
<i>extricata</i> (Collin, 1937)	B M S
<i>imparata</i> (Collin, 1937)	B M S
<i>rufipes</i> (Meigen, 1824)	M S
sg. <i>Pipunculina</i> Albrecht, 1990	
<i>borealis</i> (Wahlgren, 1910)	M S
<i>strandii</i> (Duda, 1940)	
<i>maculata</i> (Walker, 1834)	B M S
<i>Tomosvaryella</i> Aczél, 1939	
<i>eilitarsis</i> (Strobl, 1910)	M
<i>coquillettii</i> (Kertész, 1907)	B M S
<i>geniculata</i> (Meigen, 1824)	B M S
<i>kuthyi</i> Aczél, 1944	M S
<i>minima</i> (Becker, 1898)	S
<i>palliditarsis</i> (Collin, 1931)	B M S
<i>sylvatica</i> (Meigen, 1824)	B M S

PSEUDOPOMYZIDAE

Jindřich Roháček

<i>Pseudopomyza</i> Strobl, 1893	
<i>atrinana</i> (Meigen, 1830)	B M S

MICROPEZIDAE

Jindřich Roháček

MICROPEZINAE

<i>Micropeza</i> Meigen, 1803	
<i>brevipennis</i> von Roser, 1840	B M S
<i>corrigiolata</i> (Linnaeus, 1767)	B M S

CALOBATINAE

<i>Calobata</i> Meigen, 1803	
<i>petronella</i> (Linnaeus, 1761)	B M S
<i>Calobatella</i> Mik, 1898	
<i>longiceps</i> (Loew, 1870)	B M S
<i>Cnodacophora</i> Czerny, 1930	
<i>sellata</i> (Meigen, 1826)	B M S
<i>stylifera</i> (Loew, 1870)	S
<i>mutans</i> (Soós, 1975)	
<i>Compsobata</i> Czerny, 1930	
<i>Trilophyrobata</i> Hennig, 1938	
<i>cibaria</i> (Linnaeus, 1761)	B M S
<i>femoralis</i> (Meigen, 1826)	B M S
<i>commutata</i> (Czerny, 1930)	B M S
– <i>nigricornis</i> auct. nec Zetterstedt, 1838	
<i>Neria</i> Robineau-Desvoidy, 1830	
<i>ephippium</i> (Fabricius, 1794)	B M S

TAENIAPTERINAE

<i>Rainieria</i> Rondani, 1843	
<i>calceata</i> (Fallén, 1820)	B M S
<i>latifrons</i> (Loew, 1870)	S

TANYPEZIDAE

Jindřich Roháček	
<i>Tanypeza</i> Fallén, 1820	
<i>longimana</i> Fallén, 1820	B M S

STRONGYLOPHTHALMYIIDAE

Jindřich Roháček	
<i>Strongylophthalmyia</i> Heller, 1902	
<i>ustulata</i> (Zetterstedt, 1847)	B M S

MEGAMERINIDAE

Jindřich Roháček	
<i>Megamerina</i> Rondani, 1861	
<i>dolium</i> (Fabricius, 1805)	B M S

PSILIDAE

Vladislav Martinek

Chamaepsila Hendel, 1917

sg. <i>Chamaepsila</i> s.str.	
<i>atra</i> (Meigen, 1826)	B M S
<i>bicolor</i> (Meigen, 1826)	B M S
– <i>nigromaculata</i> (Strobl, 1909)	
<i>chunalis</i> (Collin, 1944)	B M S
<i>gracilis</i> (Meigen, 1826)	B M S
<i>humeralis</i> (Zetterstedt, 1847)	B M S
<i>kaszabi</i> (Soós, 1974)	B
<i>limbatella</i> (Zetterstedt, 1847)	B M S
<i>huteola</i> (Collin, 1944)	B S
<i>martineki</i> Carles-Tolrà, 1993	B
<i>morio</i> (Zetterstedt, 1835)	S
<i>nigra</i> (Fallén, 1820)	B M S
<i>nigricornis</i> (Meigen, 1826)	B M S
<i>nigrosetosa</i> Frey, 1925	B M S
<i>pallida</i> (Fallén, 1820)	B M S
<i>pectoralis</i> (Meigen, 1826)	B M S
<i>persimilis</i> (Wakerley, 1959)	B M S
<i>pseudobicolor</i> Soós, 1985	B
<i>rosae</i> (Fabricius, 1794)	B M S
<i>rufa</i> (Meigen, 1826)	B M S
<i>unilineata</i> (Zetterstedt, 1847)	B S
<i>villosula</i> (Meigen, 1826)	S
sg. <i>Tetrapsila</i> Frey, 1925	
<i>obscuritarsis</i> (Loew, 1856)	B M S
<i>Chyliza</i> Fallén, 1820	

sg. <i>Chyliza</i> s.str.			MYOPINAE	
<i>annulipes</i> Macquart, 1835	B	M	ZODIONINI	
<i>leptogaster</i> (Panzer, 1798)	B	M	<i>Zodion</i> Latreille, 1796	
<i>nova</i> Collin, 1944	B	M	<i>cinereum</i> (Fabricius, 1794)	B M S
<i>vittata</i> Meigen, 1826	B	M	<i>notatum</i> (Meigen, 1804)	B M S
sg. <i>Dasyne</i> Robineau-Desvoidy, 1830			MYOPINI	
<i>extenuata</i> (Rossi, 1790)	M	S	<i>Myopa</i> Fabricius, 1775	
<i>Loxocera</i> Meigen, 1803			<i>buccata</i> (Linnaeus, 1758)	B M S
sg. <i>Loxocera</i> s.str.			<i>dorsalis</i> Fabricius, 1794	B M S
<i>albiseta</i> (Schränk, 1803)	B	M	<i>extricata</i> Collin, 1960	B M S
<i>aristata aristata</i> (Panzer, 1801)	B	M	<i>fasciata</i> Meigen, 1804	B M S
<i>aristata maculata</i> Rondani, 1876		S	<i>morio</i> Meigen, 1804	B M S
<i>fulviventris</i> Meigen, 1826	B	M	<i>occulta</i> Wiedemann in Meigen, 1824	B M S
<i>sylvatica</i> Meigen, 1826	B	M	<i>picta</i> Panzer, 1798	B M S
sg. <i>Platystyla</i> Macquart, 1835			<i>polystigma</i> Rondani, 1857	B M S
<i>hoffmannseggii</i> Meigen, 1826	M	S	<i>stigma</i> Meigen, 1824	B M S
<i>Oxypsila</i> Frey, 1925			<i>strandii</i> Duda, 1940	B M S
<i>abdominalis</i> (Schummel, 1844)	B	M	<i>tessellatipennis</i> Motschulsky, 1859	M S
<i>Psila</i> Meigen, 1803			<i>testacea</i> (Linnaeus, 1767)	B M S
<i>finetaria</i> (Linnaeus, 1761)	B	M	<i>variegata</i> Meigen, 1804	B M S
<i>merdaria</i> Collin, 1944	B	M	<i>vicaria</i> Walker, 1849	B S
<i>Psilosoma</i> Zetterstedt, 1860			<i>Melanosoma</i> Robineau-Desvoidy, 1853	
<i>audouini</i> (Zetterstedt, 1835)	B	M	<i>bicolor</i> (Meigen, 1824)	M S
<i>lefebvrei</i> (Zetterstedt, 1835)	B	M	<i>Myopotta</i> Zimina, 1969	
			<i>pallipes</i> (Wiedemann in Meigen, 1824)	S
CONOPIDAE			<i>rubripes</i> (Villeneuve, 1909)	B M S
Milan Chvála			<i>Thecophora</i> Rondani, 1845	
CONOPINAE			<i>atra</i> (Fabricius, 1775)	B M S
<i>Abrachyglossum</i> Kröber, 1919			<i>distincta</i> (Wiedemann in Meigen, 1824)	B M S
<i>capitatum</i> (Loew, 1847)	B	M	<i>fulvipes</i> (Robineau-Desvoidy, 1830)	B M S
<i>Conops</i> Linnaeus, 1758			<i>longirostris</i> Lyneborg, 1962 ?	
<i>ceriaeformis</i> Meigen, 1824	B	M	<i>melanopa</i> Rondani, 1857	M S
<i>flavifrons</i> Meigen, 1824		S	<i>pusilla</i> (Meigen, 1824)	B M S
<i>flavipes</i> Linnaeus, 1758	B	M	SICINI	
<i>longiventris</i> Kröber, 1916		M	<i>Sicus</i> Scopoli, 1763	
<i>quadrifasciatus</i> De Geer, 1776	B	M	<i>abdominalis</i> Kröber, 1915	B M
<i>scutellatus</i> Meigen, 1804	B	M	<i>femoralis</i> Rondani, 1865	B
<i>strigatus</i> Wiedemann in Meigen, 1824	B	M	<i>ferrugineus</i> (Linnaeus, 1761)	B M S
<i>vesicularis</i> Linnaeus, 1761	B	M	<i>fusenensis</i> Ouchi, 1939	M S
<i>Leopoldius</i> Rondani, 1843			<i>nigritarsis</i> Zimina, 1975	M
<i>brevirostris</i> (Germar, 1827)	B	M	DALMANIINAE	
<i>calceatus</i> (Rondani, 1857)	B	M	<i>Dalmannia</i> Robineau-Desvoidy, 1830	
<i>coronatus</i> (Rondani, 1857)	B	M	<i>aculeata</i> (Linnaeus, 1761)	M S
<i>diadematus</i> Rondani, 1845		S	<i>dorsalis</i> (Fabricius, 1794)	B
<i>signatus</i> (Wiedemann in Meigen, 1824)	B	M	<i>marginata</i> (Meigen, 1824)	B M S
<i>Physocephala</i> Schiner, 1861			<i>punctata</i> (Fabricius, 1794)	B M S
<i>chrysorrhoea</i> (Meigen, 1824)	B	M	LONCHAEIDAE	
<i>lacera</i> (Meigen, 1824)		S	Jan Máca	
<i>laticincta</i> (Brullé, 1832)	B	M	DASIOPINAE	
<i>nigra</i> (De Geer, 1776)	B	M	<i>Dasiops</i> Rondani, 1856	
<i>pusilla</i> (Meigen, 1824)		M	<i>appendiculus</i> Morge, 1959	B S
<i>rufipes</i> (Fabricius, 1781)	B	M		
<i>truncata</i> (Loew, 1847)		M		
<i>vittata</i> (Fabricius, 1794)	B	M		

<i>calvus</i> Morge, 1959		S	<i>sororcula</i> Hackman, 1956		B
<i>facialis</i> Collin, 1953	B	M S	<i>stackelbergi</i> Czerny, 1934		B M
<i>hennigi</i> Morge, 1959	B		<i>subneatos</i> Kovalev, 1974		B
<i>latifrons</i> (Meigen, 1826)		M S	<i>sylvatica</i> Beling, 1873		B M S
<i>mucronatus</i> Morge, 1959	B	M	<i>tarsata</i> Fallén, 1820		B S
<i>nocturnus</i> Morge, 1959	B		<i>ultima</i> Collin, 1953		M
<i>perpropinquus</i> Morge, 1959	B	S	<i>zetterstedti</i> Becker, 1902		B M
<i>spatiosus</i> (Becker, 1895)	B	M S	<i>Setisquamalonchaea</i> Morge, 1963		
<i>trichosternalis</i> Morge, 1959	B		<i>fumosa</i> (Egger, 1862)		B M S
			<i>retalionis</i> (McAlpine, 1964)		B M S
LONCHAEINAE					
EAROMYIINI					
<i>Chaetolonchaea</i> Czerny, 1934			PALLOPTERIDAE		
<i>brevipilosa</i> Czerny, 1934		S	Vladislav Martinek		
<i>dasyops</i> (Meigen, 1826)	B	M S	EURYGNATHOMYIINAE		
<i>pallipennis</i> (Zetterstedt, 1855)	B	M	<i>Eurygnathomyia</i> Czerny, 1904		
<i>Earomyia</i> Zetterstedt, 1842			<i>bicolor</i> (Zetterstedt, 1837)	B	S
<i>adriatica</i> (Becker, 1895)		S			
<i>albifacies</i> (Czerny, 1934)	B		PALLOPTERINAE		
<i>crystallophila</i> (Becker, 1895)	B	S	<i>Palloptera</i> Fallén, 1820		
<i>impossibile</i> Morge, 1959	B	M S	<i>flava</i> Oldenberg, 1910	B	M S
<i>inquilina</i> (Hendel in Seitner, 1929)		M	<i>marginata</i> (Meigen, 1826)	B	M S
<i>lonchaeoides</i> Zetterstedt, 1848	B		<i>umbellatarum</i> (Fabricius, 1775)	B	M S
<i>schistopyga</i> Collin, 1953	B		<i>ustulata</i> Fallén, 1820	B	M S
<i>tomskensis</i> Morge, 1959	B		<i>Temnosira</i> Enderlein, 1936		
<i>viridana</i> (Meigen, 1826)	B	M S	<i>ambusta</i> (Meigen, 1826)	B	M S
<i>virilis</i> Collin, 1953	B	S	<i>laetabilis</i> (Loew, 1873)	B	M S
<i>Protearomyia</i> McAlpine, 1962			<i>saltuum</i> (Linnaeus, 1758)	B	M S
<i>nigra</i> (Meigen, 1826)	B	M S	<i>Toxoneura</i> Macquart, 1835		
LONCHAEINI					
<i>Lonchaea</i> Fallén, 1820			<i>basimaculata</i> (Czerny, 1934)	B	M
<i>britteni</i> Collin, 1953	B	M	<i>ephippium</i> (Zetterstedt, 1860)		S
<i>bruggeri</i> Morge, 1967	B		<i>modesta</i> (Meigen, 1830)	B	M S
<i>carpathica</i> Kovalev, 1974	B	S	<i>quinquemaculata</i> (Macquart, 1835)	B	M S
<i>caucasica</i> Kovalev, 1974	B		<i>trimacula</i> (Meigen, 1826)	B	M S
<i>chorea</i> (Fabricius, 1781)	B	M S	<i>usta</i> (Meigen, 1826)	B	M S
<i>collini</i> Hackman, 1956	B		<i>venusta</i> (Loew, 1858)		S
<i>contigua</i> Collin, 195	B	S	PIOPHILIDAE		
<i>contraria</i> Czerny, 1934	B	M S	Jan Zuska & Miroslav Barták		
<i>corusca</i> Czerny, 1934	B	M	PIOPHILINAE		
<i>deutschii</i> Zetterstedt, 1838		M	<i>Allopiophila</i> Hendel, 1917		
<i>fugax</i> Becker, 1895	B	M S	<i>luteata</i> (Haliday, 1833)	B	M S
<i>gorodkovi</i> Kovalev, 1974	B		<i>Amphipogon</i> Wahlberg, 1845		
<i>hirticeps</i> Zetterstedt, 1838		M	<i>flavum</i> (Zetterstedt, 1838)	B	S
<i>krivosheinae</i> Kovalev, 1973		M	<i>Liopiophila</i> Duda, 1924		
<i>laticornis</i> Meigen, 1826	B	M S	<i>varipes</i> (Meigen, 1830)	B	M S
<i>laxa</i> Collin, 1953		S	<i>Mycetaulus</i> Loew, 1845		
<i>limatula</i> Collin, 1953	B	M S	<i>bipunctatus</i> (Fallén, 1823)	B	M S
<i>nitens</i> (Bigot, 1885)		M	<i>latipennis</i> Ozerov & Barták, 1993	B	
<i>obscuritarsis</i> Collin, 1953	B	M	<i>Parapiophila</i> McAlpine, 1977		
<i>palposa</i> Zetterstedt, 1847	B	M S	<i>pectiniventris</i> (Duda, 1924)		M
<i>patens</i> Collin, 1953	B	S	<i>vernica</i> Ozerov & Barták, 1993	B	M S
<i>peregrina</i> Becker, 1895	B	M	<i>vulgaris</i> (Fallén, 1820)	B	M S
<i>postica</i> Collin, 1953	B	M S	<i>Piophila</i> Fallén, 1810		
<i>scutellaris</i> Rondani, 1874	B	M S	<i>casei</i> (Linnaeus, 1758)	B	M S
<i>seitneri</i> Hendel, 1928	B	M			

<i>Prochyliza</i> Walker, 1849			<i>Systata</i> Loew, 1868		
<i>nigricornis</i> (Meigen, 1826)	B	M S	<i>rivularis</i> (Fabricius, 1805)	B	S
<i>nigrimana</i> (Meigen, 1826)	B	M S			
<i>Protopiophila</i> Duda, 1924			ULIDIIDAE		
<i>latipes</i> (Meigen, 1838)	B	M S	Jan Zuska		
<i>Stearibia</i> Lioy, 1864			<i>Euxesta</i> Loew, 1868		
<i>nigriceps</i> (Meigen, 1826)	B	M S	<i>pechumani</i> Curran, 1938		S
THYREOPHORINAE			<i>Physiphora</i> Fallén, 1810		
<i>Centrophlebomyia</i> Hendel, 1903			<i>alceae</i> (Preyssler, 1791)	B	M S
<i>furcata</i> (Fabricius, 1794)	B		= <i>demandata</i> (Fabricius, 1798)		
NEOTTIOPHILINAE			<i>Ulidia</i> Meigen, 1826		
<i>Actenoptera</i> Czerny, 1904			<i>erythrophthalma</i> Meigen, 1826	B	M S
<i>hilarella</i> (Zetterstedt, 1847)	B	S			
<i>Neottiophilum</i> Frauenfeld, 1868			PLATYSTOMATIDAE		
<i>praeustum</i> (Meigen, 1826)	B	M S	Vladislav Martinek		
OTITIDAE			<i>Platystoma</i> Meigen, 1803		
Vladislav Martinek			<i>gemmationis</i> (Rondani, 1869)		M S
<i>Cephalia</i> Meigen, 1826			<i>lugubre</i> (Robineau-Desvoidy, 1830)	B	M S
<i>rufipes</i> Meigen, 1826		M S	<i>plantationis</i> (Rondani, 1869)		S
<i>Ceroxys</i> Macquart, 1835			<i>pubescens</i> Loew, 1845		M S
<i>hortulana</i> (Rossi, 1790)		M S	<i>seminationis seminationis</i> (Fabricius, 1775)	B	M S
<i>munda</i> (Loew, 1868)		M S	<i>seminationis biseta</i> Loew, 1868		S
<i>urticae</i> (Linnaeus, 1758)	B	M S	<i>Rivellia</i> Robineau-Desvoidy, 1830		
<i>Dorycera</i> Meigen, 1830			<i>syngenesiae</i> (Fabricius, 1781)	B	M S
<i>graminum</i> (Fabricius, 1794)		S			
<i>Herina</i> Robineau-Desvoidy, 1830			TEPHRITIDAE		
<i>frondescentiae</i> (Linnaeus, 1758)	B	M S	Judita Kinkorová		
<i>germinations</i> (Rossi, 1790)		M S	MYOPITINAE		
<i>lugubris</i> (Meigen, 1826)	B	M S	<i>Myopites</i> Blot, 1827		
<i>oscillans</i> (Meigen, 1826)		S	<i>apicatus</i> Freidberg, 1979	B	S
<i>paludum</i> (Fallén, 1820)		M S	<i>inulaedyssentericae</i> Blot, 1827		M S
<i>palustris</i> (Meigen, 1826)	B	M S	<i>olii</i> Dirlbek, 1974		M
<i>parva</i> (Loew, 1864)		S	<i>tenellus</i> Frauenfeld, 1863	B	M
<i>scutellaris</i> Robineau-Desvoidy, 1830	B	M S	<i>Urophora</i> Robineau-Desvoidy, 1830		
<i>Melieria</i> Robineau-Desvoidy, 1830			<i>affinis</i> (Frauenfeld, 1857)	B	M S
<i>acuticornis</i> (Loew, 1854)		M S	<i>aprica</i> (Fallén, 1820)	B	M S
<i>crassipennis</i> (Fabricius, 1794)	B	M S	<i>cardui</i> (Linnaeus, 1758)	B	M S
<i>omissa</i> (Meigen, 1826)	B	M S	<i>congrua</i> Loew, 1862	B	M S
<i>pieta</i> (Meigen, 1826)	B	M S	<i>cuspidata</i> (Meigen, 1824)	B	M
<i>Myennis</i> Robineau-Desvoidy, 1830			<i>jaceana</i> Hering, 1935	B	M S
<i>octopunctata</i> (Coquebert, 1798)	B	M S	<i>macrura</i> (Loew, 1855)	B	
<i>Orites</i> Latreille, 1804			<i>maura</i> (Frauenfeld, 1857)		M S
<i>bacescui</i> Gheorghiu, 1987	B	M S	<i>mauritanica</i> Macquart, 1851	B	
<i>ruficeps</i> auct. nec Fabricius, 1805			<i>quadrifasciata</i> (Meigen, 1826)	B	M S
<i>centralis</i> (Fabricius, 1805)	B	M S	<i>solstitialis</i> (Linnaeus, 1758)	B	M S
<i>formosa</i> (Panzer, 1798)	B	M S	<i>spoliata</i> (Haliday, 1938)		S
<i>guttata</i> (Meigen, 1830)	B	M S	<i>stigma</i> (Loew, 1840)	B	M S
<i>lamed</i> (Schränk, 1781)	B	M S	<i>stylata</i> (Fabricius, 1775)	B	M S
<i>levigata</i> (Loew, 1873)		S	<i>terebrans</i> (Loew 1850)	B	M S
<i>nebulosa</i> (Latreille, 1811)	B	M S			
<i>Seioptera</i> Kirby, 1817					
<i>vibrans</i> (Linnaeus, 1758)	B	M S			

TEPHRITINAE

DITHRYCINI

- Dithryca* Rondani, 1856
guttularis (Meigen, 1826) B M S
Noeeta Robineau-Desvoidy, 1830
crepidis Hering, 1936 B M
pupillata (Fallén, 1814) B M S

TEPHRITINI

- Acanthiophilus* Becker, 1908
helianthi (Rossi, 1790) B M S
Acinia Robineau-Desvoidy, 1830
biflexa (Loew, 1844) M S
corniculata (Zetterstedt, 1819) B M S
Actinoptera Rondani, 1871
discoidea (Fallén, 1814) M
Campiglossa Rondani, 1870
absinthii (Fabricius, 1805) B M S
achyrophori (Loew, 1869) B M S
difficilis (Hendel, 1927) M S
doronici (Loew, 1856) B M S
grandinata (Rondani, 1870) M
guttella (Rondani, 1870) M S
irrorata (Fallén, 1814) M S
loewiana (Hendel, 1927) B M S
producta (Loew, 1844) B M S
punctella (Fallén, 1814) B S
solidaginis White, 1986 S
Dioxyna Frey, 1945
bidentis (Robineau-Desvoidy, 1830) B M S
Ensina Robineau-Desvoidy, 1830
sonchi (Linnaeus, 1767) B M S
Heringina Aczél, 1940
guttata (Fallén, 1814) B
Ictericoïdes Hering, 1942
japonicus (Wiedemann, 1830) M S
zelleri (Loew, 1844) B
Oxya Robineau-Desvoidy, 1830
flavipennis (Loew, 1844) B M S
nebulosa (Wiedemann, 1817) B M S
parietina (Linnaeus, 1758) B M S
Sphenella Robineau-Desvoidy, 1830
marginata (Fallén, 1814) B M S
Euaesta Loew, 1873
bullans (Wiedemann, 1830) S
Tephritis Latreille, 1804
acanthiophilopsis Hering, 1938 M S
angustipennis (Loew, 1844) B M S
arnicae (Linnaeus, 1758) B M S
bardanae (Schrank, 1803) B M S
cometa (Loew, 1840) B M S
conura (Loew, 1844) B M S
conyzifoliae Merz, 1992 B
crepidis Hendel, 1927 B M S
dilacerata (Loew, 1846) B M S
dioscurea (Loew, 1856) B M S
fallax (Loew, 1844) B M S

- formosa* (Loew, 1844) B M S
hyoscyami (Linnaeus, 1758) B M S
leontodontis (DeGeer, 1776) B M S
matricariae (Loew, 1844) B S
mutabilis Merz, 1992 B
neesii (Meigen, 1830) B M S
nigricauda (Loew, 1856) B M
postica (Loew, 1844) M S
pulchra (Loew, 1844) S
ruralis (Loew, 1844) B M S
separata Rondani, 1871 B
tanacetii Hering, 1956 B
truncata (Loew, 1844) B M S
vespertina (Loew, 1844) B M S
zernyi Hendel, 1927 S
Trupanea Schrank, 1795
amoena (Frauenfeld, 1857) M S
stellata (Fuessly, 1775) B M
Xyphosia Robineau-Desvoidy, 1830
miliaria (Schrank, 1781) B M S

TERELLIINAE

- Chaetorellia* Hendel, 1927
acrolophi White (Macquart, 1989) M
jaceae (Robineau-Desvoidy, 1830) B M S
loricata (Rondani, 1870) B M S
Chaetostomella Hendel, 1927
cylindrica (Robineau-Desvoidy, 1830) B M S
Orellia Robineau-Desvoidy, 1830
distans (Loew, 1847) B M
falcata (Scopoli, 1763) B M S
punctata (Schrank, 1781) M
Terellia Robineau-Desvoidy, 1830
sg. *Cerajocera* Rondani, 1856
ceratocera (Hendel 1913) B M S
lappae (Cederhjelm, 1798) B M
tussilaginis (Fabricius, 1775) B M S
sg. *Terellia* s. str.
colon (Meigen, 1826) B M S
longicauda (Meigen, 1838) B M S
ruficauda (Fabricius, 1794) B M S
serratulae (Linnaeus, 1758) B M S
virens (Loew, 1846) B M S
winthemi (Meigen, 1826) B M S

TRYPETINAE

EUPHRANTINI

- Euphranta* Loew, 1862
sg. *Euphranta* s. str.
connexa (Fabricius, 1794) B M S
sg. *Rhacochlaena* Loew, 1862
toxoneura (Loew, 1846) B M S

TRYPETINI

- Acidia* Robineau-Desvoidy, 1830
cognata (Wiedemann, 1817) B M S
Anomoia Walker, 1835
pormunda (Harris, 1776) B M S

<i>Cryptaciura</i> Hendel, 1927			<i>Lauxania</i> Latreille, 1804	
<i>rotundriventris</i> (Fallén, 1814)	B	M	sg. <i>Callixania</i> Papp, 1978	
<i>Euleia</i> Walker, 1853			<i>minor</i> Martinek, 1974	B M S
<i>heracleii</i> (Linnaeus, 1758)	B	M S	sg. <i>Lauxania</i> s.str.	
<i>Myoleja</i> Rondani, 1856			<i>cylindricornis</i> (Fabricius, 1794)	B M S
<i>lucida</i> (Fallén, 1826)	B	M S	<i>Lyciella</i> Collin, 1948	
<i>Philophylla</i> Rondani, 1870			<i>affinis</i> (Zetterstedt, 1847)	B M S
<i>caesio</i> Harris, 1776	B	M S	<i>conjugata</i> (Becker, 1895)	B M S
<i>Platyparea</i> Loew, 1862			<i>decempunctata</i> (Fallén, 1820)	B M S
<i>discoidea</i> (Fabricius, 1787)	B	M S	<i>decipiens</i> (Loew, 1847)	B M S
<i>Plioreocepta</i> Korneyev, 1987			<i>illota</i> (Loew, 1847)	B M S
<i>poeciloptera</i> (Schränk, 1776)	B		<i>laeta</i> (Zetterstedt, 1838)	B M S
<i>Rhagoletis</i> Loew, 1862			<i>mihalyii</i> Papp, 1978	B M S
<i>alternata</i> (Fallén, 1814)	B	M S	<i>pallidiventris</i> (Fallén, 1820)	B M S
<i>cerasi</i> (Linnaeus, 1758)	B	M S	<i>platycephala</i> (Loew, 1847)	B M S
<i>meigenii</i> (Loew, 1844)	B	M	<i>rorida</i> (Fallén, 1820)	B M S
<i>Stemonocera</i> Rondani, 1870			<i>stylata</i> Papp, 1978	B M S
<i>cornuta</i> (Scopoli, 1772)		S	<i>subfasciata</i> (Zetterstedt, 1838)	B M S
<i>Trypeta</i> Meigen, 1803			<i>vittata</i> (Walker, 1849)	B M S
<i>artemisiae</i> (Fabricius, 1794)	B	M S	<i>Minettia</i> Robineau-Desvoidy, 1830	
<i>immaculata</i> (Macquart, 1835)	B	M	sg. <i>Frendelia</i> Collin, 1948	
<i>zoe</i> Meigen, 1826	B	M S	<i>longipennis</i> (Fabricius, 1794)	B M S
LAUXANIIDAE			sg. <i>Minettia</i> s.str.	
Vladislav Martinek			<i>austriaca</i> Hennig, 1951	M S
HOMONEURINAE			<i>fasciata</i> (Fallén, 1826)	B M S
<i>Homoneura</i> van der Wulp, 1891			<i>filia</i> (Becker, 1895)	B M S
<i>biumbrata</i> (Loew, 1847)	B	M S	<i>flaviventris</i> (Costa, 1844)	B M S
<i>consobrina</i> (Zetterstedt, 1847)		M	<i>helvola</i> (Becker, 1895)	S
<i>interstincta</i> (Fallén, 1820)	B	M S	<i>inusta</i> (Meigen, 1826)	B M S
<i>lamellata</i> (Becker, 1895)	B		<i>loewi</i> (Schiner, 1864)	B M S
<i>limnea</i> (Becker, 1895)	B	M S	<i>lupulina</i> (Fabricius, 1787)	B M S
<i>minor</i> (Becker, 1895)		S	<i>plumicornis</i> (Fallén, 1820)	B M S
<i>notata</i> (Fallén, 1820)		M S	<i>rivosa</i> (Meigen, 1826)	B M S
<i>patelliformis</i> (Becker, 1895)	B	M S	<i>Pachycerina</i> Macquart, 1835	
<i>remmi</i> Papp, 1978	B	M S	<i>pulchra</i> (Loew, 1850)	B M
<i>subnotata</i> Papp, 1978	B	M S	<i>seticornis</i> (Fallén, 1820)	B M S
<i>tesquae</i> (Becker, 1895)	B		<i>Paroecus</i> Becker, 1895	
<i>thammeri</i> Papp, 1978		S	<i>signatipes</i> (Loew, 1856)	B M S
LAUXANIINAE			<i>Peplomyza</i> Haliday, 1837	
<i>Aulogastromyia</i> Hendel, 1925			<i>discoidea</i> (Meigen, 1830)	B M S
<i>anisodactyla</i> (Loew, 1845)	B	M S	<i>intermedia</i> E. Remm, 1979	B M S
<i>Calliopum</i> Strand, 1928			<i>litura</i> (Meigen, 1826)	B M S
<i>aeneum</i> (Fallén, 1820)	B	M S	<i>Sapromyza</i> Fallén, 1810	
<i>elisae</i> (Meigen, 1826)	B	M S	sg. <i>Sapromyza</i> s.str.	
<i>geniculatum</i> (Fabricius, 1805)	B	M	<i>albiceps</i> Fallén, 1820	B M S
<i>simillimum</i> (Collin, 1933)	B	M S	<i>apicalis</i> Loew, 1847	B M S
<i>splendidum</i> Papp, 1978	B	M S	<i>basalis</i> Zetterstedt, 1847	B M S
<i>Cnemaacantha</i> Macquart, 1835			<i>intonsa</i> Loew, 1847	B M S
<i>muscaria</i> (Fallén, 1823)	B	M S	<i>obscuripennis</i> Loew, 1847	B M S
<i>Eusapromyza</i> Malloch, 1923			<i>obsoleta</i> Fallén, 1820	B M S
<i>halioptera</i> Czerny, 1932		M S	<i>opaca</i> Becker, 1895	B M S
<i>multipunctata</i> (Fallén, 1820)	B	M S	<i>palpella</i> Rondani, 1868	M S
<i>poeciloptera</i> (Loew, 1873)		M	<i>setiventris</i> Zetterstedt, 1847	B M S
			<i>sempunctata</i> Meigen, 1826	B M S
			<i>simplex</i> Loew, 1847	M S
			= <i>simplicior</i> Hendel, 1908	
			<i>visciespunctata</i> Czerny, 1932	B

<i>zetterstedti</i> Hendel, 1908	B M S	<i>decrepita</i> Zetterstedt, 1838	B M S
sg. <i>Schumannimyia</i> Papp, 1978		<i>flaveola</i> (Fabricius, 1794)	B M S
<i>hyalinata</i> (Meigen, 1826)	B M S		
Sapromyzosoma Lioy, 1864			
<i>bipunctata</i> Meigen, 1830	B M S	PHAEOMYIIDAE	
<i>quadrimeta</i> Becker, 1895		Rudolf Rozkošný	
<i>quadripunctata</i> (Linnaeus, 1767)	B M S		
Trigonometopus Macquart, 1835		Pelidnoptera Rondani, 1856	
<i>frontalis</i> (Meigen, 1830)	B M S	<i>fuscipennis</i> (Meigen, 1830)	B M S
Tricholauxania Hendel, 1925		<i>leptiformis</i> (Schiner, 1864)	B M S
<i>praeusta</i> (Fallén, 1820)	B M S	<i>nigripennis</i> (Fabricius, 1794)	B M S
CHAMAEMYIIDAE		SCIOMYZIDAE	
Vlastimil Lapáček		Rudolf Rozkošný	
Acrometopia Schiner, 1862		SALTICELLINAE	
<i>wahlbergi</i> (Zetterstedt, 1846)	B M S	Salticella Robineau-Desvoidy, 1830	
Parochthiphila Czerny, 1904		<i>fasciata</i> (Meigen, 1830)	S
<i>coronata</i> (Loew, 1858)	M S		
Chamaemyia Meigen, 1803		SCIOMYZINAE	
<i>aestiva</i> Tanasijtshuk, 1970	B S	SCIOMYZINI	
<i>aridella</i> (Fallén, 1823)	B M S	Colobaea Zetterstedt, 1837	
<i>elegans</i> (Panzer, 1809)	B M S	<i>beckeri</i> (Hendel, 1902)	S
<i>emiliae</i> Tanasijtshuk, 1970	M	<i>bifasciella</i> (Fallén, 1820)	B M S
<i>fasciata</i> (Loew, 1858)	B M	<i>distincta</i> (Meigen, 1830)	B M S
<i>flavicornis</i> (Strobl, 1902)	S	<i>pectoralis</i> (Zetterstedt, 1847)	B M S
<i>flavipalpis</i> (Haliday, 1838)	B M S	<i>punctata</i> (Lundbeck, 1923)	M S
<i>geniculata</i> (Zetterstedt, 1838)	B M S	Ditaeniella Sack, 1939	
<i>herbarum</i> (Robineau-Desvoidy, 1830)	B M S	<i>grisescens</i> (Meigen, 1830)	B M S
<i>juncorum</i> (Fallén, 1823)	B M S	Pherbellia Robineau-Desvoidy, 1830	
<i>paludosa</i> Collin, 1966	B M S	<i>albocostata</i> (Fallén, 1820)	B M S
<i>polystigma</i> (Meigen, 1830)	B M S	<i>annulipes</i> (Zetterstedt, 1846)	B M S
Leucopis Meigen, 1830		<i>argyra</i> Verbeke, 1967	B M S
sg. <i>Leucopis</i> s. str.		<i>austera</i> (Meigen, 1830)	B M S
<i>annulipes</i> Zetterstedt, 1848	M S	<i>brunnipes</i> (Meigen, 1838)	B M S
<i>glyphinivora</i> Tanasijtshuk, 1958	S	<i>cinerella</i> (Fallén, 1820)	B M S
<i>griseola</i> (Fallén, 1823)	M S	<i>czernyi</i> (Hendel, 1902)	M S
<i>puncticornis</i> Meigen, 1830	M	<i>dorsata</i> (Zetterstedt, 1846)	B M S
<i>salicis</i> Tanasijtshuk, 1962	S	<i>hubia</i> (Fallén, 1820)	B M S
sg. <i>Leucopomyia</i> Malloch, 1921		<i>griseola</i> (Fallén, 1820)	B M S
<i>silestaca</i> Egger, 1862	B	<i>limbata</i> (Meigen, 1830)	B M S
sg. <i>Neoleucopis</i> Malloch, 1921		<i>majuscula</i> (Rondani, 1868)	S
<i>atranula</i> (Ratzeburg, 1844)	M S	<i>hungarica</i> Rozkošný, 1972	
<i>obscura</i> Haliday, 1833	S	<i>nana</i> (Fallén, 1820)	B M S
		<i>obtusa</i> (Fallén, 1820)	B M S
CREMIFANIIDAE		<i>pallidicarpa</i> (Rondani, 1868)	M S
Jan Máca		<i>lichtwardti</i> (Hendel, 1902)	
		<i>pallidiventris</i> (Fallén, 1820)	B M S
Cremifania Czerny, 1904		<i>pilosa</i> (Hendel, 1902)	M
<i>nigrocellulata</i> Czerny, 1904	S	<i>rozkošnyi</i> Verbeke, 1967	S
		<i>schoenherri</i> (Fallén, 1826)	B M S
DRYOMYZIDAE		<i>scutellaris</i> (von Roser, 1840)	B M S
Vladislav Martinek		<i>silana</i> Rivosecchi, 1989	M S
		<i>sordida</i> (Hendel, 1902)	M
Dryomyza Fallén, 1820		<i>steyskali</i> Rozkošný & Zuska, 1965	B M S
<i>analís</i> Fallén, 1820	B M S	<i>ventralis</i> (Fallén, 1820)	B M S

<i>Pteromicra</i> Lioy, 1864				<i>Renocera</i> Hendel, 1900			
<i>angustipennis</i> (Staeger, 1845)	B	M	S	<i>pallida</i> (Fallén, 1820)		B	M S
<i>glabricula</i> (Fallén, 1820)	B	M	S	<i>striata</i> (Meigen, 1830)		B	M
<i>leucopeza</i> (Meigen, 1838)	B	M	S	<i>stroblii</i> Hendel, 1900		B	M S
<i>aldenbergi</i> (Hendel, 1902)			S	<i>Sepedon</i> Latreille, 1804			
<i>pectorosa</i> (Hendel, 1902)		M	S	<i>sphegea</i> (Fabricius, 1775)		B	M S
<i>Sciomyza</i> Fallén, 1820				<i>spinipes</i> (Scopoli, 1763)		B	M S
<i>dryomyzina</i> Zetterstedt, 1846	B	M	S	<i>Tetanocera</i> Duméril, 1800			
<i>pulchra</i> Roller, 1996			S	<i>arrogans</i> Meigen, 1830		B	M S
<i>simplex</i> Fallén, 1820	B	M	S	<i>elata</i> (Fabricius, 1781)		B	M S
<i>testacea</i> Macquart, 1835	B	M	S	<i>ferruginea</i> Fallén, 1820		B	M S
<i>Tetanura</i> Fallén, 1820				<i>fuscinervis</i> (Zetterstedt, 1838)		B	M S
<i>pallidiventris</i> Fallén, 1820	B	M	S	<i>hyalipennis</i> von Roser, 1840		B	M S
TETANOCERINI				<i>montana</i> Day, 1881			M S
<i>Anticheta</i> Haliday, 1838				<i>phyllophora</i> Melander, 1920		B	M S
<i>Antichaeta</i> auct. (emend.)				<i>punctifrons</i> Rondani, 1868		B	M S
<i>analís</i> (Meigen, 1830)	B	M	S	<i>robusta</i> Loew, 1847		B	M S
<i>atriseta</i> (Loew, 1849)	B			<i>silvatica</i> Meigen, 1830		B	M S
<i>brevipennis</i> (Zetterstedt, 1846)		M	S	<i>Trypetoptera</i> Hendel, 1900			
<i>obliviosa</i> Enderlein, 1939			S	<i>punctulata</i> (Scopoli, 1763)		B	M S
<i>Coremacera</i> Rondani, 1856				SEPSIDAE			
<i>catenata</i> (Loew, 1847)		M	S	Jan Zuska & Miroslav Barták			
<i>fabricii</i> Rozkošný, 1981	B	M	S	<i>Meroplius</i> Rondani, 1874			
<i>marginata</i> (Fabricius, 1775)	B	M	S	<i>minutus</i> (Wiedemann, 1830)		B	M S
<i>Dichetophora</i> Rondani, 1868				<i>Nemopoda</i> Robineau-Desvoidy, 1830			
<i>Dichetophora</i> auct. (emend.)				<i>nitidula</i> (Fallén, 1820)		B	M S
<i>finlandica</i> Verbeke, 1964	B	M	S	<i>pectinulata</i> Loew, 1873		B	M S
<i>obliterata</i> (Fabricius, 1805)	B	M	S	<i>speiseri</i> (Duda, 1926)			M S
<i>Dictya</i> Meigen, 1803				<i>Saltella</i> Robineau-Desvoidy, 1830			
<i>umbrarum</i> (Linnaeus, 1758)	B	M	S	<i>nigripes</i> Robineau-Desvoidy, 1830		B	S
<i>Ectinocera</i> Zetterstedt, 1838				<i>sphondylii</i> (Schrank, 1803)		B	M S
<i>borealis</i> Zetterstedt, 1838			S	<i>Sepsis</i> Fallén, 1810			
<i>Elgiva</i> Meigen, 1838				<i>barbata</i> Becker, 1907			S
<i>cueularia</i> (Linnaeus, 1767)	B	M	S	<i>biflexuosa</i> Strobl, 1893		B	M S
<i>solicita</i> (Harris, 1780)	B	M	S	<i>cynipsea</i> (Linnaeus, 1758)		B	M S
<i>Euthycera</i> Latreille, 1829				<i>duplicata</i> Haliday, 1838		B	M S
<i>chaerophylli</i> (Fabricius, 1798)	B	M	S	<i>fissa</i> Becker, 1903			S
<i>morio</i> (Mayer, 1953)				<i>flavimana</i> Meigen, 1826		B	M S
<i>fumigata</i> (Scopoli, 1763)	B	M	S	<i>fulgens</i> Hoffmannsegg in Meigen, 1826		B	M S
<i>suctica</i> (Fabricius, 1805)			S	<i>neocynipsea</i> Melander & Spuler, 1917			M S
<i>Hydromya</i> Robineau-Desvoidy, 1830				<i>orthocnemis</i> Frey, 1908		B	M S
<i>dorsalis</i> (Fabricius, 1775)	B	M	S	<i>pseudomonostigma</i> Ursu, 1969		B	M
<i>Ilione</i> Haliday, 1837				<i>punctum</i> (Fabricius, 1794)		B	M S
<i>Kuntsonia</i> Verbeke, 1964				<i>thoracica</i> (Robineau-Desvoidy, 1830)		B	M S
<i>albiseta</i> (Scopoli, 1763)	B	M	S	<i>violacea</i> Meigen, 1826		B	M S
<i>lineata</i> (Fallén, 1820)	B	M		<i>Themira</i> Robineau-Desvoidy, 1830			
<i>Limnia</i> Robineau-Desvoidy, 1830				<i>annulipes</i> (Meigen, 1826)		B	M S
<i>paludicola</i> Elberg, 1965	B	M	S	<i>biloba</i> Andersson, 1975		B	
<i>unguicornis</i> (Scopoli, 1763)	B	M	S	<i>gracilis</i> (Zetterstedt, 1847)			S
<i>Pherbina</i> Robineau-Desvoidy, 1830				<i>leachi</i> (Meigen, 1826)		B	M S
<i>coryleti</i> (Scopoli, 1763)	B	M	S	<i>lucida</i> (Staeger, 1844)		B	M S
<i>intermedia</i> Verbeke, 1948	B	M	S	<i>minor</i> (Haliday, 1833)		B	M S
<i>Psacadina</i> Enderlein, 1939				<i>nigricornis</i> (Meigen, 1826)		B	M S
<i>verbekei</i> Rozkošný, 1975	B	M	S	<i>putris</i> (Linnaeus, 1758)		B	M S
<i>vittigera</i> (Schiner, 1864)	B	M	S				
<i>zernyi</i> (Mayer, 1953)	B	M	S				

<i>superba</i> (Haliday, 1833)	B M	<i>albipennis</i> Meigen, 1830	B M S
<i>Xenosepsis</i> Malloch, 1925		<i>albitarsis</i> Meigen, 1830	B M S
<i>fukuharai</i> Iwasa, 1984	B S	<i>alnihetulae</i> Hendel, 1931	M S
		<i>alnivora</i> Spencer, 1969	M S
CLUSIIDAE		<i>ambigua</i> Fallén, 1823	M S
Jindřich Roháček		<i>anderssoni</i> Spencer, 1976	S
<i>Clusia</i> Haliday, 1838		<i>anthracina</i> Meigen, 1830	B M S
<i>flava</i> (Meigen, 1830)	B M S	<i>bicaudata</i> (Hendel, 1920)	B M S
<i>Clusiodes</i> Coquillett, 1904		<i>bromi</i> Spencer, 1966	B M S
sg. <i>Clusiodes</i> s.str.		<i>cinerascens</i> Macquart, 1835	B M S
<i>albumanus</i> (Meigen, 1830)	B M S	<i>conjuncta</i> Spencer, 1966	B M S
<i>caledonicus</i> (Collin, 1912)	M S	<i>dipsaci</i> Hendel, 1927	M S
sg. <i>Clusiarina</i> Malloch, 1922		<i>erythrocephala</i> Hendel, 1920	M
<i>geomyzinus</i> (Fallén, 1823)	B M S	<i>felleri</i> Hering, 1941	M
<i>pictipes</i> (Zetterstedt, 1855)	B M S	<i>ferruginosa</i> van der Wulp, 1871	B M S
<i>ruficollis</i> (Meigen, 1830)	B M S	<i>filipendulae</i> Spencer, 1976	M
<i>facialis</i> (Collin, 1953)		<i>flaviceps</i> Fallén, 1823	B M S
sg. <i>Columbiella</i> Malloch, 1922		<i>flavipennis</i> Hendel, 1920	B M S
<i>verticalis</i> (Collin, 1912)	B M S	<i>frontella</i> (Rondani, 1875)	B M S
<i>Hendelia</i> Czerny, 1903		<i>hendeli</i> Griffiths, 1963	M S
<i>heckeri</i> Czerny, 1903	B M S	<i>igniceps</i> Hendel, 1920	M S
<i>Heteromeringia</i> Czerny, 1903		<i>intermittens</i> (Becker, 1907)	B M S
<i>nigrimana</i> (Loew, 1864)	M	<i>johannae</i> de Meijere, 1924	M
<i>Paraclusia</i> Czerny, 1903		<i>lathyri</i> Hendel, 1923	M
<i>tigrina</i> (Fallén, 1820)	B M S	<i>lucida</i> Hendel, 1920	B M S
		<i>luteitarsis</i> (Rondani, 1875)	B M S
ACARTOPHTHALMIDAE		<i>megalopsis</i> Hering, 1933	B M S
Jindřich Roháček		<i>mobilis</i> Meigen, 1830	B M S
<i>Acartophthalmus</i> Czerny, 1902		<i>myosotidis</i> Kaltenbach, 1864	B
<i>bicolor</i> Oldenberg, 1910	B M S	<i>nana</i> Meigen, 1830	B M S
<i>nigrinus</i> (Zetterstedt, 1847)	B M S	<i>nigrella</i> (Rondani, 1875)	B M S
		<i>nigrescens</i> Hendel, 1920	B M
ODINIIDAE		<i>nigripes</i> Meigen, 1830	B M S
Jan Máca		<i>nigrociliata</i> Hendel, 1931	B M S
<i>Neoalticomerus</i> Hendel, 1903		<i>orobi</i> Hendel, 1920	M
<i>formosus</i> (Loew, 1844)	B S	<i>phragmitidis</i> Hendel, 1922	B M S
<i>Odinia</i> Robineau-Desvoidy, 1830		<i>potentillae</i> (Kaltenbach, 1864)	B M S
<i>boletina</i> (Zetterstedt, 1848)	B M S	<i>prespana</i> Spencer, 1957	M S
<i>czernyi</i> Collin, 1952	M S	<i>pseudoreptans</i> Nowakowski, 1967	B M S
<i>loewi</i> Collin, 1952	B	<i>pseudorufipes</i> Nowakowski, 1964	B M S
<i>maculata</i> (Meigen, 1830)	B M S	<i>pulla</i> Meigen, 1830	M S
<i>mejerei</i> Collin, 1952	B M S	<i>reptans</i> Fallén, 1823	B M S
<i>xanthocera</i> Collin, 1952	B	<i>rondensis</i> Strobl, 1900	B M S
		<i>rufipes</i> Meigen, 1830	B
AGROMYZIDAE		<i>spenceri</i> Griffiths, 1963	M
Miloslav Vála & Miloš Černý		<i>spiraeoidearum</i> Hering, 1954	M S
AGROMYZINAE		<i>sulfuriceps</i> Strobl, 1898	B M S
<i>Agromyza</i> Fallén, 1810		<i>vicifoliae</i> Hering, 1932	B M S
<i>abiens</i> Zetterstedt, 1848	B M S	<i>woerzi</i> Groschke, 1957	M S
<i>alandensis</i> Spencer, 1976	M	<i>Hexomyza</i> Enderlein, 1936	
		<i>sarothamni</i> (Hendel, 1923)	M
		<i>Melanagromyza</i> Hendel, 1920	
		<i>aenea</i> (Meigen, 1830)	B M
		<i>aeneoventris</i> (Fallén, 1823)	B M S
		<i>albocilia</i> Hendel, 1931	M
		<i>angeliciphaga</i> Spencer, 1969	M S
		<i>artemisiae</i> Spencer, 1957	M

sg. <i>Icteromyza</i> Hendel, 1931			<i>globulariae</i> Hendel, 1931	M
<i>capitata</i> (Zetterstedt, 1848)	B	M	<i>hampsteadensis</i> Spencer, 1971	B
<i>churchillensis</i> Spencer, 1969		M	<i>huidobrensis</i> (Blanchard, 1926)	B
<i>geniculata</i> (Fallén, 1823)	B	M	<i>infusata</i> Hering, 1926	B
<i>lineella</i> (Zetterstedt, 1840)		M	<i>lutea</i> (Meigen, 1830)	B
sg. <i>Phytomyza</i> Hendel, 1920			<i>occipitalis</i> Hendel, 1931	S
<i>flavocingulata</i> (Strobl, 1909)	B	M	<i>orbona</i> (Meigen, 1830)	B
sg. <i>Poemyza</i> Hendel, 1931			<i>phryne</i> Hendel, 1931	B
<i>atra</i> (Meigen, 1830)	B	M	<i>pisivora</i> Hering, 1954	M
<i>calamagrostidis</i> Nowakowski, 1967	B	M	<i>ptarmicae</i> de Meijere, 1925	B
<i>imbuta</i> (Meigen, 1838)		M	<i>puella</i> (Meigen, 1830)	S
<i>deschampsiae</i> (Spencer, 1957)			<i>pusilla</i> (Meigen, 1830)	M
<i>incisa</i> (Meigen, 1830)	B	M	<i>pusio</i> (Meigen, 1830)	B
<i>inconspicua</i> (Malloch, 1913)		M	<i>richteri</i> Hering, 1927	B
<i>lapplandica</i> (Rydén, 1956)	B	M	<i>scorzonerae</i> Rydén, 1951	B
<i>lateralis</i> (Macquart, 1835)	B	M	<i>sonchi</i> Hendel, 1931	S
<i>melicae</i> Nowakowski, 1973	B	M	<i>strigata</i> (Meigen, 1830)	B
<i>muscina</i> (Meigen, 1830)	B	M	<i>taraxaci</i> Hering, 1927	B
<i>phalaridis</i> Nowakowski, 1967	B		<i>trifolii</i> (Burgess, 1880)	B
<i>phragmitidis</i> Nowakowski, 1967	B	M	<i>virgo</i> (Zetterstedt, 1838)	M
<i>pygmaea</i> (Meigen, 1830)	B	M	<i>xanthocera</i> (Czerny & Strobl, 1909)	B
<i>superciliosa</i> (Zetterstedt, 1860)	B	M	Metopomyza Enderlein, 1936	
<i>thunbergi</i> Nowakowski, 1967		M	<i>flavonotata</i> (Haliday, 1833)	B
<i>zuskai</i> Nowakowski, 1973		S	<i>interfrontalis</i> (Melander, 1913)	B
<i>zoeneri</i> Nowakowski, 1973	B		= <i>xanthaspida</i> (Hendel, 1920)	
sg. <i>Xenophytomyza</i> Frey, 1946			<i>nigriorbita</i> (Hendel, 1931)	S
<i>atronitens</i> (Hendel, 1920)	B	M	<i>nigrohumeralis</i> (Hendel, 1931)	S
<i>biseta</i> (Hendel, 1920)	B	M	<i>ornata</i> (Meigen, 1830)	S
<i>venturii</i> Nowakowski, 1967	B	M	<i>scutellata</i> (Fallén, 1823)	B
Chromatomyia Hardy, 1849			<i>xanthaspioides</i> (Frey, 1946)	S
<i>fuscula</i> (Zetterstedt, 1838)	B	M	<i>xanthaspis</i> (Loew, 1858)	M
<i>gentianae</i> (Hendel, 1920)	B	M	Napomyza Westwood, 1840	
<i>horticola</i> (Goureau, 1851)	B	M	<i>albipennis</i> (Fallén, 1823)	M
= <i>atricornis</i> auct. nec Meigen, 1838			<i>elegans</i> (Meigen, 1830)	B
<i>istae</i> (Hering, 1962)		M	<i>evanescens</i> (Hendel, 1920)	B
<i>lonicerae</i> (Robineau-Desvoidy, 1851)	B		<i>glabra</i> (Hendel, 1935)	M
<i>mili</i> (Kaltenbach, 1864)	B	M	<i>gymnostoma</i> (Loew, 1858)	M
<i>nigra</i> (Meigen, 1830)	B	M	<i>hirticornis</i> Hendel, 1932	M
<i>opacella</i> (Hendel, 1935)		M	<i>lateralis</i> (Fallén, 1823)	B
<i>primulae</i> (Robineau-Desvoidy, 1851)		M	<i>nigriceps</i> van der Wulp, 1871	B
<i>pseudomili</i> Griffiths, 1980	B		<i>scrophulariae</i> Spencer, 1966	B
<i>ramosa</i> (Hendel, 1923)	B	M	Nemorimyza Frey, 1946	
Galiomyza Spencer, 1981			<i>posticata</i> (Meigen, 1830)	M
<i>morio</i> (Brischke, 1880)		M	Phytobia Lioy, 1864	
Liriomyza Mik, 1894			<i>humulata</i> (Hendel, 1920)	S
<i>amoena</i> (Meigen, 1830)		M	Phytoliriomyza Hendel, 1931	
<i>artemisticola</i> de Meijere, 1924	B	M	<i>arctica</i> (Lundbeck, 1901)	M
<i>balcanica</i> (Strobl, 1898)		M	<i>hilarella</i> (Zetterstedt, 1848)	S
<i>buhri</i> Hering, 1937		M	<i>melampyga</i> (Loew, 1869)	B
<i>centaureae</i> Hering, 1927		S	<i>perpusilla</i> (Meigen, 1830)	B
<i>cicerina</i> (Rondani, 1875)		S	<i>pteridii</i> Spencer, 1973	S
<i>congesta</i> (Becker, 1903)		M	<i>variegata</i> (Meigen, 1830)	B
<i>demeijerei</i> Hering, 1930		M	Phytomyza Fallén, 1810	
<i>eupatoriana</i> Spencer, 1954		M	<i>abdominalis</i> Zetterstedt, 1848	B
<i>eupatorii</i> (Kaltenbach, 1873)	B	M	<i>aconiti</i> Hendel, 1920	M
<i>flaveola</i> (Fallén, 1823)	B	M	<i>adjuncta</i> Hering, 1928	B
<i>flavopicta</i> Hendel, 1931		M	<i>affinis</i> Fallén, 1823	S

<i>albiceps</i> Meigen, 1830	M	<i>tanacetii</i> Hendel, 1923	S
<i>alpina</i> Groschke, 1957	B	<i>tussilaginis</i> Hendel, 1925	B
<i>angelicae</i> Kaltenbach, 1872	B	<i>veronicicola</i> Hering, 1925	M
<i>angelicastris</i> Hering, 1932	B M	<i>virgaureae</i> Hering, 1926	M
<i>aquilegiae</i> Hardy, 1849	S	<i>wahlgreni</i> Rydén, 1944	B M
<i>araciocecis</i> Hering, 1958	S	<i>Pseudonapomyza</i> Hendel, 1920	
<i>arnicae</i> Hering, 1925	B	<i>atra</i> (Meigen, 1830)	B M S
<i>arnicicola</i> Lundqvist, 1949	B	<i>errata</i> Zlobin, 1993	B M
<i>aronici</i> Nowakowski, 1962	M	<i>europaea</i> Spencer, 1973	B M
<i>artemisivora</i> Spencer, 1971	B M	<i>mohelnica</i> Černý, 1992	M
<i>aurei</i> Hering, 1931	M S	<i>moraviae</i> Černý, 1992	M
<i>brunnipes</i> Brischke, 1880	M	<i>palliditarsis</i> Černý, 1992	M
<i>buhriella</i> Spencer, 1969	B S	<i>spenceri</i> Černý, 1992	M
<i>caltophila</i> Hering, 1931	B M	<i>strobliana</i> Spencer, 1973	M
<i>chaerophylli</i> Kaltenbach, 1856	B M S	<i>Selachops</i> Wahlberg, 1844	
<i>continua</i> Hendel, 1920	B S	<i>flavocincta</i> Wahlberg, 1844	B
<i>conyzae</i> Hendel, 1920	M		
<i>crassiseti</i> Zetterstedt, 1860	B M S		
<i>elsae</i> Hendel, 1927	M		
<i>eupatorii</i> Hendel, 1927	M		
<i>facialis</i> Kaltenbach, 1872	M		
<i>fallaciosa</i> Brischke, 1880	M		
<i>farfarae</i> Hendel, 1935	M		
<i>flavicornis</i> Fallén, 1823	B M S		
<i>flavofemorata</i> Strobl, 1893	M		
<i>glechomae</i> Kaltenbach, 1862	M S		
<i>griffithsi</i> Spencer, 1963	M		
<i>krygeri</i> Hering, 1949	B		
<i>lappae</i> Goureau, 1851	M S		
<i>lithospermi</i> Nowakowski, 1959	S		
<i>lycopi</i> Nowakowski, 1959	S		
<i>marginella</i> Fallén, 1823	M		
<i>myosotica</i> Nowakowski, 1959	B M		
<i>nigrifemur</i> Hering, 1934	S		
<i>nigripennis</i> Fallén, 1823	B M S		
<i>nigritella</i> Zetterstedt, 1848	M		
<i>obscura</i> Hendel, 1920	S		
<i>pastinacae</i> Hendel, 1923	B M		
<i>pauliloewii</i> Hendel, 1920	B M		
<i>pimpinellae</i> Hendel, 1924	B M		
<i>plantaginis</i> Robineau-Desvoidy, 1851	M S		
<i>podagrariae</i> Hering, 1954	M		
<i>ptarmicae</i> Hering, 1937	S		
<i>pubicornis</i> Hendel, 1920	B M		
<i>pullula</i> Zetterstedt, 1848	B S		
<i>ranunculi</i> (Schrank, 1803)	B M S		
<i>ranunculicola</i> Hering, 1949	M S		
<i>ranunculivora</i> Hering, 1932	M S		
<i>rhabdophora</i> Griffiths, 1964	S		
<i>robustella</i> Hendel, 1936	M		
<i>rostrata</i> Hering, 1934	M		
<i>rufipes</i> Meigen, 1830	B M S		
<i>salviae</i> (Hering, 1924)	M		
<i>senecionis</i> Kaltenbach, 1869	B S		
<i>spinaciae</i> Hendel, 1935	M		
<i>spoliata</i> Strobl, 1906	B		
<i>spondylii</i> Robineau-Desvoidy, 1851	M		
		OPOMYZIDAE	
		Vladislav Martinek	
		<i>Geomyza</i> Fallén, 1810	
		<i>acutipennis</i> Czerny, 1928	S
		<i>angustipennis</i> Zetterstedt, 1847	M
		<i>annae</i> Martinek, 1978	B
		<i>apicalis</i> (Meigen, 1830)	B M S
		<i>balachowskyi</i> Mesnil, 1934	B M S
		<i>breviseti</i> Czerny, 1928	B M S
		<i>hackmani</i> Nartshuk, 1984	B
		<i>hendeli</i> Czerny, 1928	B S
		<i>majuscula</i> (Loew, 1864)	B M
		<i>martineki</i> Drake, 1992	B M S
		= <i>combinata</i> auct. nec Linnaeus, 1767	
		<i>paganettii</i> (Strobl, 1909)	B M S
		<i>pilosula</i> Czerny, 1928	M S
		<i>subnigra</i> Drake, 1992	B
		<i>tripunctata</i> Fallén, 1823	B M S
		<i>venusta</i> (Meigen, 1830)	B M S
		<i>virgata</i> Czerny, 1928	B
		<i>Opomyza</i> Fallén, 1820	
		<i>florum</i> (Fabricius, 1794)	B M S
		<i>germinationis</i> (Linnaeus, 1758)	B M S
		<i>lineatopunctata</i> von Roser, 1840	B M
		<i>petrei</i> Mesnil, 1934	B M S
		<i>punctata</i> Haliday, 1833	B M S
		<i>punctella</i> Fallén, 1820	B M S
		<i>thalhammeri</i> Strobl, 1900	B S
		ANTHOMYZIDAE	
		Jindřich Roháček	
		<i>Anagnota</i> Becker, 1902	
		<i>bicolor</i> (Meigen, 1838)	B M S
		<i>major</i> Roháček & Freidberg, 1993	M S
		<i>Anthomyza</i> Fallén, 1810	
		<i>albimana</i> (Meigen, 1830)	B M S

<i>anderssoni</i> Roháček, 1984	M S	ASTEIINAE	
<i>collini</i> Andersson, 1976	B M S	<i>Asteia</i> Meigen, 1830	
<i>dissors</i> Collin, 1944	B M S	<i>amoena</i> Meigen, 1830	B M S
<i>gractlis</i> Fallén, 1823	B M S	<i>concinna</i> Meigen, 1830	B M S
<i>macra</i> Czerny, 1928	B M S	<i>elegantula</i> Zetterstedt, 1847	B M S
<i>neglecta</i> Collin, 1944	B M S		
<i>pallida</i> (Zetterstedt, 1838)	B M S		
<i>paraneglecta</i> Elberg, 1968	B M S	BRAULIDAE	
<i>pleuralis</i> Czerny, 1928	B M S	Jindřich Roháček	
<i>socculata</i> (Zetterstedt, 1847)	B		
– <i>ungulata</i> Loew, 1873			
Paranthomyza Czerny, 1902		<i>Braula</i> Nitzsch, 1818	
<i>nitida</i> (Meigen, 1838)	B M S	<i>coeca coeca</i> Nitzsch, 1818	B M S
Stiphrosoma Czerny, 1928			
– <i>Ptenotaenia</i> Enderlein, 1936			
<i>cingulatum</i> (Haliday, 1855)	B M S		
<i>laetum</i> (Meigen, 1830)	B M S	CARNIDAE	
<i>sabulosum</i> (Haliday, 1837)	B M S	Jindřich Roháček	
– <i>oldenbergi</i> Czerny, 1928			
Typhamyza Roháček, 1992		<i>Carnus</i> Nitzsch, 1818	
<i>bifasciata</i> (Wood, 1911)	M S	<i>hemapterus</i> Nitzsch, 1818	B M S
		Hemeromyia Coquillett, 1902	
		<i>anthracina</i> Collin, 1949	B
		Meoneura Rondani, 1856	
		<i>acuticerva</i> Gregor, 1973	M
		<i>alpina</i> Hennig, 1948	M S
		<i>atoma</i> Papp, 1981	B
		<i>carpathica</i> Papp, 1977	B M S
		<i>exigua</i> Collin, 1930	B S
		<i>flavifacies</i> Collin, 1930	B M S
		<i>freta</i> Collin, 1937	M S
		<i>hungarica</i> Papp, 1977	B M S
		<i>lacteipennis</i> (Fallén, 1923)	B M S
		<i>milleri</i> Gregor, 1973	M S
		<i>minutissima</i> (Zetterstedt, 1860)	M S
		<i>moravica</i> Gregor & Papp, 1981	M
		<i>neglecta</i> Collin, 1930	B M S
		<i>neottiophila</i> Collin, 1930	B M S
		<i>obscura</i> (Fallén, 1923)	M
		<i>prima</i> (Becker, 1903)	B M S
		<i>triangularis</i> Collin, 1930	B M
		<i>vagans</i> (Fallén, 1823)	B M S
AULACIGASTRIDAE			
Jan Máca			
<i>Aulacigaster</i> Macquart, 1835			
<i>leucopiza</i> (Meigen, 1830)	B M S		
PERISCCELIDIDAE			
Jan Máca & Jindřich Roháček			
<i>Periscelis</i> Loew, 1858			
sg. <i>Myodris</i> Lioy, 1864			
<i>annulata</i> (Fallén, 1813)	B M S		
sg. <i>Periscelis</i> s.str.			
<i>annulipes</i> Loew, 1858	B S		
STENOMICRIDAE			
Jindřich Roháček			
<i>Stenomicro</i> Coquillett, 1900			
<i>delicata</i> (Collin, 1944)	B M S	TETHINIDAE	
		Jindřich Roháček	
ASTEIIDAE		<i>Pelomyia</i> Williston, 1893	
Jindřich Roháček		<i>steyskali</i> Hardy & Delfinado, 1980	B M S
		– <i>coronata</i> auct. nec Loew, 1866	
SIGALOESSINAE		<i>Pelomyiella</i> Hendel, 1934	
<i>Leiomyza</i> Macquart, 1835		<i>hungarica</i> (Czerny, 1928)	S
<i>curvinervis</i> (Zetterstedt, 1838)	B M S	<i>mallochi</i> (Sturtenvant, 1923)	M S
<i>duda</i> Sabrosky, 1956	B M S	Tethina Haliday, 1838	
<i>laevigata</i> (Meigen, 1830)	B M S	sg. <i>Rhinoessa</i> Loew, 1862	
<i>scatophagina</i> (Fallén, 1823)	B M S	<i>simplex</i> (Collin, 1966)	B S

MILICHIIDAE		
Jindřich Roháček		
MILICHIINAE		
<i>Milichia</i> Meigen, 1830		
<i>ludens</i> (Wahlberg, 1847)	B M S	
<i>speciosa</i> Meigen, 1830	M S	
<i>Milichiella</i> Giglio-Tos, 1895		
<i>argyrogastrer</i> (Perris, 1876)	M S	
MADIZINAE		
<i>Desmometopa</i> Loew, 1866		
<i>m-nigrum</i> (Zetterstedt, 1848)	B M S	
<i>sordida</i> (Fallén, 1820)	B M S	
<i>Leptometopa</i> Becker, 1903		
<i>latipes</i> (Meigen, 1830)	B M S	
<i>niveipennis</i> (Strobl, 1900)	B S	
<i>Madiza</i> Fallén, 1820		
<i>eximia</i> Papp, 1993		S
<i>glabra</i> Fallén, 1820	B M S	
<i>pachymera</i> Becker, 1908	B M S	
<i>Neophyllomyza</i> Melander, 1913		
<i>acyglossa</i> (Villeneuve, 1920)	B M S	
<i>leanderi</i> Hendel, 1924	B M S	
<i>Phyllomyza</i> Fallén, 1810		
<i>beckeri</i> Kramer, 1920	B	
<i>domisthorpei</i> Schmitz, 1923	B M S	
<i>equitans</i> (Hendel, 1919)	B M S	
<i>flavitaris</i> (Meigen, 1830)	B M S	
<i>formicae</i> Schmitz, 1923	B M S	
<i>longipalpis</i> (Schmitz, 1924)	B M S	
<i>melania</i> (Hendel, 1919)	B M S	
<i>rubricornis</i> Schmitz, 1923	B M	
<i>securicornis</i> Fallén, 1823	B M S	
<i>tetragona</i> Hendel, 1924		S
CHLOROPIDAE		
Jindřich Roháček		
SIPHONELLOPSINAE		
<i>Apotropina</i> Hendel, 1907		
= <i>Lastiopleura</i> Becker, 1910		
<i>brevivenosa</i> (Dely-Draskovits, 1977)		S
<i>longepilosa</i> (Strobl, 1893)		S
OSCINELLINAE		
<i>Aphanotrigonum</i> Duda, 1932		
<i>beschovski</i> Dely-Draskovits, 1981	B M S	
<i>brachypterum</i> (Zetterstedt, 1848)		S
<i>fasciellum</i> (Zetterstedt, 1855)	B S	
<i>femorellum</i> Collin, 1946		M S
<i>nigripes</i> (Zetterstedt, 1848)	B M S	
<i>trilineanum</i> (Meigen, 1830)	B M S	
<i>Calamoncosis</i> Enderlein, 1911		
sg. <i>Calamoncosis</i> s.str.		
<i>aprica</i> (Meigen, 1830)	M S	
<i>aspistylina</i> Duda, 1935		B M S
<i>duinensis</i> (Strobl, 1909)		M S
<i>minima</i> (Strobl, 1893)		M S
sg. <i>Rhaphiopyga</i> Nartshuk, 1971		
<i>glyceriae</i> Nartshuk, 1958		B M S
<i>laminiformis</i> (Becker, 1908)		B M S
= <i>stipae</i> Nartshuk, 1962		
<i>Conioscinella</i> Duda, 1929		
<i>frontella</i> (Fallén, 1820)		B M S
<i>gallarum</i> (Duda, 1933)		B M S
<i>livida</i> Nartshuk, 1970		M S
<i>mimula</i> Collin, 1946		S
<i>obscuripila</i> Duda, 1933		S
<i>sordidella</i> (Zetterstedt, 1848)		B M
<i>zetterstedti</i> Andersson, 1966		M S
<i>Crassivenula</i> Sabrosky, 1941		
<i>brachyptera</i> (Thalhammer, 1913)		B
<i>Dasyopa</i> Malloch, 1918		
= <i>Trachysiphonella</i> auct. nec Enderlein, 1936		
<i>pygmaea</i> (Meigen, 1838)		S
<i>ruficeps</i> (Macquart, 1835)		M S
<i>scutellata</i> (von Roser, 1840)		B M S
<i>Dicraeus</i> Loew, 1873		
sg. <i>Dicraeus</i> s.str.		
<i>ingratus</i> (Loew, 1866)		B M S
<i>nigropilosus</i> Becker, 1910		S
<i>raptus</i> (Haliday, 1838)		M S
<i>tibialis</i> (Macquart, 1835)		S
sg. <i>Oedesiella</i> Becker, 1910		
<i>fennicus</i> Duda, 1933		B M S
<i>rossicus</i> Stackelberg, 1955		B M S
sg. <i>Paroedesiella</i> Sabrosky, 1941		
<i>napaeus</i> Collin, 1946		M
<i>vagans</i> (Meigen, 1838)		B M S
<i>vallis</i> Collin, 1946		B M
<i>Elachiptera</i> Macquart, 1835		
<i>brevipennis</i> (Meigen, 1830)		B M S
<i>cornuta</i> (Fallén, 1820)		B M S
<i>diastema</i> Collin, 1946		B M S
<i>scrobiculata</i> (Strobl, 1901)		B M S
<i>sibirica</i> (Loew, 1858)		M S
<i>tuberculifera</i> (Corti, 1909)		B M S
<i>uniseta</i> Collin, 1939		B M S
<i>Eribolus</i> Becker, 1910		
<i>gracilior</i> (de Meijere, 1918)		B M S
<i>hungaricus</i> Becker, 1910		M S
<i>nanus</i> (Zetterstedt, 1838)		B M
<i>Gampsocera</i> Schiner, 1862		
<i>numerata</i> (Heeger, 1858)		B M S
<i>Gaurax</i> Loew, 1863		
<i>britannicus</i> Deeming, 1980		M
<i>dubius</i> (Macquart, 1835)		M
<i>flavoscutellatus</i> (Stackelberg, 1955)		B
<i>maculipennis</i> (Zetterstedt, 1848)		B M S
<i>niger</i> Czerny, 1906		M
<i>Hapleginella</i> Duda, 1933		
<i>laevifrons</i> (Loew, 1858)		B M S

<i>Incertella</i> Sabrosky, 1980			<i>femorale</i> (Collin, 1946)	M S
<i>albipalpis</i> (Meigen, 1830)	B M S		<i>platythorax</i> (Nartshuk, 1958)	B M S
<i>antennata</i> (Collin, 1946)	B M		<i>Siphonella</i> Macquart, 1835	
<i>kerteszi</i> (Becker, 1910)	B M S		<i>oscinina</i> (Fallén, 1820)	B M S
<i>scotica</i> (Collin, 1946)	M S		<i>Siphunculina</i> Rondani, 1856	
<i>zuercheri</i> (Duda, 1933)	M S		<i>aenea</i> (Macquart, 1835)	B M S
<i>Lasiambia</i> Sabrosky, 1941			<i>quinquangula</i> (Loew, 1873)	M
<i>Fiebrigella</i> auct. nec Duda, 1921			<i>simulata</i> Kanmiya, 1982	S
<i>coxalis</i> (von Roser, 1840)	M S		<i>Speccafrons</i> Sabrosky, 1980	
<i>euphila</i> (Hennig, 1941)			<i>halophila</i> (Duda, 1933)	B M S
<i>palposa</i> (Fallén, 1820)	B S		<i>Tricimba</i> Lioy, 1864	
<i>Lipara</i> Meigen, 1830			sg. <i>Tricimba</i> s.str.	
<i>lucens</i> Meigen, 1830	B M S		<i>lineella</i> (Fallén, 1820)	B M S
<i>pulltarsis</i> Doskočil & Chvála, 1971	B M S		sg. <i>Nartshukiella</i> Beschovski, 1981	
<i>rufitarsis</i> Loew, 1858	B M		<i>albisetia</i> Dely-Draskovits, 1983	B M
<i>similis</i> Schiner, 1854	B M S		<i>apicalis</i> (von Roser, 1840)	M S
<i>Melanochaeta</i> Bezzi, 1906			<i>fungicola</i> Dely-Draskovits, 1983	M S
<i>capreola</i> (Haliday, 1838)	M S		<i>humeralis</i> (Loew, 1858)	B M S
<i>pubescens</i> (Thalhammer, 1898)	B M S		= <i>aristolochiae</i> (Rondani, 1869)	
<i>Microcercis</i> Beschovski, 1978			<i>sulcella</i> (Zetterstedt, 1848)	B M S
<i>trigonella</i> (Duda, 1933)	B M S			
<i>Oscinella</i> Becker, 1909			CHLOROPINAE	
<i>alopeuri</i> Mesnil in Balachowsky			<i>Camarota</i> Meigen, 1830	
& Mesnil, 1935	S		<i>curvipennis</i> (Latreille, 1805)	B M S
<i>angularis</i> Collin, 1946	S		<i>Centorisoma</i> Becker, 1910	
<i>angustipennis</i> Duda, 1933	M S		<i>elegantulum</i> Becker, 1910	B
<i>festucae</i> Mesnil in Balachowsky			<i>Cetema</i> Hendel, 1907	
& Mesnil, 1935	S		<i>cereris</i> (Fallén, 1820)	B M S
<i>frut</i> (Linnaeus, 1758)	B M S		<i>elongata</i> (Meigen, 1830)	B M S
<i>hortensis</i> Collin, 1946	B S		<i>neglecta</i> Tonnoir, 1921	B M S
<i>maura</i> (Fallén, 1820)	B M S		<i>myopina</i> (Loew, 1866)	B M S
<i>nigerrima</i> (Macquart, 1835)	M S		<i>Chlorops</i> Meigen, 1803	
<i>nitidissima</i> (Meigen, 1838)	B M S		sg. <i>Chlorops</i> s.str.	
<i>pusilla</i> (Meigen, 1830)	B M S		<i>adjunctus</i> Becker, 1910	S
<i>trochanterata</i> Collin, 1946	B S		<i>anthracophagoideus</i> Strobl, 1901	B S
<i>vastator</i> (Curtis, 1845)	S		<i>bohemicus</i> Zuska, 1960	B M
<i>ventricosi</i> Nartshuk, 1955	S		<i>calceatus</i> Meigen, 1830	B M S
<i>Oscinimorpha</i> Lioy, 1864			<i>centromaculatus</i> Duda, 1933	B S
<i>albisetosa</i> (Duda, 1932)	M S		<i>citrinellus</i> (Zetterstedt, 1848)	S
<i>arcuata</i> (Duda, 1932)	B M S		<i>dasycerus</i> Loew, 1866	M
<i>koeleriae</i> Nartshuk, 1970	S		<i>fasciatus</i> Meigen, 1830	B M
<i>minutissima</i> (Strobl, 1900)	B M S		<i>finitimus</i> Becker, 1910	S
<i>sordidissima</i> (Strobl, 1893)	M		<i>frontosus</i> Meigen, 1830	M S
<i>Oscinisoma</i> Lioy, 1864			<i>geminatus</i> Meigen, 1830	B M S
<i>cognatum</i> (Meigen, 1830)	B M S		<i>gracilis</i> Meigen, 1830	B M S
<i>Polyodaspis</i> Duda, 1933			<i>hypostigma</i> Meigen, 1830	B M S
<i>ruficornis</i> (Macquart, 1835)	S		<i>interruptus</i> Meigen, 1830	B S
<i>sulcicollis</i> (Meigen, 1838)	B M S		<i>laetus</i> Meigen, 1830	M
<i>Pseudogaurax</i> Malloch, 1915			<i>novakii</i> Strobl, 1902	S
<i>venustus</i> (Czerny, 1906)	B		<i>pannonicus</i> Strobl, 1893	M S
<i>Rhodesiella</i> Adams, 1905			<i>planifrons</i> Loew, 1866	B M S
<i>plumigera</i> (Meigen, 1830)	M S		<i>pumilionis</i> (Bjerkander, 1778)	B M S
<i>Rhopalopterum</i> Duda, 1929			<i>ringens</i> Loew, 1866	B M S
<i>anthracinum</i> (Meigen, 1830)	M S		<i>rufescens</i> Oldenberg, 1923	B M S
<i>atricillum</i> (Zetterstedt, 1838)	B S		= <i>meigenii</i> Loew, 1866 nec Fallén, 1823	
<i>atricorne</i> (Zetterstedt, 1838)	B		<i>rufinus</i> (Zetterstedt, 1848)	S
<i>fasciolum</i> (Meigen, 1830)	B M S		<i>scalaris</i> Meigen, 1830	B M S

<i>serenus</i> Loew, 1866	B M S	<i>glabra</i> (Duda, 1933)	M S
<i>speciosus</i> Meigen, 1830	B M S	<i>tarsata</i> (Fallén, 1820)	B M S
<i>strigulus</i> (Fabricius, 1794)	B M S	<i>Parectecephala</i> Becker, 1910	
<i>trogodytes</i> (Zetterstedt, 1848)	B M	<i>longicornis</i> (Fallén, 1820)	B M
sg. <i>Sclerophallus</i> Beschovski, 1978		<i>Platycephala</i> Fallén, 1820	
<i>brevimanus</i> Loew, 1866	B S	<i>planifrons</i> (Fabricius, 1798)	B M S
<i>pallidiventris</i> Duda, 1933	S	<i>umbraculata</i> (Fabricius, 1794)	B M S
<i>Cryptonevra</i> Lioy, 1864		<i>Pseudopachychaeta</i> Strobl, 1902	
<i>consimilis</i> (Collin, 1932)	M S	<i>approximatonervis</i> (Zetterstedt, 1848)	B S
<i>diadema</i> (Meigen, 1830)	M S	<i>heleocharis</i> (Nartshuk, 1964)	M
<i>flavitaris</i> (Meigen, 1830)	B M S	<i>ruficeps</i> (Zetterstedt, 1838)	B M
<i>migrataris</i> (Duda, 1933)	M	<i>Thaumatomyia</i> Zenker, 1833	
<i>Diplotoxa</i> Loew, 1863		<i>elongatula</i> (Becker, 1910)	B S
sg. <i>Diplotoxa</i> s.str.		<i>glabra</i> (Meigen, 1830)	B M S
<i>messoria</i> (Fallén, 1820)	B M S	<i>hallandica</i> Andersson, 1966	M S
sg. <i>Diplotoxoides</i> Andersson, 1977		<i>notata</i> (Meigen, 1830)	B M S
<i>dalmatina</i> (Strobl, 1900)	B M S	<i>rufa</i> (Macquart, 1835)	B M S
<i>Epichlorops</i> Becker, 1910		<i>trifasciata</i> (Zetterstedt, 1848)	B M
<i>puncticollis</i> (Zetterstedt, 1848)	B S	<i>Trichieurina</i> Duda, 1933	
<i>Eurina</i> Meigen, 1830		<i>pubescens</i> (Meigen, 1830)	B M S
<i>calva</i> Egger, 1862	B M		
<i>ducalis</i> A. Costa, 1885	M		
<i>Eutropha</i> Loew, 1866		HELEOMYZIDAE	
<i>variegata</i> Loew, 1866	S	Vladislav Martinek	
<i>Homalura</i> Meigen, 1826			
<i>tarsata</i> Meigen, 1826	B M S	HELEOMYZINAE	
<i>Lusiosina</i> Becker, 1910		BORBOROPSIDINI	
<i>albipila</i> (Loew, 1866)	B M S	<i>Borboropsis</i> Czerny, 1902	
<i>brevistylata</i> Dely-Draskovits, 1977	M S	<i>puberula</i> (Zetterstedt, 1838)	B M S
<i>cinctipes</i> (Meigen, 1830)	B M S		
<i>immaculata</i> Becker, 1912	M S	HELEOMYZINI	
<i>intermedia</i> Dely-Draskovits, 1977	B M S	<i>Acantholeria</i> Garret, 1921	
<i>nigripes</i> Dely-Draskovits, 1977	S	<i>cineraria</i> (Loew, 1862)	B M
<i>subnigripes</i> Dely-Draskovits, 1977	B	<i>Heleomyza</i> Fallén, 1810	
<i>Melanum</i> Becker, 1910		sg. <i>Anypotacta</i> Czerny, 1924	
<i>laterale</i> (Haliday, 1833)	S	<i>setulosa</i> (Czerny, 1924)	B S
<i>Meromyza</i> Meigen, 1830		sg. <i>Heleomyza</i> s.str.	
<i>bohémica</i> Fedoseeva, 1962	B S	<i>captiosa</i> (Gorodkov, 1962)	B M S
<i>curvinervis</i> (Zetterstedt, 1848)	M	<i>modesta</i> (Meigen, 1838)	B M S
<i>hybrida</i> Péterfi, 1961		<i>serrata</i> (Linnaeus, 1758)	B S
<i>elbergi</i> Fedoseeva, 1979	M	<i>Morpholeria</i> Garret, 1921	
<i>femorata</i> Macquart, 1835	B M S	sg. <i>Morpholeria</i> s.str.	
<i>laeta</i> Meigen, 1838	B M S	<i>dudai</i> (Czerny, 1924)	B M S
<i>mosquensis</i> Fedoseeva, 1960	M	sg. <i>Spanoparea</i> Czerny, 1924	
<i>nigriseta</i> Fedoseeva, 1960	B M S	<i>innotata</i> (Czerny, 1933)	B M
<i>nigriventris</i> Macquart, 1835	B M S	<i>kerteszi</i> (Czerny, 1924)	B M S
<i>palposa</i> Fedoseeva, 1960	S	<i>ruficornis</i> (Meigen, 1830)	B M S
<i>pluriseta</i> Péterfi, 1961	B	<i>variabilis</i> (Loew, 1862)	B M S
<i>pratorum</i> Meigen, 1830	B M S	<i>Neoleria</i> Malloch, 1919	
<i>saltatrix</i> (Linnaeus, 1761)	B M S	<i>flavicornis</i> (Loew, 1862)	M S
<i>sororecula</i> Fedoseeva, 1962	B	<i>inscripta</i> (Meigen, 1830)	B M S
<i>triangulina</i> Fedoseeva, 1960	B M	<i>ruficauda</i> (Zetterstedt, 1847)	B M S
<i>variegata</i> Meigen, 1830	B M S	<i>ruficeps</i> (Zetterstedt, 1838)	B M
<i>zachvatkini</i> Fedoseeva, 1960	M S	<i>Scoliocentra</i> Loew, 1862	
<i>Metopostigma</i> Becker, 1903		sg. <i>Chaetomus</i> Czerny, 1924	
<i>polonicum</i> (Schnabl, 1884)	S	<i>confusa</i> (Wahlgren, 1918)	B M S
<i>Neohaplegis</i> Beschovski, 1981		<i>flavotestacea</i> (Zetterstedt, 1838)	B M S

sg. <i>Gymnomus</i> Loew, 1893			<i>imberbis</i> Czerny, 1924	B	M	S
<i>caesia</i> (Meigen, 1830)	B	M	<i>inornata</i> (Loew, 1862)	B	M	S
<i>ceianui</i> Martinec, 1985	B		<i>laevifrons</i> (Loew, 1862)	B	M	S
<i>czernyi</i> (Papp & Woznica, 1993)	B		<i>lurida</i> (Meigen, 1830)	B	M	S
<i>europaea</i> (Papp & Woznica, 1993)	B	M	<i>mikii</i> (Pokorny, 1886)	B	M	S
<i>sabroskyi</i> (Gill, 1962)	B		<i>nemorum</i> (Meigen, 1830)	B	M	S
<i>spectabilis</i> (Loew, 1862)	B	M	<i>notata</i> (Meigen, 1830)	B		S
sg. <i>Leriola</i> Gorodkov, 1962			<i>oldenbergii</i> (Czerny, 1904)	B	M	S
<i>brachypterna</i> (Loew, 1873)	B	M	<i>oxyphora</i> (Mik, 1900)	B	M	S
<i>nigrinervis</i> (Wahlgren, 1918)			<i>pallida</i> (Fallén, 1820)	B	M	S
sg. <i>Scoliocentra</i> s.str.			<i>parva</i> (Loew, 1862)	B	M	S
<i>amplicornis</i> (Czerny, 1924)	B	M	<i>pilimana</i> (Loew, 1862)	B	M	S
<i>dupliciseta</i> (Strobl, 1894)	B	M	<i>quadrilineata</i> Czerny, 1924	B	M	S
<i>scutellaris</i> (Zetterstedt, 1838)	B		<i>similis</i> (Meigen, 1838)	B	M	
<i>villosa</i> (Meigen, 1830)	B	M	<i>stroblii</i> (Czerny, 1904)			S
<i>Schroederella</i> Enderlein, 1920			<i>umbratica</i> (Meigen, 1838)	B	M	S
<i>iners</i> (Meigen, 1830)	B	M	<i>univittata</i> (von Roser, 1840)	B	M	S
<i>minuta</i> Papp & Carles-Tolrá, 1994		M	<i>ustulata</i> (Meigen, 1830)	B	M	S
			<i>vaginata</i> (Loew, 1862)	B	M	S
			<i>variegata</i> (Loew, 1862)			S
OECOTHEINI						
<i>Eccoptomera</i> Loew, 1862			HETEROMYZINAE			
<i>emarginata</i> Loew, 1862	B	M	<i>Heteromyza</i> Fallén, 1820			
<i>filata</i> Loew, 1862	B	M	<i>atricornis</i> Meigen, 1830			S
<i>infusata</i> Wahlgren, 1918	B		<i>commixta</i> Collin, 1901	B	M	S
<i>longiseta</i> (Meigen, 1830)	B	M	<i>oculata</i> Fallén, 1820	B	M	S
<i>microps</i> (Meigen, 1830)	B	M	<i>rotundicornis</i> (Zetterstedt, 1846)	B	M	S
<i>nigricornis</i> Strobl, 1906	B		<i>Tephrochlaena</i> Czerny, 1924			
<i>obscura</i> (Meigen, 1830)	B	M	<i>halterata</i> (Meigen, 1830)	B		
<i>ornata</i> Loew, 1862	B	M	<i>Tephrochlamys</i> Loew, 1862			
<i>pallescens</i> (Meigen, 1830)	B	M	<i>flavipes</i> (Zetterstedt, 1838)	B	M	S
<i>triseia</i> Czerny, 1924	B		<i>laeta</i> (Meigen, 1838)	B		S
<i>Oecothia</i> Haliday, 1837			<i>rufiventris</i> (Meigen, 1830)	B	M	S
<i>fenestralis</i> (Fallén, 1820)	B	M	<i>tarsalis</i> (Zetterstedt, 1847)	B	M	S
<i>praecox</i> Loew, 1862		M				
ORBELLIINI			TRIXOSCELIDIDAE			
<i>Orbellia</i> Robineau-Desvoidy, 1830			Jindřich Roháček			
<i>hiemalis</i> (Loew, 1862)	B					
<i>myiopsiformis</i> Robineau-Desvoidy, 1830	B	M				
<i>Oldenbergiella</i> Czerny, 1924			<i>Trixoscelis</i> Rondani, 1856			
<i>brunalis</i> Czerny, 1924	B	M	<i>frontalis</i> (Fallén, 1823)	B	M	S
<i>calcarifera</i> Papp, 1980	B	M	<i>marginella</i> (Fallén, 1823)	B	M	S
<i>seticercus</i> Papp, 1980	B		<i>obscura</i> (Fallén, 1823)	B	M	S
			<i>similis</i> Hackman, 1970	B	M	S
SUILLIINAE						
<i>Suillia</i> Robineau-Desvoidy, 1830			CHYROMYIDAE			
<i>affinis</i> (Meigen, 1830)	B	M	Jindřich Roháček			
<i>atricornis</i> (Meigen, 1830)	B	M				
<i>bicolor</i> (Zetterstedt, 1838)	B	M	<i>Aphaniosoma</i> Becker, 1903			
<i>cepelaki</i> Martinec, 1985			<i>propinquans</i> Collin, 1949	B		
<i>crinimana</i> (Czerny, 1904)			<i>socium</i> Collin, 1949	B		
<i>femoralis</i> (Loew, 1862)	B	M	<i>unicolor</i> Hendel, 1933	B		
<i>flava</i> (Meigen, 1830)	B	M	<i>Chyromya</i> Robineau-Desvoidy, 1830			
<i>flavifrons</i> (Zetterstedt, 1838)	B	M	<i>femorella</i> (Fallén, 1820)	B	M	
<i>fuscicornis</i> (Zetterstedt, 1847)	B	M	<i>flava</i> (Linnaeus, 1758)	B	M	S
<i>gigantea</i> (Meigen, 1830)	B	M	<i>miladae</i> Andersson, 1976	B	M	S
<i>humilis</i> (Meigen, 1830)		M				

<i>oppidana</i> (Scopoli, 1763)	B M S	<i>Sphaerocera</i> Latreille, 1804	
<i>Gymnochiromyia</i> Hendel, 1933		<i>curvipes</i> Latreille, 1805	B M S
<i>flavella</i> (Zetterstedt, 1848)	S	<i>monilis</i> Haliday, 1836	B M S
<i>inermis</i> (Collin, 1933)	B M S		
SPHAEROCERIDAE		LIMOSININAE	
Jindřich Roháček		<i>Apteromyia</i> Vimmer, 1929	
		<i>claviventris</i> (Strobl, 1909)	B M S
COPROMYZINAE		<i>Aptilotus</i> Mik, 1894	
<i>Alloborborus</i> Duda, 1923		<i>paradoxus</i> Mik, 1894	B M S
<i>pallifrons</i> (Fallén, 1820)	B M S	<i>Chaetopodella</i> Duda, 1920	
<i>Borborillus</i> Duda, 1923		<i>scutellaris</i> (Haliday, 1836)	B M S
<i>uncinatus</i> (Duda, 1923)	M S	<i>Coproica</i> Rondani, 1861	
<i>viripennis</i> (Meigen, 1830)	S	<i>acutangula</i> (Zetterstedt, 1847)	B M S
<i>Copromyza</i> Fallén, 1810		<i>digitata</i> (Duda, 1918)	M S
<i>equina</i> Fallén, 1820	B M S	<i>ferruginata</i> (Stenhammar, 1855)	B M S
<i>montana</i> Roháček, 1992	B M S	<i>hirticula</i> Collin, 1956	B M S
<i>neglecta</i> (Malloch, 1913)	B M S	<i>hirtula</i> (Rondani, 1880)	B M S
= <i>calcitrans</i> (Spuler, 1925)		<i>lugubris</i> (Haliday, 1836)	B M S
= <i>parastercoraria</i> Papp, 1973		<i>pustio</i> (Zetterstedt, 1847)	M S
<i>nigrina</i> (Gimmerthal, 1847)	B M S	<i>vagans</i> (Haliday, 1833)	B M S
= <i>similis</i> (Collin, 1930)		<i>Elachisoma</i> Rondani, 1880	
<i>stercoraria</i> (Meigen, 1830)	B M S	<i>aterrimum</i> (Haliday, 1833)	B M S
<i>Crumomyia</i> Macquart, 1835		<i>bajzae</i> Papp, 1983	B M S
<i>glabrifrons</i> (Meigen, 1830)	B M S	<i>pilosum</i> (Duda, 1924)	B M S
<i>fimetaria</i> (Meigen, 1830)	B M S	<i>Gigalimosina</i> Roháček, 1983	
<i>nigra</i> (Meigen, 1830)	B M S	<i>flaviceps</i> (Zetterstedt, 1847)	B M S
<i>nitida</i> (Meigen, 1830)	B M S	<i>Gonioneura</i> Rondani, 1880	
<i>notabilis</i> (Collin, 1902)	B M S	= <i>Halidayina</i> Duda, 1918	
<i>parentela alpicola</i> Roháček in Troger		<i>spinipennis</i> (Haliday, 1836)	B M S
& Roháček, 1980	S	<i>Herniosina</i> Roháček, 1983	
<i>pedestris</i> (Meigen, 1830)	B M S	<i>bequaerti</i> (Villeneuve, 1917)	B M S
<i>prunosa</i> (Richards, 1932)	B M S	<i>horrida</i> (Roháček, 1978)	B M S
<i>rohaceki</i> Norrbom & Kim, 1985	B M S	<i>pollex</i> Roháček, 1993	B M S
= <i>glacialis</i> auct. Meigen, 1830		<i>Leptocera</i> Olivier, 1813	
<i>setitibialis</i> (Spuler, 1925)	S	sg. <i>Leptocera</i> s.str.	
= <i>freyi</i> (Hackman, 1965)		<i>alpina</i> Roháček, 1982	B S
<i>zuskai</i> (Roháček, 1976)	B M S	<i>caenosa</i> (Rondani, 1880)	B M S
<i>Lotophila</i> Lioy, 1864		<i>finalis</i> (Collin, 1956)	M
<i>atra</i> (Meigen, 1830)	B M S	<i>fontinalis</i> (Fallén, 1826)	B M S
<i>Norrbomia</i> Papp, 1988		<i>nigra</i> Olivier, 1813	B M S
<i>costalis</i> (Zetterstedt, 1847)	B M S	<i>oldenbergi</i> (Duda, 1918)	B M S
<i>nitidifrons</i> (Duda, 1923)	M	sg. <i>Rachispoda</i> Lioy, 1864	
<i>sordida</i> (Zetterstedt, 1847)	B M S	<i>anceps</i> (Stenhammar, 1855)	B M
		<i>breviceps</i> (Stenhammar, 1855)	B M S
SPHAEROCERINAE		<i>brevior</i> Roháček, 1991	M S
<i>Ischiolepta</i> Lioy, 1864		<i>cilifera</i> (Rondani, 1880)	M S
<i>crenata</i> (Meigen, 1838)	B M S	<i>duplex</i> Roháček, 1991	S
<i>denticulata</i> (Meigen, 1830)	B M	<i>fuscipennis</i> (Haliday, 1833)	B M S
<i>micropyga</i> (Duda, 1938)	B M S	<i>gel</i> Papp, 1978	B M S
<i>nitida</i> (Duda, 1920)	B M S	<i>hostica</i> Villeneuve, 1917	B M S
<i>pusilla</i> (Fallén, 1820)	B M S	<i>intermedia</i> (Duda, 1918)	S
<i>scabricula</i> (Haliday, 1836)	B M S	<i>limosa</i> (Fallén, 1820)	B M S
<i>vaporariorum</i> (Haliday, 1836)	B M S	<i>longior</i> Roháček, 1991	M S
<i>Lotobia</i> Lioy, 1864		<i>lugubrina</i> (Zetterstedt, 1847)	M S
<i>pallidiventris</i> (Meigen, 1830)	M S	<i>lutosa</i> (Stenhammar, 1855)	B M S
		<i>lutosoidea</i> (Duda, 1938)	B M S
		<i>modesta</i> (Duda, 1924)	B M S

<i>opinata</i> Roháček, 1991	M S	<i>Pteremis</i> Rondani, 1856	
<i>pseudohostica</i> (Duda, 1924)	M S	<i>fenestralis</i> (Fallén, 1820)	B M S
<i>segem</i> Roháček, 1991	B M S	<i>Pullimosina</i> Roháček, 1983	
<i>tuberosa</i> (Duda, 1938)	M	sg. <i>Dahlimosina</i> Roháček, 1983	
<i>Limosina</i> Macquart, 1835		<i>dahli</i> (Duda, 1918)	B M
<i>silvatica</i> (Meigen, 1830)	B M S	sg. <i>Pullimosina</i> s.str.	
<i>Minilimosina</i> Roháček, 1983		<i>antennata</i> (Duda, 1918)	B M S
sg. <i>Allolimosina</i> Roháček, 1983		<i>heteroneura</i> (Haliday, 1836)	B M S
<i>albinervis</i> (Duda, 1918)	S	<i>mejerei</i> (Duda, 1918)	B M S
<i>alloneura</i> (Richards, 1952)	M S	<i>moesta</i> (Villeneuve, 1918)	B M S
sg. <i>Minilimosina</i> s.str.		<i>pulhula</i> (Zetterstedt, 1847)	B M S
<i>bicuspis</i> Roháček, 1993	B M S	<i>Puncticorpus</i> Duda, 1918	
<i>fungicola</i> (Haliday, 1836)	B M S	<i>cribratum</i> (Villeneuve, 1918)	B M S
<i>gemella</i> Roháček, 1983	B M	<i>Rudolfina</i> Roháček, 1987	
<i>parvula</i> (Stenhammar, 1855)	B M S	= <i>Rudolfia</i> Roháček, 1982 nec Wilson, 1924	
<i>tenera</i> Roháček, 1983	M	<i>rozkosnyi</i> (Roháček, 1975)	M S
sg. <i>Svarciella</i> Roháček, 1983		<i>Spelobia</i> Spuler, 1924	
<i>floreni</i> Roháček & Marshall, 1988	M	sg. <i>Bifronsina</i> Roháček, 1983	
<i>splendens</i> (Duda, 1928)	B M S	<i>bifrons</i> (Stenhammar, 1855)	B M S
<i>v-atrum</i> (Villeneuve, 1917)	B M S	sg. <i>Eulimosina</i> Roháček, 1983	
<i>vitripennis</i> (Zetterstedt, 1847)	B M S	<i>ochripes</i> (Meigen, 1830)	B M S
<i>Opacifrons</i> Duda, 1918		sg. <i>Spelobia</i> s.str.	
<i>coxata</i> (Stenhammar, 1855)	B M S	<i>belanica</i> Roháček, 1983	M S
<i>maculifrons</i> (Becker, 1907)	M	<i>cambrica</i> (Richards, 1929)	B M S
<i>moravica</i> (Roháček, 1975)	M S	<i>clunipes</i> (Meigen, 1830)	B M S
<i>Opalimosina</i> Roháček, 1983		<i>czizeki</i> (Duda, 1918)	B M S
sg. <i>Dentilimosina</i> Roháček, 1983		<i>faeroensis</i> (Deeming, 1966)	S
<i>denticulata</i> (Duda, 1924)	B M S	<i>ibrida</i> Roháček, 1983	B M S
sg. <i>Hackmanina</i> Roháček, 1983		<i>luteilabris</i> (Rondani, 1880)	B M S
<i>czernyi</i> (Duda, 1918)	B M S	<i>manicata</i> (Richards, 1927)	B M S
sg. <i>Opalimosina</i> s.str.		<i>nana</i> (Rondani, 1880)	B M S
<i>ealcarifera</i> (Roháček, 1975)	M S	<i>palmata</i> (Richards, 1927)	B M S
<i>collini</i> (Richards, 1929)	B M S	<i>pappi</i> Roháček, 1983	B M
<i>mirabilis</i> (Collin, 1902)	B M S	<i>parapusio</i> (Dahl, 1909)	B M S
<i>simplex</i> (Richards, 1929)	B M S	<i>pseudonivalis</i> (Dahl, 1909)	B M
sg. <i>Pappiella</i> Roháček, 1983		<i>pseudosetaria</i> (Duda, 1918)	B M S
<i>liliputana</i> (Rondani, 1880)	B M S	<i>rufilabris</i> (Stenhammar, 1855)	B M S
sg. <i>Paralimosina</i> Papp, 1973		<i>simplicipes</i> (Duda, 1925)	S
<i>fucata</i> (Rondani, 1880)	B M S	<i>talis</i> Roháček, 1983	B S
<i>kaszabi</i> Papp, 1973	B M S	<i>talparum</i> (Richards, 1927)	B M S
= <i>latigena</i> (Roháček, 1977)		<i>Spinilimosina</i> Roháček, 1983	
<i>suberibrata</i> (Roháček, 1977)	B M S	<i>brevicostata</i> (Duda, 1918)	M
<i>trichopyga</i> (Richards, 1952)	B M S	<i>Telomerina</i> Roháček, 1983	
<i>Philocoprella</i> Richards, 1929		<i>eburnea</i> Roháček, 1983	B M
<i>italica</i> (Deeming, 1964)	S	<i>flavipes</i> (Meigen, 1830)	B M S
<i>quadrispina</i> (Laurence, 1952)	M S	<i>levifrons</i> (Spuler, 1925)	M S
<i>Phthitia</i> Enderlein 1938		= <i>antonini</i> Roháček, 1983	
sg. <i>Alimosina</i> Roháček, 1983		<i>pseudoleucoptera</i> (Duda, 1924)	B M S
<i>empirica</i> (Hutton, 1901)	B M S	<i>ursina</i> Roháček, 1983	S
sg. <i>Collimosina</i> Roháček, 1983		<i>Terrilimosina</i> Roháček, 1983	
<i>spinosa</i> (Collin, 1930)	B M	<i>racovitzae</i> (Bezzi, 1911)	B M S
sg. <i>Kimosina</i> Roháček, 1983		<i>schmitzi</i> (Duda, 1918)	B M S
<i>longisetosa</i> (Dahl, 1909)	B M S	<i>sudetica</i> (Roháček, 1975)	B M S
<i>phumosula</i> (Rondani, 1880)	B M S	<i>Thoracochaeta</i> Duda, 1918	
<i>Pseudocollinella</i> Duda, 1924		<i>brachystoma</i> (Stenhammar, 1855)	B M
= <i>Opacifrons</i> auct. nec Duda, 1918		<i>zosteriae</i> (Haliday, 1833)	B M S
<i>humida</i> (Haliday, 1836)	B M S		

<i>Trachyopella</i> Duda, 1918				sg. <i>Amiota</i> s. str.			
sg. <i>Insulomyia</i> Papp, 1972				<i>albilabris</i> (Roth in Zetterstedt, 1860)	B	M	
<i>artivena</i> Roháček & Marshall, 1986	B	M	S	<i>alboguttata</i> (Wahlberg, 1838)	B	M	S
sg. <i>Minuscula</i> Roháček & Marshall, 1986				<i>basdeni</i> Fonseca, 1965	B	M	S
<i>minuscula</i> Collin, 1956	B	M		<i>collini</i> Beuk & Máca, 1995	B		S
sg. <i>Nudopella</i> Roháček & Marshall, 1986				<i>filipes</i> Máca, 1980	B	M	
<i>hem</i> Roháček & Marshall, 1986	B			<i>flavopruinosa</i> Duda, 1934	B		
<i>leucoptera</i> (Haliday, 1836)	B	M	S	<i>rufescens</i> (Oldenberg, 1914)	B		S
sg. <i>Trachyopella</i> s.str.				<i>subtusradiata</i> Duda, 1934		M	
<i>atomus</i> (Rondani, 1880)	B	M	S	sg. <i>Phortica</i> Schiner, 1862			
<i>bovilla</i> Collin, 1954		M	S	<i>semivirgo</i> Máca, 1977	B	M	S
<i>coprina</i> (Duda, 1918)	B			<i>variegata</i> (Fallén, 1823)	B	M	S
<i>kuntzei</i> (Duda, 1918)	B	M	S	<i>Cacoxenus</i> Loew, 1858			
<i>lineafrons</i> (Spuler, 1925)	B	M	S	sg. <i>Cacoxenus</i> s. str.			
<i>melania</i> (Haliday, 1836)	B	M	S	<i>indagator</i> Loew, 1858	B	M	
<i>straminea</i> Roháček & Marshall, 1986	B	M	S	sg. <i>Paracacoxenus</i> D. E. Hardy, 1960			
				<i>kaszabi</i> (Okada, 1973)	B		S
				<i>Gitona</i> Meigen, 1830			
				<i>distigma</i> Meigen, 1830	B	M	S
CURTONOTIDAE							
Jan Máca				DROSOPHILINAE			
<i>Curtonotum</i> Macquart, 1843				<i>Chymomyza</i> Czerny, 1903			
<i>anus</i> (Meigen, 1830)		M	S	<i>amoena</i> (Loew, 1862)	B	M	S
				<i>caudatula</i> Oldenberg, 1914	B	M	S
CAMILLIDAE				<i>costata</i> (Zetterstedt, 1838)	B	M	S
Jan Máca				<i>distincta</i> (Egger, 1862)	B	M	S
<i>Camilla</i> Haliday, 1838				<i>fuscimana</i> (Zetterstedt, 1838)	B	M	S
<i>atrimana</i> Strobl, 1910	B	M	S	<i>Drosophila</i> Fallén, 1823			
<i>flavicauda</i> Duda, 1922	B	M		sg. <i>Dorsilopha</i> Sturtevant, 1942			
<i>glabrata</i> Collin, 1956				<i>busckii</i> Coquillett, 1901	B	M	S
<i>glabra</i> (Fallén, 1823)	B	M	S	sg. <i>Drosophila</i> s.str.			
<i>nigrifrons</i> Collin, 1933	B			<i>funbris</i> (Fabricius, 1789)	B	M	S
				<i>histrio</i> Meigen, 1830	B	M	S
DROSOPHILIDAE				<i>hydei</i> Sturtevant, 1921	B	M	S
Jan Máca				<i>immigrans</i> Sturtevant, 1921	B	M	S
STEGANINAE				<i>kuntzei</i> Duda, 1924	B	M	S
STEGANINI				<i>limbata</i> von Roser, 1840	B	M	S
<i>Leucophenga</i> Mik, 1886				<i>littoralis</i> Meigen, 1830	B	M	
<i>maculata</i> (Dufour, 1839)	B	M	S	<i>hummei</i> Hackman, 1972	B	M	S
<i>quinquemaculata</i> Strobl, 1893	B	M	S	<i>phalerata</i> Meigen, 1830	B	M	S
<i>Stegana</i> Meigen, 1830				<i>picta</i> Zetterstedt, 1847		M	S
sg. <i>Stegana</i> s. str.				<i>repleta</i> Wollaston, 1858	B	M	S
<i>furta</i> (Linnaeus, 1767)	B	M	S	<i>testacea</i> von Roser, 1840	B	M	S
sg. <i>Steganina</i> Wheeler, 1960				<i>transversa</i> Fallén, 1823	B	M	S
<i>coleoprata</i> (Scopoli, 1763)	B		S	<i>unimaculata</i> Strobl, 1893	B	M	S
<i>hypoleuca</i> Meigen, 1830	B		S	<i>virilis</i> Sturtevant, 1921	B	M	S
<i>longifibula</i> Takada, 1968	B	M	S	sg. <i>Sophophora</i> Sturtevant, 1939			
<i>mehadiae</i> Duda, 1924	B	M	S	<i>alpina</i> Burla, 1948	B		S
<i>nigrithorax</i> Strobl, 1898			S	<i>ambigua</i> Pomini, 1940	B	M	S
<i>similis</i> Laštovka & Máca, 1982	B	M	S	<i>bifasciata</i> Pomini, 1940	B	M	
				<i>helvetica</i> Burla, 1948	B		
GITONINI				<i>melanogaster</i> Meigen, 1830	B	M	S
<i>Acetoxenus</i> Frauenfeld, 1868				<i>obscura</i> Fallén, 1823	B	M	S
<i>formosus</i> (Loew, 1846)	B			<i>silvestris</i> Basden, 1954	B	M	S
<i>Amiota</i> Loew, 1862				= <i>subsilvestris</i> D.E. Hardy & Kaneshiro, 1968			
				<i>simulans</i> Sturtevant, 1919	B	M	
				<i>subobscura</i> Collin in Gordon, 1936	B	M	S
				<i>tristis</i> Fallén, 1823	B	M	

sg. <i>Spinodrosophila</i> Duda, 1924		<i>obscuripennis</i> (Meigen, 1830)	B M S
<i>nigrosparva</i> Strobl, 1898	B M S	<i>punctum</i> (Meigen, 1830)	B M S
Hirtodrosophila Duda, 1924			
<i>cameraria</i> (Haliday, 1833)	B M		
<i>confusa</i> (Staeger, 1844)	B M S	EPHYDRIDAE	
<i>lundstroemi</i> (Duda, 1935)	S	Tadeusz Zatwarnicki	
<i>oldenbergi</i> (Duda, 1924)	S		
<i>toyohiokadai</i> (Sidorenko, 1990)	S	DISCOMYZINAE	
<i>trivittata</i> (Strobl, 1893)	B M	DISCOMYZINI	
Lordiphosa Basden, 1961		<i>Discomyza</i> Meigen, 1830	
<i>acuminata</i> (Collin, 1952)	M S	<i>incurva</i> (Fallén, 1813)	B M S
<i>andalusiaca</i> (Strobl, 1906)	B M S		
<i>fenestrarum</i> (Fallén, 1823)	B M S	PSILOPINI	
<i>hexasticha</i> (Papp, 1971)	B M S	<i>Clanoneurum</i> Becker, 1903	
<i>miki</i> (Duda, 1924)	M S	<i>cimiciforme</i> (Haliday, 1855)	S
Microdrosophila Malloch, 1921		Cnestrum Becker, 1896	
<i>congesta</i> (Zetterstedt, 1847)	B M S	<i>lepidopis</i> Becker, 1896	M S
Mycodrosophila Oldenberg, 1914		Psilopa Fallén, 1823	
<i>poecilogastra</i> (Loew, 1874)	S	<i>apicalis</i> (Perris, 1847)	S
Scaptodrosophila Duda, 1923		<i>compta</i> (Meigen, 1830)	B M S
<i>defflexa</i> (Duda, 1924)	B M S	<i>leucostoma</i> (Meigen, 1830)	B
<i>rufifrons</i> (Loew, 1873)	B M S	<i>marginella</i> Fallén, 1823	M S
Scaptomyza J. Hardy, 1849		<i>nana</i> Loew, 1860	B M
sg. <i>Parascaptomyza</i> Duda, 1924		<i>nigritella</i> Stenhammar, 1844	B M
<i>pallida</i> (Zetterstedt, 1847)	B M S	<i>nitidula</i> (Fallén, 1813)	B M S
sg. <i>Scaptomyza</i> s. str.		<i>pappi</i> Canzoneri & Meneghini, 1965	S
<i>consimilis</i> Hackman, 1955	B	<i>polita</i> (Macquart, 1835)	B M S
<i>flava</i> (Fallén, 1823)	B M S	<i>pulicaria</i> (Haliday, 1839)	B
<i>graminum</i> (Fallén, 1823)	B M S	Trimerina Macquart, 1835	
<i>griseola</i> (Zetterstedt, 1847)	B M S	<i>madizans</i> (Fallén, 1813)	B M S
		<i>microchaeta</i> Hendel, 1932	M S
DIASTATIDAE			
Jindřich Roháček & Jan Mác		HYDRELLIINAE	
		ATISSINI	
DIASTATINAE		<i>Atissa</i> Haliday in Curtis, 1837	
Diastata Meigen, 1830		<i>limosina</i> Becker, 1896	M
<i>Trichoptera</i> Lioy, 1864		<i>pygmaea</i> (Haliday, 1839)	B S
= <i>Callopteryella</i> Coquillett, 1910		Ptilomyia Coquillett, 1900	
<i>adusta</i> Meigen, 1830	B M S	<i>orsovana</i> Enderlein, 1922	B M S
= <i>unipunctata</i> Zetterstedt, 1847			
<i>cervinala</i> Chandler, 1987	B M S	HYDRELLIINI	
<i>costata</i> Meigen, 1830	B M S	Hydrellia Robineau-Desvoidy, 1830	
<i>fuscula</i> auct. nec Fallén, 1823		<i>albifrons</i> (Fallén, 1813)	M
<i>modesta</i> Melander, 1913		= <i>hydrocharitis</i> Hering, 1925	
<i>flavivosta</i> Chandler, 1987	B M S	<i>albilabris</i> (Meigen, 1830)	B S
<i>fuscula</i> (Fallén, 1823)	B M S	<i>argyrogenis</i> Becker, 1896	B
<i>fulvifrons</i> Haliday, 1837		<i>cardamines</i> Haliday, 1839	B S
= <i>ornata</i> Loew, 1864		<i>concolor</i> (Stenhammar, 1844)	M
<i>nebulosa</i> (Fallén, 1823)	B M S	= <i>butomi</i> Hering, 1925	
= <i>ornata</i> auct. nec Meigen, 1830		<i>fascitibia</i> (von Roser, 1840)	B
<i>ornata</i> Meigen, 1830	B M	<i>flaviceps</i> (Meigen, 1830)	B S
<i>vagans</i> Loew, 1864	B M S	= <i>discoirs</i> Collin, 1966	
		<i>flavicornis</i> (Fallén, 1823)	B S
CAMPICHOETINAE		<i>frontalis</i> Loew, 1860	B
Campichoeta Macquart, 1835		<i>frontosa</i> Becker, 1926	S
<i>griseola</i> (Zetterstedt, 1855)	B M S	<i>fulviceps</i> (Stenhammar, 1844)	B M S
		<i>fusca</i> (Stenhammar, 1844)	B

EPHYDRINAE

PARYDRINI

Eutaenionotum Oldenberg, 1923*guttipennis* (Stenhammar, 1844)*Parydra* Stenhammar, 1844sg. *Chaetoapnea* Hendel, 1930*fossarium* (Haliday, 1833)*hecate* (Haliday, 1833)*mitis* (Cresson, 1930)*pusilla* (Meigen, 1830)sg. *Parydra* s.str.*aquila* (Fallén, 1813)*coarctata* (Fallén, 1813)*littoralis* (Meigen, 1830)*nigritarsis* Strobl, 1893*mibecula* Becker, 1896*quadripunctata* (Meigen, 1830)*quinquemaculata* Becker, 1896*unicolor* (Becker, 1926)

EPHYDRINI

Coenia Robineau-Desvoidy, 1830*curvicauda* (Meigen, 1830)*palustris* (Fallén, 1823)*Ephydra* Fallén, 1810*macellaria* Egger, 1862*riparia* Fallén, 1813*Paracoenia* Cresson, 1935*fumosa* (Stenhammar, 1844)*Setacera* Cresson, 1930*aurata* (Stenhammar, 1844)*breviventris* (Loew, 1860)*micans* (Haliday, 1833)*trina* Collin, 1964

SCATELLINI

Haloscatella Mathis, 1979*dichaeta* (Loew, 1860)*Lamproscatella* Hendel, 1917*bimaculata* Hendel, 1933*sibilans* (Haliday, 1833)*Limmellia* Malloch, 1925*fallax* (Czerny, 1910)*quadrata* (Fallén, 1813)*stenhammari* (Zetterstedt, 1846)*Philotelma* Becker, 1896*defecta* (Haliday, 1833)*nigripenne* (Meigen, 1830)*strandi* Duda, 1942*Scatella* Robineau-Desvoidy, 1830sg. *Neoscatella* Malloch, 1933*crassicauda* Becker, 1896= *lundbergi* Dahl, 1959*subguttata* (Meigen, 1830)sg. *Scatella* s.str.*indistincta* Becker, 1896*paludum* (Meigen, 1830)*stagnalis* (Fallén, 1813)*tenuicosta* Collin, 1930*Scatophila* Becker, 1896*caviceps* (Stenhammar, 1844)*contaminata* (Stenhammar, 1844)= *hamifera* Becker, 1896*cribrata* (Stenhammar, 1844)*despecta* (Haliday, 1839)*iowana* Wheeler, 1961*noctula* (Meigen, 1830)= *silesiaca* Becker, 1926= *nasuta* Papp, 1975*quadriguttata* (Meigen, 1830)= *variegata* Loew, 1860*unicornis* Czerny, 1900

SCATHOPHAGIDAE

František Šifner

NORELLIINAE

Norellisoma Wahlgren, 1917*alpestre* (Schiner, 1864)*femorale* (Loew, 1864)*litturatum* (Wiedemann in Meigen, 1826)*mirusae* Šifner, 1974*nervosum* (Meigen, 1826)*spinimamum* (Fallén, 1819)*striolatum* (Meigen, 1826)*Norellia* Robineau-Desvoidy, 1830*spinipes* (Meigen, 1826)

CORDILURINAE

Leptopa Zetterstedt, 1838*filiformis* Zetterstedt, 1838*Americina* Malloch, 1923sg. *Chylizosoma* Hendel, 1924*media* (Becker, 1894)*vittata* (Meigen, 1826)*Parallelomma* Becker, 1894*albipes* (Fallén, 1819)*Phrosia* Robineau-Desvoidy, 1830*albilabris* (Fabricius, 1794)*Megaphthalma* Becker, 1894*pallida* (Fallén, 1819)*Cordilura* Fallén, 1819*aemula* Collin, 1958*atrata* (Zetterstedt, 1846)*ciliata* (Meigen, 1826)*picipes* (Meigen, 1826)*pubera* (Linnaeus, 1758)*pudica* (Meigen, 1826)*rufimana* (Meigen, 1826)*umbrosa* (Loew, 1873)*Micropselapha* Becker, 1894*filiformis* (Zetterstedt, 1846)*Scoliaphleps* Becker, 1894*ustulata* (Zetterstedt, 1838)

<i>Gonatherus</i> Rondani, 1856			<i>obscurum</i> (Fallén, 1819)	B	M	S
<i>planiceps</i> (Fallén, 1819)		M	<i>tinctinervis</i> Becker, 1894			B
<i>Megaphthalmoides</i> Ringdahl, 1936			<i>Scathophaga</i> Meigen, 1803			
<i>unilineata</i> (Zetterstedt, 1838)		B	<i>analís</i> (Meigen, 1826)		B	M
<i>Nanna</i> Becker, 1894			<i>cineraria</i> (Meigen, 1826)		B	M
<i>armillata</i> (Zetterstedt, 1846)		B	<i>furcata</i> (Say, 1823)		B	M
<i>articulata</i> (Becker, 1894)		B	<i>inquinata</i> (Meigen, 1826)		B	M
<i>brevifrons</i> (Zetterstedt, 1838)		B	<i>lutaria</i> (Fabricius, 1794)		B	M
<i>fasciata</i> (Meigen, 1926)		B	<i>pictipennis</i> (Oldenberg, 1923)		B	
<i>flavipes</i> (Fallén, 1819)		B	<i>scybalaria</i> (Linnaeus, 1758)		B	M
<i>inermis</i> (Becker, 1894)		B	<i>suilla</i> (Fabricius, 1794)		B	M
<i>leucostoma</i> (Zetterstedt, 1846)		B	<i>stercoraria</i> (Linnaeus, 1758)		B	M
<i>minuta</i> (Becker, 1894)		B				
<i>puberulla</i> (Becker, 1894)		B				
<i>tibiella</i> (Zetterstedt, 1838)		B				
<i>Gonarticus</i> Becker, 1894			DELININAE			
<i>abdominalis</i> (Zetterstedt, 1846)		B	<i>Delina</i> Robineau-Desvoidy, 1830			
<i>Spathephilus</i> Becker, 1894			<i>anthrax</i> (Schiner, 1864)		B	M
<i>breviventris</i> (Loew, 1873)		B	<i>nigriceps</i> (Becker, 1894)		B	
<i>Orthacheta</i> Becker, 1894			<i>nigrita</i> (Fallén, 1819)		B	M
<i>pilosa</i> (Zetterstedt, 1838)		B				
<i>Cleigastrea</i> Macquart, 1835			ANTHOMYIIDAE			
<i>apicalis</i> (Meigen, 1826)		B	Rudolf Rozkošný			
			<i>Acklandia</i> Hennig, 1976			
HYDROMYZINAE			<i>servadei</i> (Séguy, 1933)		M	S
<i>Cochliarium</i> Becker, 1894			<i>Acyglossa</i> Rondani, 1866			
<i>castanipes</i> Becker, 1894		S	<i>atramentaria</i> (Meigen, 1826)			S
<i>cuneiventris</i> (Zetterstedt, 1846)		S	<i>Adia</i> Robineau-Desvoidy, 1830			
<i>Gymnomera</i> Rondani, 1867			<i>cinerella</i> (Fallén, 1825)		B	M
<i>dorsata</i> (Zetterstedt, 1838)		S	<i>Alliopsis</i> Schnabl & Dziedzicki, 1911			
<i>Microprosopa</i> Becker, 1894			<i>billbergi</i> (Zetterstedt, 1838)		B	M
<i>haemorrhoidalis</i> (Meigen, 1826)		M	<i>brunneigena</i> (Schnabl, 1915)			M
<i>hoberlandti</i> Šifner, 1981		M	<i>conifrons</i> (Zetterstedt, 1845)			M
<i>pallidicauda</i> (Zetterstedt, 1838)		B	<i>pilitarsis</i> (Stein, 1900)			S
<i>Spaziphora</i> Rondani, 1856			<i>sepiella</i> (Zetterstedt, 1845)			M
<i>hydromyzina</i> (Fallén, 1819)		B	<i>silvestris</i> (Fallén, 1824)		B	M
<i>Trichopampus</i> Rondani, 1856			<i>Anthomyia</i> Meigen, 1803			
<i>fraternus</i> (Meigen, 1826)		B	<i>cannabina</i> (Stein, 1916)		B	
<i>Chaetosa</i> Coquillett, 1898			<i>confusanea</i> Michelsen in Michelsen			
<i>punctipes</i> (Meigen, 1826)		B	& Baez, 1985		B	M
<i>Acerocnema</i> Becker, 1894			<i>imbrida</i> Rondani, 1866		B	M
<i>macrocera</i> (Meigen, 1826)		B	<i>liturata</i> (Robineau-Desvoidy, 1830)		B	M
<i>Cosmetopus</i> Becker, 1894			<i>monilis</i> (Meigen, 1826)		B	M
<i>dentimanus</i> (Zetterstedt, 1838)		B	<i>plurinotata</i> Brullé, 1832		B	
<i>Pogonota</i> Zetterstedt, 1860			<i>pluvialis</i> (Linnaeus, 1758)		B	M
<i>barbata</i> (Zetterstedt, 1838)		B	<i>procellaris</i> Rondani, 1866		B	M
<i>Hexamitocera</i> Becker, 1894			<i>quinquemaculata</i> Macquart, 1839			M
<i>loxocerata</i> (Fallén, 1819)		B	<i>Botanophila</i> Lioy, 1864			
<i>Acanthocnema</i> Becker, 1894			<i>biciliaris</i> (Pandellé, 1900)		B	M
<i>glauescens</i> (Loew, 1864)		M	<i>brunneilinea</i> (Zetterstedt, 1845)		B	M
<i>Hydromyza</i> Fallén, 1819			<i>discreta</i> (Meigen, 1826)		B	
<i>livens</i> (Fabricius, 1794)		B	<i>dissecta</i> (Meigen, 1826)		B	M
SCATHOPHAGINAE			<i>flavisquama</i> (Stein, 1906)		B	
<i>Coniosternum</i> Becker, 1894			<i>fugax</i> (Meigen, 1826)		B	M
<i>lapponicum</i> Ringdahl, 1920		B	<i>gentianae</i> (Pandellé, 1900)		B	M
			<i>gnava</i> (Meigen, 1826)		B	M

<i>hucketti</i> (Ringdahl, 1935)	B	<i>nigrescens</i> Rondani, 1877	B	S
<i>laterella</i> (Collin, 1967)	B	<i>nuda</i> (Strobl, 1901)	B	M S
<i>latifrons</i> (Zetterstedt, 1845)	B M S	<i>penicillaris</i> (Rondani, 1866)	B	M S
<i>lobata</i> (Collin, 1967)	M S	<i>penicillosa</i> Hennig, 1974		S
<i>monticola</i> (Karl, 1932)	M	<i>planipalpis</i> (Stein, 1898)		M
<i>phreomine</i> (Séguy, 1937)	B M S	<i>platura</i> (Meigen, 1826)	B	M S
<i>profuga</i> (Stein, 1916)	M S	<i>radicum</i> (Linnaeus, 1758)	B	M S
<i>retusa</i> (Ringdahl, 1952)	M	<i>segmentata</i> (van der Wulp, 1896)		S
<i>rubrifrons</i> (Ringdahl, 1933)		<i>tarsifimbria</i> (Pandellé, 1900)	B	
<i>rubrigena</i> (Schnabl, 1915)		<i>tenuiventris</i> (Zetterstedt, 1860)	B	M S
<i>sanctimarci</i> (Czerny, 1906)	M S	= <i>angustitarsis</i> Malloch, 1920		
<i>seneciella</i> (Mead, 1892)	B M S	<i>uniseriata</i> (Stein, 1914)	B	M S
<i>sericea</i> (Malloch, 1920)	M S	Egle Robineau-Desvoidy, 1830		
<i>stylatica</i> (Robineau-Desvoidy, 1830)	B	<i>ciliata</i> (Walker, 1840)	B	M S
<i>spinosa</i> (Rondani, 1866)	B	<i>concomitans</i> (Pandellé, 1900)	B	M S
<i>striolata</i> (Fallén, 1824)	B M S	<i>minuta</i> (Meigen, 1826)	B	M S
<i>varicolor</i> (Meigen, 1826)	B M S	<i>parva</i> Robineau-Desvoidy, 1830	B	M
<i>verticella</i> (Zetterstedt, 1838)	B	<i>rhinometra</i> (Pandellé, 1900)	B	
Calythea Schnabl & Dziedzicki, 1911		Emmesomyia Malloch, 1917		
<i>nigricans</i> (Robineau-Desvoidy, 1830)	B M S	<i>grisea</i> (Robineau-Desvoidy, 1830)	B	M S
<i>pratincola</i> (Panzer, 1809)	B M S	= <i>socia</i> auct. nec Fallén, 1825		
Chiastochaeta Pokorný, 1889		<i>socia</i> (Fallén, 1825)		M S
<i>lophota</i> Karl, 1943	M	Enneastigma Stein, 1916		
= <i>trollii</i> auct. nec Zetterstedt, 1845		<i>compressum</i> (Stein, 1916)		S
Chirosia Rondani, 1856		Eustalomyia Kowarz, 1873		
<i>albitarsis</i> (Zetterstedt, 1845)	B M S	<i>festiva</i> (Zetterstedt, 1845)	B	M S
<i>bendleri</i> (Ringdahl, 1935)		<i>hilaris</i> (Fallén, 1823)	B	M S
<i>cepelaki</i> Teschner, 1978		<i>histrion</i> (Zetterstedt, 1838)	B	M S
<i>cinerosa</i> (Zetterstedt, 1845)	B M S	<i>vittipes</i> (Zetterstedt, 1845)		M
<i>flavipennis</i> (Fallén, 1823)	B M	Eutrichota Kowarz, 1893		
<i>grossicauda</i> Strobl, 1899	B M S	= <i>Parapegomyia</i> Griffiths, 1984		
<i>histrion</i> (Rondani, 1866)	B M S	<i>inornata</i> (Loew, 1873)		M S
<i>latipennis</i> (Zetterstedt, 1838)	B M S	<i>longimana</i> (Pokorný, 1887)	B	M S
<i>similata</i> (Tienstuu, 1839)	M	<i>pilimana</i> (Ringdahl, 1918)		M
Delia Robineau-Desvoidy, 1830		<i>praeclara</i> (Stein, 1914)		S
<i>albula</i> (Fallén, 1825)	B	<i>praepotens</i> (Wiedemann, 1817)	B	M S
<i>antiqua</i> (Meigen, 1826)	B M S	<i>schineri</i> (Schnabl in Schnabl		
<i>bracteata</i> (Rondani, 1866)		& Dziedzicki, 1911)	B	M S
<i>brunnescens</i> (Zetterstedt, 1845)	B M S	<i>soculata</i> (Zetterstedt, 1845)	B	
<i>cardui</i> (Meigen, 1826)	B M S	Heterostylodes Hennig, 1967		
<i>carduiformis</i> Schnabl in Schnabl		<i>macrura</i> (Schnabl, 1911)		S
& Dziedzicki, 1911		<i>nominabilis</i> (Collin, 1947)	B	
<i>coarctata</i> (Fallén, 1825)	B M S	<i>obscura</i> (Macquart, 1835)	B	M S
<i>coronariae</i> (Hendel, 1925)	M S	= <i>denigrata</i> (Zetterstedt, 1845)		
<i>crumventris</i> (Zetterstedt, 1860)	M S	= <i>pilifera</i> (Zetterstedt, 1845)		
<i>echinata</i> (Séguy, 1923)	B M S	= <i>pratensis</i> auct. nec Meigen, 1826		
<i>fallax</i> (Loew, 1873)		<i>pratensis</i> (Meigen, 1826)	B	S
<i>floralis</i> (Fallén, 1824)	B M S	Hydrophoria Robineau-Desvoidy, 1830		
<i>floricola</i> Robineau-Desvoidy, 1830	B M S	<i>albiceps</i> (Meigen, 1826)	B	M S
<i>florilega</i> (Zetterstedt, 1845)	B M S	<i>lancifer</i> (Harris, 1780)	B	M S
<i>frontella</i> (Zetterstedt, 1838)	B M S	= <i>virginea</i> (Meigen, 1826)		
<i>interflua</i> (Pandellé, 1900)	M S	<i>linogrisea</i> (Meigen, 1826)	B	M S
<i>lamelliseta</i> (Stein, 1900)	B	<i>ruralis</i> (Meigen, 1826)	B	M S
<i>linearis</i> (Stein, 1898)	B M S	<i>silvicola</i> (Robineau-Desvoidy, 1830)	B	M S
<i>lineariventris</i> (Zetterstedt, 1845)	B M	= <i>annulata</i> auct. nec Pandellé, 1899		
<i>longicauda</i> (Strobl, 1898)	B M S	Hylemya Robineau-Desvoidy, 1830		
<i>majuscula</i> (Pokorný, 1889)	S	<i>nigrimana</i> (Meigen, 1826)	B	M S

<i>urbica</i> van der Wulp, 1896	B M S	<i>meridiana</i> (Villeneuve, 1923)	B M S
= <i>latifrons</i> (Schnabl in Schnabl & Dziedzicki, 1911)		<i>nigrisquama</i> (Stein, 1888)	B M S
<i>vagans</i> (Panzer, 1798)	B M S	<i>notabilis</i> (Zetterstedt, 1846)	B M S
<i>variata</i> (Fallén, 1823)	B M S	<i>pallidoscutellata</i> (Zetterstedt, 1852)	B S
Hylemyza Schnabl & Dziedzicki, 1911		<i>pulchripes</i> (Loew, 1857)	B M
<i>partita</i> (Meigen, 1826)	B M S	<i>rubivora</i> (Coquillett, 1897)	B M S
Hyporites Pokorný, 1893		<i>rufina</i> (Fallén, 1825)	B M S
<i>montanus</i> (Schiner, 1862)	B M S	<i>rugulosa</i> (Zetterstedt, 1845)	M
Lasiomma Stein, 1916		<i>seitenstettensis</i> (Strobl, 1880)	B M S
<i>anthomyinum</i> (Rondani, 1866)	S	<i>setaria</i> (Meigen, 1826)	B M S
<i>cuneicorne</i> (Zetterstedt, 1838)	M	<i>solenis</i> (Meigen, 1826)	B M S
<i>octomaculatum</i> (Ringdahl, 1937)	B	<i>steini</i> (Hendel, 1925)	B M S
<i>picipes</i> (Meigen, 1826)	B M	<i>testacea</i> (De Geer, 1776)	B M S
<i>semitidum</i> (Zetterstedt, 1845)	B M S	<i>transversa</i> (Fallén, 1825)	B M S
<i>strigilatum</i> (Zetterstedt, 1838)	B M S	<i>ulmaria</i> (Rondani, 1866)	B S
Leucophora Robineau-Desvoidy, 1830		<i>vanduzeei</i> Malloch, 1919	B M
<i>cinerea</i> Robineau-Desvoidy, 1830	B M S	<i>vittigera</i> (Zetterstedt, 1838)	B S
<i>grisella</i> Hennig, 1967	B M	<i>winthemi</i> (Meigen, 1826)	B M S
<i>obtusata</i> (Zetterstedt, 1838)	B S	<i>zonata</i> (Zetterstedt, 1838)	B M
<i>personata</i> (Collin, 1922)	B M S	Pegoplata Schnabl & Dziedzicki, 1911	
<i>sericea</i> Robineau-Desvoidy, 1830	B M S	<i>aestiva</i> (Meigen, 1826)	B M S
<i>sociata</i> (Meigen, 1826)	B	<i>debilis</i> (Stein, 1916)	B S
<i>sponsa</i> (Meigen, 1826)	B S	<i>infirma</i> (Meigen, 1826)	B M S
<i>unistriata</i> (Zetterstedt, 1838)	B M S	<i>juvenilis nitidicauda</i> Schnabl, 1911	B M S
Mycophaga Rondani, 1856		= <i>virginea</i> auct. nec Meigen, 1826	
<i>testacea</i> (Gimmerthal, 1834)	B M S	<i>nigroscutellata</i> (Stein, 1920)	B
Myopina Robineau-Desvoidy, 1830		<i>palposa</i> (Stein, 1897)	M S
<i>myopina</i> (Fallén, 1824)	B M S	<i>patellans</i> (Pandellé, 1900)	M
Paradelia Ringdahl, 1933		Phorbia Robineau-Desvoidy, 1830	
<i>intersecta</i> (Meigen, 1826)	B M S	<i>atrogrisea</i> Tiensuu, 1936	B M S
<i>lunatifrons</i> (Zetterstedt, 1845)	B M S	<i>bartaki</i> Ackland & Michelsen, 1987	B M S
Paregle Schnabl, 1911		<i>curvicauda</i> (Zetterstedt, 1845)	B M S
<i>audacula</i> (Harris, 1780)	B M S	<i>fascicularis</i> Tiensuu, 1936	B M S
<i>pilipes</i> (Stein, 1916)	B	<i>fumigata</i> (Meigen, 1826)	B M S
<i>vetula</i> (Zetterstedt, 1838)	B M S	<i>genitalis</i> (Schnabl, 1911)	B M S
Pegomya Robineau-Desvoidy, 1830		<i>haberlandti</i> (Schiner, 1865)	B M S
<i>argyrocephala</i> (Meigen, 1826)	B M	<i>hadyensis</i> Ackland, 1993	M
<i>betae</i> (Curtis, 1847)	B M S	<i>kulai</i> Ackland, 1993	B
<i>bicolor</i> (Wiedemann, 1817)	B M S	<i>longipilis</i> (Pandellé, 1900)	B M
<i>caesia</i> (Stein, 1906)	B M S	<i>melania</i> Ackland & Michelsen, 1987	B
<i>calypttrata</i> (Zetterstedt, 1846)	B M	<i>moliniaris</i> (Karl, 1917)	B M S
<i>conformis</i> (Fallén, 1825)	B M S	<i>nuceicornis</i> (Pandellé, 1900)	M
<i>cunicularia</i> (Rondani, 1866)	B	<i>penicillaris</i> (Stein, 1916)	B M S
<i>deprimata</i> (Zetterstedt, 1845)	M	<i>ponti</i> Hennig, 1969	M
<i>flavifrons</i> (Walker, 1849)	B M	<i>sepia</i> (Meigen, 1826)	B M S
<i>flavoscutellata</i> (Zetterstedt, 1838)	B	<i>unipila</i> (Karl, 1917)	B M S
<i>fulgens</i> (Meigen, 1826)	B M S	Strobilomyia Michelsen, 1988	
<i>geniculata</i> (Bouché, 1834)	B M S	<i>anthracina</i> (Czerny, 1906)	B M
<i>haemorrhoum</i> (Zetterstedt, 1838)	B M S	<i>laricicola</i> (Karl, 1928)	M
<i>holmgreni</i> (Boheman, 1859)	S	Subhylemyia Ringdahl, 1933	
<i>holostaeae</i> (Hering, 1924)	M	<i>longula</i> (Fallén, 1824)	B M S
<i>hyoscyami</i> (Panzer, 1809)	B M S	Zaphne Robineau-Desvoidy, 1830	
<i>incisiva</i> Stein, 1906	B	<i>ambigua</i> (Fallén, 1823)	B M
<i>interruptella</i> (Zetterstedt, 1855)	B	<i>caudata</i> (Zetterstedt, 1855)	M
<i>laticornis</i> (Fallén, 1825)	M	<i>divisa</i> (Meigen, 1826)	B M S
<i>maculata</i> (Stein, 1906)	M S	<i>imuncta</i> (Zetterstedt, 1838)	S
		<i>wierzejskii</i> (Mik, 1867)	B M

FANNIIDAE

František Gregor & Rudolf Rozkošný

Piezura Rondani, 1866*boletorum* (Rondani, 1866) B M S*graminicola* (Zetterstedt, 1846) B M S*Fannia* Robineau-Desvoidy, 1830*aequilineata* Ringdahl, 1945 B M S*armata* (Meigen, 1826) B M S*atra* (Stein, 1895) B M S*barbata* (Stein, 1892) M S*canicularis* (Linnaeus, 1761) B M S*carbonaria* (Meigen, 1826) B M S*carbonella* (Stein, 1895) B M S*clara* Collin, 1939 B*coracina* (Loew, 1873) B M S*corvina* (Verrall, 1892) B M*cothurnata* (Loew, 1873) M S*difficilis* (Stein, 1895) B M S*fasciculata* (Loew, 1873) M*fuscitibia* Stein, 1920 M*= coracula* Collin, 1958*fuscula* (Fallén, 1825) B M S*genualis* (Stein, 1895) B M*glaucescens* (Zetterstedt, 1845) B M S*hirticeps* (Stein, 1892) B M S*immutica* Collin, 1939 B M*incisurata* (Zetterstedt, 1838) B M S*krimensis* Ringdahl, 1934 M S*latipalpis* (Stein, 1892) B M S*lepida* (Wiedemann, 1817) B M S*leucosticta* (Meigen, 1838) B M S*limbata* (Tiensuu, 1938) B*= herolinensis* Hennig, 1955*lineata* (Stein, 1895) B M*lustrator* (Harris, 1780) B M S*manicata* (Meigen, 1826) B M S*melania* (Dufour, 1839) S*metallipennis* (Zetterstedt, 1838) B M S*minutipalpis* (Stein, 1895) B M S*mollissima* (Haliday in Westwood, 1840) B M S*monilis* (Haliday, 1838) B M S*nidica* Collin, 1939 B*nigra* Malloch, 1910 B M*ornata* (Meigen, 1826) B M S*pallitibia* (Rondani, 1866) B M S*parva* (Stein, 1895) B M S*pauli* Pont in Rozkošný, Gregor

& Pont, 1997 B M

= nitida (Stein, 1895) nec Macquart, 1835*polychaeta* (Stein, 1895) B M S*postica* (Stein, 1895) B M S*posticata* (Meigen, 1826) B M S*pruinosa* (Meigen, 1826) B M S*pseudonorvegica* d'Assis-Fonseca, 1966 B*ringdahlana* Collin, 1939 B M S*rondanii* (Strobl, 1893) B M S*scalaris* (Fabricius, 1794) B M S*serena* (Fallén, 1825) B M S*similis* (Stein, 1895) B M S*sociella* (Zetterstedt, 1845) B M S*= spinosa* Karl, 1928*speciosa* (Villeneuve, 1898) B*subpubescens* Collin, 1958 B M S*subsimilis* Ringdahl, 1934 M S*tuberculata* (Zetterstedt, 1849) B M*umbratica* Collin, 1939 B M S*umbrosa* (Stein, 1895) B M S*verrallii* (Stein, 1895) M*vesparia* (Meade, 1891) B M S

MUSCIDAE

František Gregor

ACHANTHIPTERINAE

Acanthiptera Rondani, 1856*rohrelliformis* (Robineau-Desvoidy, 1830) B M

MUSCINAE

REINWARDTIINI

Muscina Robineau-Desvoidy, 1830*angustifrons* (Loew, 1858) B*levida* (Harris, 1780) B M S*pascuorum* (Meigen, 1826) B M S*prolapsa* (Harris, 1780) B M S*stabulans* (Fallén, 1817) B M S

AZELIINI

Azelia Robineau-Desvoidy, 1830*aterrima* (Meigen, 1826) B M*cilipes* (Haliday, 1838) B M S*gibbera* (Meigen, 1826) B S*monodactyla* Loew, 1874 B M S*nebulosa* Robineau-Desvoidy, 1830 B M S*parva* Rondani, 1868 M S*triquetra* (Wiedemann, 1817) B M S*zetterstedti* Rondani, 1866 B M S*Thricops* Rondani, 1866*aculeipes* (Zetterstedt, 1838) S*beckeri* (Pokorny, 1893) B M S*culminum* (Pokorny, 1889) S*cunctans* (Meigen, 1826) B M S*diaphanus* (Wiedemann, 1817) B M S*genarum* (Zetterstedt, 1838) B M S*innocuus* (Zetterstedt, 1838) B S*lividiventris* (Zetterstedt, 1845) B M S*longipes* (Zetterstedt, 1845) B M S*nigrifrons* (Robineau-Desvoidy, 1830) B M S*nigritellus* (Zetterstedt, 1838) B M S*rostratus* (Meade, 1882) B S*semicinereus* (Wiedemann, 1817) B M S*simplex* (Wiedemann, 1817) B M S

<i>sudeticus</i> (Schnabl, 1888)	B M S	<i>Neomyia</i> Walker, 1859	
<i>tatricus</i> Gregor, 1988	S	<i>cornicina</i> (Fabricius, 1781)	B M S
Drymeia Meigen, 1826		<i>viridescens</i> (Robineau-Desvoidy, 1830)	B M S
<i>alpicola</i> (Rondani, 1871)	S	Pyrellia Robineau-Desvoidy, 1830	
<i>hamata</i> (Fallén, 1823)	B M S	<i>rapax</i> (Harris, 1780)	M S
<i>tetra</i> (Meigen, 1826)	B S	<i>vivida</i> Robineau-Desvoidy, 1830	B M S
<i>vicana</i> (Harris, 1780)	B M S	Eudasyphora Townsend, 1911	
Hydrotaea Robineau-Desvoidy, 1830		<i>cyanella</i> (Meigen, 1826)	B S
<i>aenescens</i> (Wiedemann, 1830)	B M S	<i>cyanicolor</i> (Zetterstedt, 1845)	B M S
<i>albipuncta</i> (Zetterstedt, 1845)	B M S	<i>zimini</i> (Hennig, 1963)	S
<i>armipes</i> (Fallén, 1825)	B M S	Dasyphora Robineau-Desvoidy, 1830	
<i>borussica</i> Stein, 1899	B M S	<i>albofasciata</i> (Macquart, 1839)	M S
<i>capensis</i> (Wiedemann, 1818)	B M S	<i>penicilata</i> (Egger, 1865)	B M S
<i>cinerea</i> Robineau-Desvoidy, 1830	B	<i>pratorum</i> (Meigen, 1826)	B M S
<i>cyrtoneurina</i> (Zetterstedt, 1845)	B M S		
<i>dentipes</i> (Fabricius, 1805)	B M S	STOMOXYINI	
<i>diabolus</i> (Harris, 1780)	B M S	Stomoxys Geoffroy, 1762	
<i>floccosa</i> Macquart, 1835	B M S	<i>calcitrans</i> (Linnaeus, 1758)	B M S
<i>glabricula</i> (Fallén, 1825)	S	Haematobia Le Peletier & Serville, 1828	
<i>ignava</i> (Harris, 1780)	B M S	<i>irritans</i> (Linnaeus, 1758)	B M S
<i>irritans</i> (Fallén, 1823)	B M S	Haematobosca Bezzi, 1907	
<i>meridionalis</i> Portschinsky, 1882	B M S	<i>atripalpis</i> (Bezzi, 1895)	S
<i>meteorica</i> (Linnaeus, 1758)	B M S	<i>stimulans</i> (Meigen, 1824)	B M S
<i>militaris</i> (Meigen, 1826)	S		
<i>nidicola</i> Malloch, 1925	M	PHAONIINAE	
<i>palaestrica</i> (Meigen, 1826)	B M S	EGINIINI	
<i>pandellei</i> Stein, 1899	B M S	Eginia Robineau-Desvoidy, 1830	
<i>pellucens</i> Portschinsky, 1879	B M S	<i>ocypterata</i> (Meigen, 1826)	M S
<i>pillipes</i> Stein, 1903	B M S		
<i>piliptibia</i> Stein, 1916	S	ATHERIGONINI	
<i>similis</i> Meade, 1887	B M S	Atherigona Rondani, 1856	
<i>tuberculata</i> Rondani, 1866	B M S	<i>varia</i> (Meigen, 1826)	M
<i>velutina</i> Robineau-Desvoidy, 1830	B M S		
Potamia Robineau-Desvoidy		PHAONIINI	
<i>littoralis</i> Robineau-Desvoidy, 1830	B M S	Phaonia Robineau-Desvoidy, 1830	
		<i>aeneiventris</i> (Zetterstedt, 1845)	B M
MUSCINI		<i>alpicola</i> (Zetterstedt, 1845)	S
Mesembrina Meigen, 1826		<i>amicula</i> Villeneuve, 1922	M
<i>intermedia</i> Zetterstedt, 1848	S	<i>angelicae</i> (Scopoli, 1763)	B M S
<i>meridiana</i> (Linnaeus, 1758)	B M S	<i>angulicornis</i> (Zetterstedt, 1838)	B
<i>mystacea</i> (Linnaeus, 1758)	B M S	<i>apicalis</i> Stein, 1914	B M
<i>resplendens</i> Wahlberg, 1844	S	<i>atriceps</i> (Loew, 1858)	B
Polietes Rondani, 1866		<i>bitincta</i> (Rondani, 1866)	B M
<i>domitor</i> (Harris, 1780)	B M S	<i>boleticola</i> (Rondani, 1866)	B S
<i>lardarius</i> (Fabricius, 1781)	B M S	<i>candicans</i> (Pandellé, 1898)	S
Musca Linnaeus, 1758		<i>canescens</i> Stein, 1916	M
<i>autumnalis</i> De Geer, 1776	B M S	<i>cincta</i> (Zetterstedt, 1846)	B M
<i>domestica</i> Linnaeus, 1758	B M S	<i>consobrina</i> (Zetterstedt, 1838)	B M S
<i>larvipara</i> Portschinsky, 1910	S	<i>czernyi</i> Hennig, 1963	B M S
<i>osiris</i> Wiedemann, 1830	B M S	<i>disjuncta</i> Stein, 1916	B S
<i>tempestiva</i> Fallén, 1817	B M S	<i>errans</i> (Meigen, 1826)	B M S
Morellia Robineau-Desvoidy, 1830		<i>erronea</i> (Schnabl, 1887)	B M S
<i>aenescens</i> Robineau-Desvoidy, 1830	B M S	<i>exoleta</i> (Meigen, 1826)	B
<i>asetosa</i> Baranov, 1925	B S	<i>falleni</i> Michelsen, 1977	B M S
<i>hortorum</i> (Fallén, 1817)	B M S	<i>fuscata</i> (Fallén, 1825)	B M S
<i>podagrica</i> (Loew, 1857)	B M S	<i>goberti</i> (Mik, 1881)	B M S
<i>simplex</i> (Loew, 1857)	B M S	<i>gracilis</i> Stein, 1916	B M S

<i>halterata</i> (Stein, 1893)	B M S	<i>lasiophthalma</i> (Macquart, 1835)	B M S
<i>hybrida</i> (Schnabl, 1881)	B M S	<i>latitarsis</i> Ringdahl, 1924	B M S
<i>incana</i> (Wiedemann, 1817)	B M S	<i>laxifrons</i> (Zetterstedt, 1860)	M S
<i>jugorum</i> (Strobl, 1910)	B M S	<i>maculipennis</i> (Zetterstedt, 1845)	M S
<i>laeta</i> (Fallén, 1823)	M S	<i>moedlingensis</i> (Schnabl in Schnabl & Dziedzicki, 1911)	B M S
<i>liliputa</i> Zinovev, 1990	M	<i>obscurata</i> (Meigen, 1826)	B M S
<i>lugubris</i> (Meigen, 1826)	B M S	<i>obscuratoides</i> (Schnabl, 1887)	B S
<i>magnicornis</i> (Zetterstedt, 1845)	B M S	<i>obtusipennis</i> (Fallén, 1923)	S
<i>mario</i> (Zetterstedt, 1845)	B M S	<i>pertusa</i> (Meigen, 1826)	B M S
<i>mystica</i> (Meigen, 1826)	B M	<i>protuberans</i> (Zetterstedt, 1845)	S
<i>nymphaeorum</i> (Robineau-Desvoidy, 1830)	B M S	<i>pubiseta</i> (Zetterstedt, 1845)	B M S
<i>pallida</i> (Fabricius, 1787)	B M S	<i>quadrinotata</i> (Meigen, 1826)	B S
<i>palpata</i> (Stein, 1897)	B M S	<i>quadrum</i> (Fabricius, 1805)	B M S
<i>perdita</i> (Meigen, 1830)	B M S	<i>reversio</i> (Harris, 1780)	B M S
<i>pratensis</i> (Robineau-Desvoidy, 1830)	B M S	<i>setiventris</i> Ringdahl, 1924	B M S
<i>pullata</i> (Czerny, 1900)	B S	<i>sexmaculata</i> (Preyssler, 1791)	B M S
<i>pura</i> (Loew, 1873)	M S	<i>subvittata</i> (Séguy, 1923)	B M S
<i>rufipalpis</i> (Macquart, 1835)	S	<i>tetrastigma</i> (Meigen, 1826)	B M S
<i>rufiventris</i> (Scopoli, 1763)	B M S	<i>trivittata</i> (Zetterstedt, 1860)	B M S
<i>serva</i> (Meigen, 1826)	B M S	<i>veterana</i> (Zetterstedt, 1838)	B
<i>siebecki</i> Schnabl in Schnabl & Dziedzicki, 1911	B M S	<i>vicina</i> (Czerny, 1900)	S
<i>subventa</i> (Harris, 1780)	B M S	Brontaea Kowarz, 1873	
<i>tefti</i> (Schnabl, 1888)	B M S	<i>tonitru</i> (Wiedemann, 1824)	S
<i>trimaculata</i> (Bouché, 1834)	B M S		
<i>tuguriorum</i> (Scopoli, 1763)	B M S	MYDAEINAE	
<i>valida</i> (Harris, 1780)	B M S	Mydaea Robineau-Desvoidy, 1830	
<i>vivida</i> (Rondani, 1871)	B	<i>ancilla</i> (Meigen, 1826)	B M S
<i>wahlbergi</i> Ringdahl, 1930	B S	<i>anicula</i> (Zetterstedt, 1860)	B M S
<i>zugmayeriae</i> (Schnabl, 1888)	B M S	<i>corni</i> (Scopoli, 1763)	B M S
Lophosceles Ringdahl, 1922		<i>deserta</i> (Zetterstedt, 1845)	B M S
<i>cinereiventris</i> (Zetterstedt, 1845)	B M S	<i>electa</i> (Zetterstedt, 1860)	B M S
<i>frenatus</i> (Holmgren, 1872)	S	<i>humeralis</i> Robineau-Desvoidy, 1830	B M S
<i>mutatus</i> (Fallén, 1825)	S	<i>maculiventris</i> (Zetterstedt, 1846)	B
Helina Robineau-Desvoidy, 1830		<i>nebulosa</i> (Stein, 1893)	B M S
<i>allotata</i> (Meigen, 1830)	B M	<i>nubila</i> Stein, 1919	M S
<i>annosa</i> (Zetterstedt, 1838)	B M S	<i>orthonevra</i> (Macquart, 1835)	M
<i>areolata</i> Collin, 1953	B M	<i>setifemur</i> Ringdahl, 1924	B M S
<i>atricolor</i> (Fallén, 1825)	B M S	<i>urbana</i> (Meigen, 1826)	B M S
<i>balsaci</i> (Séguy, 1946)	B	Myospila Rondani, 1856	
<i>calceata</i> (Rondani, 1866)	B M S	<i>alpina</i> Hendel, 1901	S
<i>calceataeformis</i> (Schnabl in Schnabl & Dziedzicki, 1911)	B M S	<i>bimaculata</i> (Macquart, 1834)	B M S
<i>celsa</i> (Harris, 1780)	B M S	<i>meditabunda</i> (Fabricius, 1781)	B M S
<i>ciliatocosta</i> (Zetterstedt, 1845)	B M S	Hebecnema Schnabl, 1889	
<i>cilipes</i> (Schnabl, 1902)	S	<i>fumosa</i> (Meigen, 1826)	M S
<i>concolor</i> (Czerny, 1900)	M	<i>nigra</i> (Robineau-Desvoidy, 1830)	B M S
<i>confinis</i> (Fallén, 1825)	B M S	<i>nigricolor</i> (Fallén, 1825)	B M S
<i>consimilis</i> (Fallén, 1825)	B	<i>umbratica</i> (Meigen, 1826)	B M S
<i>cothurnata</i> (Rondani, 1866)	B M S	<i>vespertina</i> (Fallén, 1823)	B M S
<i>decipiens</i> Mihályi, 1974	B M S	Graphomya Robineau-Desvoidy, 1830	
<i>deleta</i> (Stein, 1914)	B M S	<i>maculata</i> (Scopoli, 1763)	B M S
<i>depuncta</i> (Fallén, 1825)	B M S		
<i>evecta</i> (Harris, 1780)	B M S	COENOSIINAE	
<i>fratercula</i> (Zetterstedt, 1845)	B M S	LIMNOPHORINI	
<i>impuncta</i> (Fallén, 1825)	B M S	Spilogona Schnabl, 1911	
		<i>acrostichalis</i> (Stein, 1916)	B
		<i>atricans</i> (Pandellé, 1899)	B M S

<i>brunneisquama</i> (Zetterstedt, 1845)	B	S	<i>Schoenomyza</i> Haliday, 1833		
<i>carbonella</i> (Zetterstedt, 1845)	B	M	<i>litorella</i> (Fallén, 1823)	B	M
<i>contractifrons</i> (Zetterstedt, 1838)	B	M	<i>Macrororchis</i> Rondani, 1877		
<i>denigrata</i> (Meigen, 1826)	B	M	<i>meditata</i> (Fallén, 1825)	B	M
<i>dispar</i> (Fallén, 1823)	B	M	<i>Coenosia</i> Meigen, 1826		
<i>leucogaster</i> (Zetterstedt, 1838)	B		= <i>Caricea</i> Robineau-Desvoidy, 1830		
<i>litorea</i> (Fallén, 1823)	B	M	<i>acuminata</i> Strobl, 1898	B	
<i>meadei</i> (Schnabl in Becker & al., 1915)		M	<i>agromyzina</i> (Fallén, 1825)	B	M
<i>pacifica</i> (Meigen, 1826)	B	M	<i>albicornis</i> Meigen, 1826	B	
<i>scutulata</i> (Schnabl in Schnabl			<i>ambulans</i> Meigen, 1826	B	
& Dziedzicki, 1911)	B	S	<i>antennata</i> (Zetterstedt, 1849)		S
<i>solitaria</i> (Collin, 1921)	B	M	<i>atra</i> Meigen, 1830	B	M
<i>surda</i> (Zetterstedt, 1845)		M	<i>bilineella</i> (Zetterstedt, 1838)		M
<i>Limnophora</i> Robineau-Desvoidy, 1830			<i>campestris</i> (Robineau-Desvoidy, 1830)	B	M
<i>exuta</i> (Kowarz, 1893)	B	M	<i>distinguens</i> Collin, 1930	B	M
<i>latevittata</i> Schnabl in Schnabl			<i>dubiosa</i> Hennig, 1961		S
& Dziedzicki, 1911	B	M	<i>femoralis</i> (Robineau-Desvoidy, 1830)	B	M
<i>maculosa</i> (Meigen, 1826)	B	M	<i>humilis</i> Meigen, 1826	B	M
<i>nigripes</i> (Robineau-Desvoidy, 1830)	B		<i>infantula</i> Rondani, 1866	B	M
<i>pollinifrons</i> Stein, 1916	B	M	<i>intermedia</i> (Fallén, 1825)	B	M
<i>pulchriceps</i> (Loew, 1860)	B	M	<i>means</i> Meigen, 1826	B	M
<i>riparia</i> (Fallén, 1824)	B	M	<i>mollicula</i> (Fallén, 1825)	B	M
<i>tigrina</i> (Am Stein, 1860)	B	M	<i>nigridigita</i> Rondani, 1866	B	M
<i>triangula</i> (Fallén, 1825)	B	M	<i>obscuricula</i> (Rondani, 1871)		S
<i>Lispe</i> Latreille, 1796			<i>octopunctata</i> (Zetterstedt, 1838)	B	
<i>caesia</i> Meigen, 1826	B	M	<i>octosignata</i> Rondani, 1866	B	
<i>consanguinea</i> Loew, 1858	B	M	<i>paludis</i> Tiensuu, 1939	B	
<i>flavicincta</i> Loew, 1847	B	S	<i>pedella</i> (Fallén, 1825)	B	M
<i>loewi</i> Ringdahl, 1922	B		<i>perpusilla</i> Meigen, 1826	B	M
<i>longicollis</i> Meigen, 1826		M	<i>pumila</i> (Fallén, 1825)	B	M
<i>melaleuca</i> Loew, 1847	B	M	<i>rufipalpis</i> Meigen, 1826	B	M
<i>nana</i> Macquart, 1835		M	<i>sempustulata</i> Rondani, 1866	B	S
<i>pygmaea</i> Fallén, 1825	B	M	<i>styriaca</i> Hennig, 1961		S
<i>superciliosa</i> Loew, 1861		M	<i>testacea</i> (Robineau-Desvoidy, 1830)	B	M
<i>tentaculata</i> (De Geer, 1776)	B	M	<i>tigrina</i> (Fabricius, 1775)	B	M
<i>uliginosa</i> Fallén, 1825	B	M	<i>trilineella</i> (Zetterstedt, 1838)	B	M
			<i>verralli</i> Collin, 1953	B	M
COENOSIINI					
<i>Pseudocoenosia</i> Stein, 1916			CALLIPHORIDAE		
<i>abnormis</i> Stein, 1916	B		František Gregor & Rudolf Rozkošný		
<i>solitaria</i> (Zetterstedt, 1838)	B	M			
<i>Limnospila</i> Schnabl, 1902			CALLIPHORINAE		
<i>albifrons</i> (Zetterstedt, 1849)	B	M	<i>Bellardia</i> Robineau-Desvoidy, 1863		
<i>Lispocephala</i> Pokorný, 1893			<i>bayeri</i> (Jacentkovský, 1937)		M
<i>alma</i> (Meigen, 1826)	B	M	<i>obsoleta</i> (Meigen, 1824)		S
<i>brachialis</i> (Rondani, 1877)	B	M	<i>pandia</i> (Walker, 1849)		M
<i>erythrocerata</i> (Robineau-Desvoidy,			<i>polita</i> (Mik, 1884)		M
1830)	B	M	<i>stricta</i> (Villeneuve, 1926)	B	M
<i>falculata</i> Collin, 1963	B		<i>vespillo</i> (Fabricius, 1794)		
<i>pallipalpis</i> (Zetterstedt, 1845)	B	M	= <i>borotinensis</i> (Jacentkovský, 1944)		
<i>spuria</i> (Zetterstedt, 1838)	B	M	<i>viarum</i> (Robineau-Desvoidy, 1830)	B	M
<i>ungulata</i> (Rondani, 1866)		M	= <i>pusilla</i> (Meigen, 1826) nec Gmelin, 1790		
<i>verna</i> (Fabricius, 1794)	B	M	<i>vulgaris</i> (Robineau-Desvoidy, 1830)	B	M
<i>Orchisia</i> Rondani, 1878			= <i>agilis</i> (Meigen, 1826) nec Harris, 1780		
<i>costata</i> (Meigen, 1826)	B	M	<i>Calliphora</i> Robineau-Desvoidy, 1830		
<i>Spanochaeta</i> Stein, 1919			<i>loewi</i> Enderlein, 1903	B	M
<i>dorsalis</i> (von Roser, 1840)	B	M			

CHECK LIST OF DIPTERA / KAROLINUM - CHARLES UNIVERSITY PRESS, PRAGUE 1997

SARCOPHAGIDAE			OEBALIINI		
Dalibor Povolný			<i>Oebalia</i> Robineau-Desvoidy, 1863		
MACRONYCHIINAE			<i>cyllindrica</i> (Fallén, 1810)	B	M
<i>Macronychia</i> Rondani, 1859			<i>Ptychoneura</i> Brauer & Bergenstamm, 1889		
sg. <i>Macronychia</i> s. str.			<i>minuta</i> (Fallén, 1810)	M	S
<i>lemartei</i> Jacentkovský, 1941	M	S	METOPINI		
<i>strigilnervis</i> (Zetterstedt, 1844)	M	S	<i>Hilarella</i> Rondani, 1856		
sg. <i>Moschusa</i> Robineau-Desvoidy, 1863			<i>hilarella</i> (Zetterstedt, 1844)	B	M S
<i>agrestis</i> (Fallén, 1810)	B	M S	<i>stricta</i> (Meigen, 1830)	B	M S
<i>alpestris</i> Rondani, 1865	B	M S	= <i>siphonina</i> (Zetterstedt, 1844)		
<i>dumosa</i> (Pandellé, 1895)			<i>Mesomelaena</i> Rondani, 1859		
<i>griseola</i> (Fallén, 1820)	M	S	<i>mesomelaena</i> (Loew, 1848)	M	S
<i>polyodon</i> (Meigen, 1824)	B	M S	<i>Metopia</i> Meigen, 1803		
MILTOGRAMMATINAE			<i>argyrocephala</i> (Meigen, 1824)	B	M S
MILTOGRAMMATINI			<i>campestris</i> (Fallén, 1810)	B	M S
<i>Anaconthothecum</i> Rohdendorf, 1930			<i>roseri</i> Rondani, 1859	B	S
<i>testaceifrons</i> (von Roser, 1840)	B	M S	= <i>rondaniana</i> Venturi, 1953		
<i>Apodacra</i> Macquart, 1854			<i>staegeri</i> Rondani, 1859	M	S
<i>pulchra</i> Egger, 1861	B	S	<i>Phrosinella</i> Robineau-Desvoidy, 1863		
<i>Miltogramma</i> Meigen, 1803			<i>nasuta</i> (Wiedemann, 1824)	M	S
<i>brevipilum</i> Villeneuve, 1911	M	S	<i>Sphenometopa</i> Townsend, 1902		
<i>germari</i> Meigen, 1824	B	M S	= <i>Eumetopiella</i> Verves, 1986 nec Hendel, 1907		
<i>murinum</i> Meigen, 1824	B	M S	<i>fastuosa</i> (Meigen, 1824)	M	S
<i>oestraceum</i> (Fallén, 1820)	B	M S	<i>Taxigramma</i> Perris, 1852		
<i>punctatum</i> Meigen, 1824	B	M S	<i>heteroneura</i> (Meigen, 1830)	B	S
<i>villeneuvei</i> Verves, 1982	M	S	PARAMACRONYCHIINAE		
<i>Miltogrammatidium</i> Rohdendorf, 1930			PARAMACRONYCHIINI		
<i>rutilans</i> (Meigen, 1824)	M	S	<i>Agria</i> Robineau-Desvoidy, 1830		
<i>taeniatum</i> (Meigen, 1824)	M	S	<i>mamillata</i> (Pandellé, 1896)	B	M S
<i>Protomiltogramma</i> Townsend, 1916			<i>monache</i> (Kramer, 1908)	M	S
<i>fasciata</i> (Meigen, 1824)	M	S	<i>punctata</i> Robineau-Desvoidy, 1830	B	M S
<i>Pterella</i> Robineau-Desvoidy, 1863			= <i>affinis</i> (Fallén, 1817) nec Turton, 1800		
<i>grisea</i> (Meigen, 1824)	M	S	<i>Angiometopa</i> Brauer & Bergenstamm, 1889		
<i>melamura</i> (Meigen, 1824)	M	S	<i>falleni</i> Pape, 1986	B	S
<i>penicillaris</i> (Rondani, 1865)	B	M	= <i>ruralis</i> (Fallén, 1817) nec Gravenhorst, 1807		
<i>Senotainia</i> Macquart, 1846			<i>Brachicoma</i> Rondani, 1856		
sg. <i>Senotainia</i> s. str.			<i>devia</i> (Fallén, 1820)	B	M S
<i>conica</i> (Fallén, 1810)	B	M S	<i>Nyctia</i> Robineau-Desvoidy, 1830		
<i>tricuspis</i> (Meigen, 1838)	M	S	<i>halterata</i> (Panzer, 1798)	B	M S
sg. <i>Sphixapata</i> Rondani, 1859			<i>Paramacronychia</i> Brauer &		
<i>albifrons</i> (Rondani, 1859)	B	M S	Bergenstamm, 1889		
<i>puncticornis</i> (Zetterstedt, 1859)	M	S	<i>flavipalpis</i> (Girschner, 1881)	B	M S
AMOBIIINI			<i>Sarcophila</i> Rondani, 1856		
<i>Amobia</i> Robineau-Desvoidy, 1830			<i>latifrons</i> (Fallén, 1817)	B	M S
<i>oculata</i> (Zetterstedt, 1844)	M	S	<i>Wohlfahrtia</i> Brauer & Bergenstamm, 1889		
<i>distorta</i> (Allen, 1926)			<i>magnifica</i> (Schiner, 1862)		S
<i>pelopei</i> (Rondani, 1859)	M		<i>meigeni</i> (Schiner, 1862)	B	M S
<i>signata</i> (Meigen, 1824)	B	M S	SARCOPHAGINAE		
PHYLLOTELINI			PROTODEXIINI		
<i>Metopodia</i> Brauer & Bergenstamm, 1889			<i>Blesoxipha</i> Loew, 1861		
<i>grisea</i> Brauer & Bergenstamm, 1889	M		<i>cochlearis</i> (Pandellé, 1896)	B	M S
<i>Phylloteles</i> Loew, 1844			<i>lapidosa</i> Pape, 1994		M S
<i>pictipennis</i> Loew, 1844	B	M	= <i>campestris</i> auct. nec Robineau-Desvoidy, 1863		
			<i>occatrix</i> (Pandellé, 1896)	M	S

<i>plumicornis</i> (Zetterstedt, 1859)	M S	<i>dissimilis</i> (Meigen, 1826)	B M S
<i>gladiatrix</i> Pandellé, 1896		<i>haemorrhoea</i> (Meigen, 1826)	B M S
<i>ungulata</i> (Pandellé, 1896)	B M	<i>haemorrhoides</i> Boettcher, 1913	M S
Servaisia Robineau-Desvoidy, 1863		<i>hirticrus</i> (Pandellé, 1896)	B M S
<i>erythrura</i> (Meigen, 1826)	B M S	<i>lednicensis</i> Povolný in Povolný	
<i>rossica</i> Villeneuve, 1912	M S	& Verves, 1984	M
Tephromyia Robineau-Desvoidy, 1830		<i>pauciseta</i> (Pandellé, 1896)	S
<i>grisea</i> (Meigen, 1826)	M S	<i>porrecta</i> (Boettcher, 1913)	S
RAVINIINI		<i>proxima</i> (Rondani, 1860)	B M S
Ravinia Robineau-Desvoidy, 1863		<i>rohdendorfi</i> Povolný & Slamečková, 1959	M S
<i>permix</i> (Harris, 1780)	B M S	<i>rohdendorfiana</i> Mihályi, 1975	M S
<i>striata</i> (Fabricius, 1794)		<i>rondaniana</i> (Rohdendorf, 1937)	S
JOHNSONIINI		= <i>depressifrons</i> auct. nec Zetterstedt, 1845	
Sarcotachinella Townsend, 1892		<i>schineri</i> (Bezzi, 1891)	B M S
<i>sinuata</i> (Meigen, 1826)	B M S	<i>slovaca</i> Povolný & Slamečková, 1967	B M S
SARCOPHAGINI		<i>vagans</i> (Meigen, 1826)	B M S
Arachnidomyia Townsend, 1934		<i>vicina</i> (Macquart, 1835)	M S
<i>sempunctata</i> (Fabricius, 1794)	B M S	sg. <i>Pandelleola</i> Rohdendorf, 1937	
<i>clathrata</i> (Meigen, 1826)		<i>filia</i> (Rondani, 1860)	B M S
Ascelotella Enderlein, 1928		Kramerea Rohdendorf, 1937	
<i>granulata</i> (Kramer, 1908)	M S	<i>schuetzei</i> (Kramer, 1909)	B M S
Bellieriomima Rohdendorf, 1937		Krameromyia Verves, 1982	
<i>subulata</i> (Pandellé, 1896)	B M S	<i>anaces</i> (Walker, 1849)	M S
Bercaea Robineau-Desvoidy, 1863		Liopygia Enderlein, 1928	
<i>cruentata</i> (Meigen, 1826)	B M S	sg. <i>Jantia</i> Rohdendorf, 1937	
Discachaeta Enderlein, 1928		<i>crassipalpis</i> (Macquart, 1839)	M S
<i>arcipes</i> (Pandellé, 1896)	B M S	sg. Thomsonia Rohdendorf, 1937	
<i>cucullans</i> (Pandellé, 1896)	S	<i>argyrostoma</i> (Robineau-Desvoidy, 1830)	M S
<i>pumila</i> (Meigen, 1826)	B M S	sg. Varirosellea Hsue, 1979	
Helicophagella Enderlein, 1928		<i>uliginosa</i> (Kramer, 1908)	B S
sg. <i>Helicophagella</i> s.str.		Liosarcophaga Enderlein, 1928	
<i>agnata</i> (Rondani, 1860)	B M S	sg. <i>Curranea</i> Rohdendorf, 1937	
<i>crassimargo</i> (Pandellé, 1896)	B M S	<i>tibialis</i> (Macquart, 1851)	M
<i>maerura</i> (Rohdendorf, 1937)	S	sg. Liosarcophaga s.str.	
<i>novella</i> (Baranov, 1929)	S	<i>aegyptica</i> (Salem, 1935)	S
<i>novercoides</i> auct. nec Boettcher, 1913		<i>emdeni</i> (Rohdendorf, 1969)	B M S
<i>noverca</i> (Rondani, 1860)	B M S	<i>harpax</i> (Pandellé, 1896)	B M S
<i>rosellei</i> (Boettcher, 1912)	B M S	<i>jacobsoni</i> (Rohdendorf, 1937)	S
sg. Parabellieria Verves, 1987		<i>portschinskyi</i> (Rohdendorf, 1937)	M S
<i>melamara</i> (Meigen, 1826)	B M S	<i>tuberosa</i> (Pandellé, 1896)	B M S
Heteronychia Brauer & Bergenstamm, 1889		sg. Pandelleisca Rohdendorf, 1937	
sg. Boettcherella Enderlein, 1927		<i>similis</i> (Meade, 1876)	B M S
<i>munia</i> (Villeneuve, 1912)	S	Pandelleana Rohdendorf, 1937	
sg. Ctenodasyphygia Enderlein, 1928		<i>protuberans</i> (Pandellé, 1896)	M S
<i>minima</i> (Rondani, 1862)	M S	Pierretia Robineau-Desvoidy, 1863	
sg. Heteronychia s.str.		sg. Mehria Enderlein, 1928	
<i>ancilla</i> (Rondani, 1865)	M S	<i>nemoralis</i> (Kramer, 1908)	M S
<i>henaci</i> (Boettcher, 1913)	M S	sg. Pierretia s.str.	
= <i>bezziana</i> auct. nec Boettcher, 1913		<i>discifera</i> Pandellé, 1896	M S
<i>bulgarica</i> (Enderlein, 1936)	B M S	<i>lunigera</i> (Boettcher, 1914)	M S
= <i>boettcheriana</i> (Rohdendorf, 1937)		<i>nigriventris</i> (Meigen, 1826)	B M S
<i>cepelaki</i> Povolný & Slamečková, 1970	M S	<i>socrus</i> (Rondani, 1860)	M S
<i>depressifrons</i> (Zetterstedt, 1845)	B M S	= <i>rostrata</i> (Pandellé, 1896)	
<i>obscurata</i> (Rohdendorf, 1937)		<i>soror</i> (Rondani, 1860)	M S
		<i>villeneuvei</i> Boettcher, 1912	M S
		Parasarcophaga Johnston & Tiggs, 1921	
		<i>albiceps</i> (Meigen, 1826)	B M S

<i>Robineauella</i> Enderlein, 1928			
sg. <i>Digitiventr</i> Fan, 1964			
<i>pseudoscopia</i> (Kramer, 1911)	M	S	
sg. <i>Robineauella</i> s.str.			
<i>caerulescens</i> (Zetterstedt, 1838)	B	M	S
<i>scopia</i> (Pandellé, 1896)			
<i>Rosellea</i> Rohdendorf, 1937			
<i>aratrix</i> (Pandellé, 1896)	B	M	S
<i>Sarcophaga</i> Meigen, 1826			
<i>bachmayeri</i> Lehrer, 1978			S
<i>carriaria</i> (Linnaeus, 1758)	B	M	S
<i>schultzei</i> Müller, 1922			
<i>lastostyla</i> Macquart, 1843	B	M	S
<i>lehmanni</i> Müller, 1922			
<i>moldavica</i> Rohdendorf, 1937		M	S
<i>moravica</i> Povolný, 1986		M	
<i>palavae</i> Povolný, 1993		M	
<i>serbica</i> Baranov, 1930			S
<i>subvicina</i> Rohdendorf, 1937	B	M	S
<i>ukrainica</i> Rohdendorf, 1937			S
<i>variegata</i> (Scopoli, 1763)	B	M	S
<i>carriaria</i> auct. nec Linnaeus, 1758			
<i>zumptiana</i> Lehrer, 1959		M	S
<i>Stackelbergeola</i> Rohdendorf, 1937			
<i>mehadiensis</i> (Boettcher, 1912)		M	
<i>Thyrsochema</i> Enderlein, 1928			
<i>incisilobata</i> Pandellé, 1896	B	M	S
TACHINIDAE			
Juraj Čepelák & Jaromír Vaňhara			
EXORISTINAE			
EXORISTINI			
<i>Bessa</i> Robineau-Desvoidy, 1863			
<i>parallela</i> (Meigen, 1824)	B	M	S
<i>selecta</i> (Meigen, 1824)	B	M	S
<i>Diplostichus</i> Brauer & Bergenstamm, 1889			
<i>janitrix</i> (Hartig, 1838)	B		S
<i>Exorista</i> Meigen, 1803			
<i>civilis</i> (Rondani, 1859)			S
<i>fasciata</i> (Fallén, 1820)	B	M	S
<i>florentina</i> Herting, 1975			S
<i>glossatorum</i> (Rondani, 1859)			S
<i>grandis</i> (Zetterstedt, 1844)	B	M	S
<i>larvarum</i> (Linnaeus, 1758)	B	M	S
<i>mimula</i> (Meigen, 1824)	B	M	S
<i>nympharum</i> (Rondani, 1859)			S
<i>rustica</i> (Fallén, 1810)	B	M	S
<i>tubulosa</i> Herting, 1967	B	M	S
<i>Chetogena</i> Rondani, 1856			
<i>acuminata</i> Rondani, 1859	B	M	S
<i>fasciata</i> (Egger, 1856)		M	S
<i>filipalpis</i> Rondani, 1859	B	M	S
<i>media</i> Rondani, 1859		M	S
<i>obliquata</i> (Fallén, 1810)	B	M	S
<i>Neophryxe</i> Townsend, 1916			
<i>vallina</i> (Rondani, 1861)			S
<i>Parasetigena</i> Brauer & Bergenstamm, 1891			
<i>silvestris</i> (Robineau-Desvoidy, 1863)	B	M	S
<i>Phorinia</i> Robineau-Desvoidy, 1830			
<i>aurifrons</i> Robineau-Desvoidy, 1830	B	M	S
<i>Phorocera</i> Robineau-Desvoidy, 1830			
<i>assimilis</i> (Fallén, 1810)	B	M	S
<i>grandis</i> (Rondani, 1859)	B	M	S
<i>obscura</i> (Fallén, 1810)	B	M	S
BLONDELIINI			
<i>Admontia</i> Brauer & Bergenstamm, 1889			
<i>blanda</i> (Fallén, 1820)		B	M
<i>grandicornis</i> (Zetterstedt, 1849)		B	M
<i>maculisquama</i> (Zetterstedt, 1859)		B	M
<i>podomyia</i> (Brauer & Bergenstamm, 1889)	B	M	S
<i>seria</i> (Meigen, 1824)		M	S
<i>Belida</i> Robineau-Desvoidy, 1863			
<i>angelicae</i> (Meigen, 1824)		M	S
<i>latifrons</i> (Jacentkovský, 1944)		M	
<i>Blondelia</i> Robineau-Desvoidy, 1830			
<i>inclusa</i> (Hartig, 1838)		M	S
<i>nigripes</i> (Fallén, 1810)	B	M	S
<i>pinariae</i> (Hartig, 1838)	B	M	S
<i>Compsilura</i> Bouché, 1834			
<i>concinata</i> (Meigen, 1824)	B	M	S
<i>Conogaster</i> Brauer & Bergenstamm, 1891			
<i>pruinosa</i> (Meigen, 1824)	B	M	S
<i>Gastrolepta</i> Rondani, 1862			
<i>anthracina</i> (Meigen, 1826)	B	M	S
<i>Hemimacquartia</i> Brauer & Bergenstamm, 1893			
<i>paradoxa</i> Brauer & Bergenstamm, 1893		M	S
<i>Istocheta</i> Rondani, 1859			
<i>cinerea</i> (Macquart, 1850)	B	M	S
<i>longicornis</i> (Fallén, 1810)	B		
<i>Lecanipa</i> Rondani, 1859			
<i>bicincta</i> (Meigen, 1824)	B	M	S
<i>leucomelas</i> (Meigen, 1824)	B		S
<i>Leiophora</i> Robineau-Desvoidy, 1863			
<i>innoxia</i> (Meigen, 1824)	B	M	S
<i>Ligeria</i> Robineau-Desvoidy, 1863			
<i>angusticornis</i> (Loew, 1847)	B	M	S
<i>Ligeriella</i> Mesnil, 1962			
<i>aristata</i> (Villeneuve, 1911)			S
<i>Lomachantha</i> Rondani, 1859			
<i>parra</i> Rondani, 1859			S
<i>Medina</i> Robineau-Desvoidy, 1830			
<i>collaris</i> (Fallén, 1820)	B	M	S
<i>luctuosa</i> (Meigen, 1824)	B	M	S
<i>melania</i> (Meigen, 1824)	B	M	S
<i>multispina</i> (Herting, 1966)			S
<i>separata</i> (Meigen, 1824)	B	M	
<i>Meigenia</i> Robineau-Desvoidy, 1830			
<i>dorsalis</i> (Meigen, 1824)	B	M	S
<i>grandigena</i> (Pandellé, 1896)			S

<i>incana</i> (Fallén, 1810)	B M S	<i>quadripustulata</i> (Fabricius, 1794)	B M S
<i>majuscula</i> (Rondani, 1859)	M S	<i>speciosa</i> (Egger, 1861)	M S
<i>mutabilis</i> (Fallén, 1810)	B M S	<i>variegata</i> (Meigen, 1824)	B M S
<i>uncinata</i> Mesnil, 1967	B M S	<i>venusta</i> (Meigen, 1824)	B M S
Oswaldia Robineau-Desvoidy, 1863			
<i>eggeri</i> (Brauer & Bergenstamm, 1889)	M S	ERYCIINI	
<i>muscaria</i> (Fallén, 1810)	B M S	Alsomyia Brauer & Bergenstamm, 1891	
<i>reducta</i> (Villeneuve, 1930)	M S	<i>capillata</i> (Rondani, 1859)	B M
<i>spectabilis</i> (Meigen, 1824)	B M S	Amelibaea Mesnil, 1955	
Paracraspedothrix Villeneuve, 1919		<i>tultschensis</i> (Brauer & Bergenstamm, 1891)	M S
<i>montivaga</i> Villeneuve, 1919	B	Aplomya Robineau-Desvoidy, 1830	
Paratrixa Brauer & Bergenstamm, 1891		<i>confinis</i> (Fallén, 1820)	B M S
<i>polonica</i> Brauer & Bergenstamm, 1891	M	Bactromyia Brauer & Bergenstamm, 1891	
Picconia Robineau-Desvoidy, 1863		<i>aurulenta</i> (Meigen, 1824)	B S
<i>incurva</i> (Zetterstedt, 1844)	M S	Buquetia Robineau-Desvoidy, 1847	
Policheta Rondani, 1856		<i>musca</i> Robineau-Desvoidy, 1847	B M S
<i>unicolor</i> (Fallén, 1820)	B M S	Cadurciella Villeneuve, 1927	
Staurochaeta Brauer & Bergenstamm, 1889		<i>tritaeniata</i> (Rondani, 1859)	M
<i>albocingulata</i> (Fallén, 1820)	S	Carcelia Robineau-Desvoidy, 1830	
Vibrissina Rondani, 1861		<i>atricosta</i> Herting, 1961	B M
<i>debilitata</i> (Pandellé, 1896)	M	<i>bombylans</i> Robineau-Desvoidy, 1830	B M S
<i>turrata</i> (Meigen, 1824)	B S	<i>dubia</i> (Brauer & Bergenstamm, 1891)	S
Zaira Robineau-Desvoidy, 1830		<i>falenaria</i> (Rondani, 1859)	M S
<i>cinerea</i> (Fallén, 1810)	B M S	<i>gnava</i> (Meigen, 1824)	B M S
ACEMYINI			
Acemya Robineau-Desvoidy, 1830		<i>iliaca</i> (Ratzeburg, 1840)	S
<i>acuticornis</i> (Meigen, 1824)	B M S	<i>kowarzi</i> Villeneuve, 1912	M S
<i>rufitibia</i> (von Roser, 1840)	B M S	<i>laxifrons</i> Villeneuve, 1912	B M S
ETHILLINI			
Atylomyia Brauer, 1898		<i>lucorum</i> (Meigen, 1824)	B M S
<i>loewi</i> Brauer, 1898	B M S	<i>puberula</i> Mesnil, 1941	B M S
Ethilla Robineau-Desvoidy, 1863		<i>rasa</i> (Macquart, 1849)	S
<i>aemula</i> (Meigen, 1824)	B M S	<i>tibialis</i> (Robineau-Desvoidy, 1863)	S
Paratryphera Brauer & Bergenstamm, 1891		Catagonia Brauer & Bergenstamm, 1891	
<i>barbatula</i> (Rondani, 1859)	B M S	<i>aberrans</i> (Rondani, 1859)	M S
<i>bisetosa</i> (Brauer & Bergenstamm, 1891)	S	Drino Robineau-Desvoidy, 1863	
<i>palpalis</i> (Rondani, 1859)	B S	<i>atropivora</i> Robineau-Desvoidy, 1830	S
Prosethilla Herting, 1984		<i>bohémica</i> Mesnil, 1949	B
<i>kramerella</i> (Stein, 1924)	S	<i>galii</i> (Brauer & Bergenstamm, 1891)	B
WINTHEMIINI			
Nemorilla Rondani, 1856		<i>gilva</i> (Hartig, 1838)	B M S
<i>floralis</i> (Fallén, 1810)	B M S	<i>inconspicua</i> (Meigen, 1830)	B M S
<i>maculosa</i> (Meigen, 1824)	B M S	<i>lota</i> (Meigen, 1824)	B M S
Rhaphiochaeta Brauer & Bergenstamm, 1889		<i>vicina</i> (Zetterstedt, 1849)	B M S
<i>brevisetata</i> (Zetterstedt, 1838)	B M S	Epicampocera Macquart, 1849	
Smidia Robineau-Desvoidy, 1830		<i>succincta</i> (Meigen, 1824)	B M S
<i>conspersa</i> (Meigen, 1824)	B M S	Erycia Robineau-Desvoidy, 1830	
Timavia Robineau-Desvoidy, 1863		<i>fasciata</i> Villeneuve, 1924	B S
<i>amoena</i> (Meigen, 1824)	B M S	<i>fatua</i> (Meigen, 1824)	B M S
Winthemia Robineau-Desvoidy, 1830		<i>festinans</i> (Meigen, 1824)	S
<i>bohemanii</i> (Zetterstedt, 1844)	B M S	<i>furibunda</i> (Zetterstedt, 1844)	B M S
<i>eruentata</i> (Rondani, 1859)	B M S	Huebneria Robineau-Desvoidy, 1847	
<i>erythrura</i> (Meigen, 1838)	B M S	<i>affinis</i> (Fallén, 1810)	B M S
<i>jacentkovskyi</i> Mesnil, 1949	B M S	Lydella Robineau-Desvoidy, 1830	
		<i>griseocens</i> Robineau-Desvoidy, 1830	B M S
		<i>stabulans</i> (Meigen, 1824)	B M S
		<i>thompsoni</i> Herting, 1959	M S
		Nilea Robineau-Desvoidy, 1863	
		<i>hortulana</i> (Meigen, 1824)	B S

<i>innoxia</i> Robineau-Desvoidy, 1863	B	S	<i>Brachicheta</i> Rondani, 1861	
<i>rufiscutellaris</i> (Zetterstedt, 1859)	B	M	<i>strigata</i> (Meigen, 1824)	B M S
<i>Periarchiclops</i> Villeneuve, 1924			<i>Cadurcia</i> Villeneuve, 1926	
<i>scutellaris</i> (Fallén, 1820)	B	M S	<i>casta</i> (Rondani, 1861)	S
<i>Phebellia</i> Robineau-Desvoidy, 1846			<i>Ceromasia</i> Rondani, 1856	
<i>clavellariae</i> (Brauer & Bergenstamm, 1891)	B		<i>rubrifrons</i> (Macquart, 1834)	B M S
<i>glauca</i> (Meigen, 1824)	B	M S	<i>Clemelis</i> Robineau-Desvoidy, 1863	
<i>glaucoides</i> Herting, 1961	B	S	<i>pullata</i> (Meigen, 1824)	B M S
<i>glirina</i> (Rondani, 1859)	B	M	<i>Cyzenis</i> Robineau-Desvoidy, 1863	
<i>nigripalpis</i> (Robineau-Desvoidy, 1847)	B	M S	<i>albicans</i> (Fallén, 1810)	B M S
<i>snulta</i> (Zetterstedt, 1844)		M	<i>jucunda</i> (Meigen, 1838)	S
<i>triseta</i> (Pandellé, 1896)	B		<i>Elodia</i> Robineau-Desvoidy, 1863	
<i>villica</i> (Zetterstedt, 1838)	B	S	<i>ambulatoria</i> (Meigen, 1824)	B M S
<i>Phonomyia</i> Brauer & Bergenstamm, 1893			<i>morio</i> (Fallén, 1820)	B M S
<i>aristata</i> (Rondani, 1861)		M S	<i>Erycilla</i> Mesnil, 1957	
<i>Phryxe</i> Robineau-Desvoidy, 1830			<i>ferruginea</i> (Meigen, 1824)	B M S
<i>erythrostoma</i> (Hartig, 1838)	B	M S	<i>Erythroceras</i> Robineau-Desvoidy, 1848	
<i>heraclei</i> (Meigen, 1824)	B	M S	<i>nigripes</i> (Robineau-Desvoidy, 1830)	B M S
<i>hirta</i> (Bigot, 1880)		M	<i>Eumea</i> Robineau-Desvoidy, 1863	
<i>magnicornis</i> (Zetterstedt, 1838)	B	S	<i>linearicornis</i> (Zetterstedt, 1844)	B M S
<i>nemea</i> (Meigen, 1824)	B	M S	<i>mitis</i> (Meigen, 1824)	B M S
<i>prima</i> (Brauer & Bergenstamm, 1889)	B	M S	<i>Eurysthaea</i> Robineau-Desvoidy, 1863	
<i>vulgaris</i> (Fallén, 1810)	B	M S	<i>scutellaris</i> (Robineau-Desvoidy, 1848)	B M S
<i>Pseudoperichaeta</i> Brauer & Bergenstamm, 1889			<i>Frontina</i> Meigen, 1838	
<i>nigrolineata</i> (Walker, 1853)	B	M S	<i>laeta</i> (Meigen, 1824)	B M S
<i>palesoidea</i> (Robineau-Desvoidy, 1830)		M S	<i>Gaedia</i> Meigen, 1838	
<i>Ptesiomyia</i> Brauer & Bergenstamm, 1893			<i>connexa</i> (Meigen, 1824)	M S
<i>alacris</i> (Meigen, 1824)	B	M S	<i>distincta</i> Egger, 1861	M S
<i>Senometopia</i> Macquart, 1834			<i>Gonia</i> Meigen, 1803	
<i>confundens</i> (Rondani, 1859)		S	<i>atra</i> Meigen, 1826	M
<i>excisa</i> (Fallén, 1820)	B	M S	<i>capitata</i> (DeGeer, 1776)	B M S
<i>pilosa</i> (Baranov, 1931)		M	<i>distinguenda</i> Herting, 1963	S
<i>pollinosa</i> (Mesnil, 1941)	B	M S	<i>divisa</i> Meigen, 1826	B M S
<i>separata</i> (Rondani, 1859)	B	M S	<i>foersteri</i> Meigen, 1838	M S
<i>susurrans</i> (Rondani, 1859)		M S	<i>ornata</i> Meigen, 1826	B M S
<i>Thecocarcelia</i> Townsend, 1933			<i>picea</i> (Robineau-Desvoidy, 1830)	B M S
<i>acutangulata</i> (Macquart, 1850)	B	M	<i>vacua</i> Meigen, 1826	M S
<i>Thelymyia</i> Brauer & Bergenstamm, 1891			<i>Hebia</i> Robineau-Desvoidy, 1830	
<i>alium</i> (Meigen, 1824)	B	M	<i>flavipes</i> Robineau-Desvoidy, 1830	B M S
<i>Tlephusa</i> Robineau-Desvoidy, 1863			<i>Masicera</i> Macquart, 1834	
<i>cincinna</i> (Rondani, 1859)	B	M S	<i>pavoniae</i> (Robineau-Desvoidy, 1830)	B M S
<i>Townsendiellomyia</i> Baranov, 1932			<i>silvatica</i> (Fallén, 1810)	B M S
<i>nidicola</i> (Townsend, 1908)	B	M S	<i>sphingivora</i> (Robineau-Desvoidy, 1830)	S
<i>Xylotachina</i> Brauer & Bergenstamm, 1891			<i>Masistylum</i> Brauer & Bergenstamm, 1893	
<i>dihuta</i> (Meigen, 1824)	B	M S	<i>arcuatum</i> (Mik, 1863)	S
GONIINI			<i>Myxexoristops</i> Townsend, 1911	
<i>Baumhaueria</i> Meigen, 1838			<i>abietis</i> Herting, 1964	B M
<i>goniaeformis</i> (Meigen, 1824)	B	M S	<i>bicolor</i> (Villeneuve, 1908)	B M S
<i>Blepharipa</i> Rondani, 1856			<i>blondeli</i> (Robineau-Desvoidy, 1830)	B M S
<i>pratensis</i> (Meigen, 1824)	B	M S	<i>bonsdorffi</i> (Zetterstedt, 1859)	B M S
<i>schineri</i> (Mesnil, 1939)	B	M S	<i>stolida</i> (Stein, 1924)	B S
<i>Bothria</i> Rondani, 1856			<i>Ocytata</i> Gistel, 1848	
<i>frontosa</i> (Meigen, 1824)	B	M S	<i>pallipes</i> (Fallén, 1820)	B M S
<i>subalpina</i> Villeneuve, 1910	B	M S	<i>Onychogonia</i> Brauer & Bergenstamm, 1889	
			<i>flaviceps</i> (Zetterstedt, 1838)	B S
			<i>Pales</i> Robineau-Desvoidy, 1830	
			<i>pavida</i> (Meigen, 1824)	B M S

<i>processioneae</i> (Ratzeburg, 1840)	S	<i>Linnaemya</i> Robineau-Desvoidy, 1830	
<i>pumicata</i> (Meigen, 1824)	M	<i>comta</i> (Fallén, 1810)	B M S
<i>Pexopsis</i> Brauer & Bergenstamm, 1889		<i>fissiglobula</i> Pandellé, 1895	S
<i>aprica</i> (Meigen, 1824)	M S	<i>frater</i> (Rondani, 1859)	M S
<i>Phryno</i> Robineau-Desvoidy, 1830		<i>haemorrhoidalis</i> (Fallén, 1810)	B M S
<i>vetula</i> (Meigen, 1824)	B M S	<i>helvetica</i> Herting, 1963	S
<i>Platymya</i> Robineau-Desvoidy, 1830		<i>impudica</i> (Rondani, 1859)	B M S
<i>fimbriata</i> (Meigen, 1824)	B M S	<i>media</i> Zimin, 1954	B S
<i>Prosopea</i> Rondani, 1861		<i>olsufjevi</i> Zimin, 1954	B
<i>nigricans</i> (Egger, 1861)	M S	<i>perinealis</i> Pandellé, 1895	S
<i>Pseudogonia</i> Brauer & Bergenstamm, 1889		<i>picta</i> (Meigen, 1824)	B M S
<i>paristaca</i> (Robineau-Desvoidy, 1851)	S	<i>rossica</i> Zimin, 1954	S
<i>rufifrons</i> (Wiedemann, 1830)	B S	<i>steini</i> Jacentkovský, 1944	M S
<i>Spallanzania</i> Robineau-Desvoidy, 1830		<i>tessellans</i> (Robineau-Desvoidy, 1830)	B M S
<i>hebes</i> (Fallén, 1820)	M S	<i>vulpina</i> (Fallén, 1810)	B M S
<i>Sturmia</i> Robineau-Desvoidy, 1830		<i>Lydina</i> Robineau-Desvoidy, 1830	
<i>bella</i> (Meigen, 1824)	B M S	<i>aenea</i> (Meigen, 1824)	B M S
<i>Thelymorpha</i> Brauer & Bergenstamm, 1889		<i>Lypha</i> Robineau-Desvoidy, 1830	
<i>marmorata</i> (Fabricius, 1805)	B M S	<i>dubia</i> (Fallén, 1810)	B M S
<i>Zenillia</i> Robineau-Desvoidy, 1830		<i>ruficauda</i> (Zetterstedt, 1838)	B M S
<i>dolosa</i> (Meigen, 1824)	M S	<i>Petagnia</i> Rondani, 1856	
<i>libatrix</i> (Panzer, 1798)	B M S	<i>subpetiolata</i> Rondani, 1859	M
TACHININAE		ERNESTIINI	
TACHININI		<i>Appendicia</i> Stein, 1924	
<i>Germania</i> Robineau-Desvoidy, 1830		<i>truncata</i> (Zetterstedt, 1838)	B
<i>angustata</i> (Zetterstedt, 1844)	S	<i>Cleonice</i> Robineau-Desvoidy, 1863	
<i>ruficeps</i> (Fallén, 1820)	B M S	<i>callida</i> (Meigen, 1824)	B M S
<i>Nowickia</i> Wachtl, 1894		<i>nitidiuscula</i> (Zetterstedt, 1859)	B
<i>atripalpis</i> (Robineau-Desvoidy, 1863)	M S	<i>Eloceria</i> Robineau-Desvoidy, 1863	
<i>ferox</i> (Panzer, 1809)	B M S	<i>delecta</i> (Meigen, 1824)	M
<i>marklini</i> (Zetterstedt, 1838)	B M S	<i>Ernestia</i> Robineau-Desvoidy, 1830	
<i>Peleteria</i> Robineau-Desvoidy, 1830		<i>argentifera</i> (Meigen, 1824)	B M S
<i>ferina</i> (Zetterstedt, 1844)	B M S	<i>laevigata</i> (Meigen, 1838)	B M S
<i>rubescens</i> (Robineau-Desvoidy, 1830)	B M S	<i>puparum</i> (Fabricius, 1794)	B M S
<i>ruficornis</i> Macquart, 1835	S	<i>rudis</i> (Fallén, 1810)	B M S
<i>varia</i> (Fabricius, 1794)	M S	<i>vagans</i> (Meigen, 1824)	B M S
<i>Schineria</i> Rondani, 1857		<i>Eurithia</i> Robineau-Desvoidy, 1830	
<i>tergestina</i> Rondani, 1859	S	<i>anthophila</i> (Robineau-Desvoidy, 1830)	B M S
<i>Tachina</i> Meigen, 1803		<i>caesia</i> (Fallén, 1810)	B M S
<i>casta</i> Rondani, 1859	B M S	<i>connivens</i> (Zetterstedt, 1844)	B M S
<i>fera</i> (Linnaeus, 1761)	B M S	<i>consobrina</i> (Meigen, 1824)	B M S
<i>grossa</i> (Linnaeus, 1758)	B M S	<i>indigens</i> (Pandellé, 1896)	M S
<i>lurida</i> (Fabricius, 1781)	B M S	<i>intermedia</i> (Zetterstedt, 1844)	B M S
<i>magnicornis</i> (Zetterstedt, 1844)	B M S	<i>vivida</i> (Zetterstedt, 1838)	B M S
<i>nigrohirta</i> (Stein, 1924)	M S	<i>Fausta</i> Robineau-Desvoidy, 1830	
<i>nupta</i> (Rondani, 1859)	M	<i>nemorum</i> (Meigen, 1824)	B M S
<i>praeceps</i> Meigen, 1824	B S	<i>Gymnocheta</i> Robineau-Desvoidy, 1830	
<i>ursina</i> (Meigen, 1824)	B M S	<i>viridis</i> (Fallén, 1810)	B M S
NEMORAEINI		<i>Hyalurgus</i> Brauer & Bergenstamm, 1893	
<i>Nemoraeta</i> Robineau-Desvoidy, 1830		<i>lucidus</i> (Meigen, 1824)	B M S
<i>pellucida</i> (Meigen, 1824)	B M S	<i>tomostethi</i> Čepelák, 1963	M
LINNAEMYINI		<i>Loewia</i> Egger, 1856	
<i>Chrysosomopsis</i> Townsend, 1916		<i>brevifrons</i> Rondani, 1856	M S
<i>auratus</i> (Fallén, 1820)	B M S	<i>foeda</i> (Meigen, 1824)	M S
		<i>mudigena</i> Mesnil, 1973	M
		<i>phaeoptera</i> (Meigen, 1824)	B M S

<i>piligena</i> Mesnil, 1973	S	<i>lichtwardtiana</i> (Villeneuve, 1931)	B
<i>Synactia</i> Villeneuve, 1916		<i>samarensis</i> (Villeneuve, 1921)	B
<i>parvula</i> (Rondani, 1861)	B M S	<i>selecta</i> (Pandellé, 1894)	M S
<i>Zophomyia</i> Macquart, 1835		<i>siphonoides</i> (Strobl, 1898)	B M S
<i>temula</i> (Scopoli, 1763)	B M S	<i>starkei</i> (Mesnil, 1952)	B
		<i>temuipalpis</i> (Villeneuve, 1921)	M
BRACHYMERINI		<i>Ceromya</i> Robineau-Desvoidy, 1830	
<i>Pseudopachystylum</i> Mik, 1891		<i>bicolor</i> (Meigen, 1824)	B
<i>gonioides</i> (Zetterstedt, 1838)	B M S	<i>flaviceps</i> (Ratzeburg, 1844)	B S
		<i>silacea</i> (Meigen, 1824)	B M
PELATACHININI		<i>Entomophaga</i> Lioy, 1864	
<i>Pelatachina</i> Meade, 1894		<i>exoleta</i> (Meigen, 1824)	S
<i>tibialis</i> (Fallén, 1810)	B M S	<i>nigrohalterata</i> (Villeneuve, 1921)	B M S
		<i>Goniocera</i> Brauer & Bergenstamm, 1891	
MACQUARTIINI		<i>versicolor</i> (Fallén, 1820)	B M S
<i>Macquartia</i> Robineau-Desvoidy, 1830		<i>Peribaea</i> Robineau-Desvoidy, 1863	
<i>dispar</i> (Fallén, 1820)	B M S	<i>apicalis</i> Robineau-Desvoidy, 1863	S
<i>grisea</i> (Fallén, 1810)	B M S	<i>fissicornis</i> (Strobl, 1909)	B M S
<i>chalconota</i> (Meigen, 1824)	B M S	<i>tibialis</i> (Robineau-Desvoidy, 1851)	B M S
<i>macularis</i> Villeneuve, 1926	S	<i>Siphona</i> Meigen, 1803	
<i>nudigena</i> Mesnil, 1972	B S	<i>boreata</i> Mesnil, 1960	B M S
<i>praefica</i> (Meigen, 1824)	B M S	<i>collini</i> Mesnil, 1960	B S
<i>pubiceps</i> (Zetterstedt, 1845)	B M S	<i>confusa</i> Mesnil, 1961	B M S
<i>tenebricosa</i> (Meigen, 1824)	B M S	<i>cristata</i> (Fabricius, 1805)	B M S
<i>tessellum</i> (Meigen, 1824)	S	<i>flavifrons</i> Staeger, 1849	B M S
<i>viridana</i> Robineau-Desvoidy, 1863	B M S	<i>geniculata</i> (Degeer, 1776)	B M S
<i>Macroprosopa</i> Brauer & Bergenstamm, 1889		<i>grandistylum</i> Pandellé, 1894	S
<i>atrata</i> (Fallén, 1810)	B M S	<i>ingerae</i> Andersen, 1982	S
		<i>maculata</i> Staeger, 1849	B M S
TRIARTHRIINI		<i>nigricans</i> (Villeneuve, 1930)	M
<i>Triarthria</i> Stephens, 1829		<i>paludosa</i> Mesnil, 1960	B S
<i>setipennis</i> (Fallén, 1810)	B M S	<i>pauciseta</i> Rondani, 1865	B S
<i>Trichactia</i> Stein, 1824		<i>setosa</i> Mesnil, 1960	B S
<i>pictiventris</i> (Zetterstedt, 1852)	B M S		
		LESKIINI	
NEARINI		<i>Aphria</i> Robineau-Desvoidy, 1830	
<i>Elfia</i> Robineau-Desvoidy, 1850		<i>latifrons</i> Villeneuve, 1908	M S
<i>bohémica</i> (Kramer, 1907)	B	<i>longilingua</i> Rondani, 1861	S
<i>cingulata</i> (Robineau-Desvoidy, 1830)	B M S	<i>longirostris</i> (Meigen, 1824)	B M S
<i>minutissima</i> (Zetterstedt, 1844)	M	<i>xyphias</i> Pandellé, 1896	M
<i>zonella</i> (Zetterstedt, 1844)	B M S	<i>Atylostoma</i> Brauer & Bergenstamm, 1889	
<i>Phytomyptera</i> Rondani, 1845		<i>tricolor</i> (Mik, 1884)	S
<i>nigrina</i> (Meigen, 1824)	S	<i>Bithia</i> Robineau-Desvoidy, 1863	
		<i>acanthophora</i> (Rondani, 1861)	S
SIPHONINI		<i>demotica</i> (Egger, 1861)	M S
<i>Actia</i> Robineau-Desvoidy, 1830		<i>glirina</i> (Rondani, 1861)	B M S
<i>crassicornis</i> (Meigen, 1824)	B M S	<i>immaculata</i> (Herting, 1971)	S
<i>dubitata</i> Herting, 1971	B M S	<i>jacentkovskyi</i> (Villeneuve, 1937)	M
<i>infantula</i> (Zetterstedt, 1844)	B M S	<i>spreta</i> (Meigen, 1824)	B M S
<i>lamia</i> (Meigen, 1838)	B M S	<i>Clausicella</i> Rondani, 1856	
<i>maksymovi</i> Mesnil, 1952	S	<i>suturata</i> Rondani, 1859	S
<i>nigroscutellata</i> Lundbeck, 1927	B	<i>Demoticus</i> Macquart, 1854	
<i>nudibasis</i> Stein, 1924	M S	<i>plebejus</i> (Fallén, 1810)	B M S
<i>pilipennis</i> (Fallén, 1810)	B M S	<i>Leskia</i> Robineau-Desvoidy, 1830	
<i>Ceranthia</i> Robineau-Desvoidy, 1830		<i>aurea</i> (Fallén, 1820)	B M S
<i>abdominalis</i> (Robineau-Desvoidy, 1830)	M S	<i>Solieria</i> Robineau-Desvoidy, 1848	
<i>brunnescens</i> (Villeneuve, 1921)	B	<i>fenestrata</i> (Meigen, 1824)	B M S

<i>inanis</i> (Fallén, 1810)	B M S	<i>inepta</i> (Meigen, 1824)	B M S
<i>pacifica</i> (Meigen, 1824)	B M S	<i>praecox</i> (Meigen, 1824)	B M S
<i>vacua</i> (Rondani, 1861)	M S	<i>Cyrtophleba</i> Rondani, 1856	
		<i>ruricola</i> (Meigen, 1824)	B M S
		<i>vernalis</i> (Kramer, 1917)	B S
MINTHOINI		<i>Eriothrix</i> Meigen, 1803	
<i>Mintho</i> Robineau-Desvoidy, 1830		<i>apenninus</i> (Rondani, 1862)	M S
<i>rufiventris</i> (Fallén, 1816)	B M S	<i>argyreatus</i> (Meigen, 1824)	B M S
		<i>prolixa</i> (Meigen, 1824)	B M S
MICROPTHALMINI		<i>rufomaculatus</i> (Degeer, 1776)	B M S
<i>Dexiosoma</i> Rondani, 1856		<i>Hyleorus</i> Aldrich, 1926	
<i>caninum</i> (Fabricius, 1781)	B M S	<i>elatus</i> (Meigen, 1838)	B
<i>Melisoneura</i> Rondani, 1861		<i>Klugia</i> Robineau-Desvoidy, 1863	
<i>leucoptera</i> (Meigen, 1824)	M	<i>marginata</i> (Meigen, 1824)	B M S
<i>Microphthalma</i> Macquart, 1843		<i>Periscepsia</i> Gistel, 1848	
<i>europaea</i> Egger, 1860	B M S	<i>carbonaria</i> (Panzer, 1798)	B M S
		<i>Peteina</i> Meigen, 1838	
DEXIINAE		<i>erinaceus</i> (Fabricius, 1794)	B S
DEXIINI		<i>Phenicellia</i> Robineau-Desvoidy, 1830	
<i>Billaea</i> Robineau-Desvoidy, 1830		<i>haematodes</i> (Meigen, 1824)	B
<i>adelpha</i> (Loew, 1873)	M S	<i>Phyllomya</i> Robineau-Desvoidy, 1830	
<i>irrorata</i> (Meigen, 1826)	B M S	<i>volvulus</i> (Fabricius, 1794)	B M S
<i>pectinata</i> (Meigen, 1826)	B M S	<i>Ramonda</i> Robineau-Desvoidy, 1863	
<i>triangulifera</i> (Zetterstedt, 1844)	B M S	<i>delphinensis</i> (Villeneuve, 1922)	S
<i>Dexia</i> Meigen, 1826		<i>latifrons</i> (Zetterstedt, 1844)	B S
<i>rustica</i> (Fabricius, 1775)	B M S	<i>prunaria</i> (Rondani, 1861)	B M S
<i>vacua</i> (Fallén, 1816)	B S	<i>prunicia</i> (Herting, 1969)	S
<i>Dinera</i> Robineau-Desvoidy, 1863		<i>spathulata</i> (Fallén, 1820)	B M S
<i>carinifrons</i> (Fallén, 1816)	B M S	<i>Stomina</i> Robineau-Desvoidy, 1830	
<i>ferina</i> (Fallén, 1816)	B M S	<i>caliendrata</i> (Rondani, 1862)	S
<i>grisescens</i> (Fallén, 1816)	B M S	<i>iners</i> (Meigen, 1824)	S
<i>Estheria</i> Robineau-Desvoidy, 1830		<i>tachinoides</i> (Fallén, 1816)	S
<i>bohemanii</i> (Rondani, 1862)	B M	<i>Thelaira</i> Robineau-Desvoidy, 1830	
<i>cristata</i> (Meigen, 1826)	B M S	<i>nigripes</i> (Fabricius, 1794)	B M S
<i>microcera</i> Robineau-Desvoidy, 1830	S	<i>solivaga</i> (Harris, 1780)	B S
<i>petiolata</i> (Bonsdorff, 1866)	B M S	<i>Trafoia</i> Brauer & Bergenstamm, 1893	
<i>picta</i> (Meigen, 1826)	M S	<i>monticola</i> Brauer & Bergenstamm, 1893	M S
<i>Prosenia</i> LePeletier & Serville, 1828		<i>Voria</i> Robineau-Desvoidy, 1830	
<i>siberita</i> (Fabricius, 1775)	B M S	<i>ruralis</i> (Fallén, 1810)	B M S
<i>Trixa</i> Meigen, 1824		<i>Wagneria</i> Robineau-Desvoidy, 1830	
<i>alpina</i> Meigen, 1824	M S	<i>costata</i> (Fallén, 1815)	M S
<i>caerulescens</i> Meigen, 1824	M S	<i>cunctans</i> (Meigen, 1824)	M S
<i>conspersa</i> (Harris, 1776)	B M S	<i>gagatea</i> Robineau-Desvoidy, 1830	M S
<i>Zeuxia</i> Meigen, 1826			
<i>brevicornis</i> (Egger, 1860)	S		
<i>cinerea</i> Meigen, 1826	B M S	DUFOURIINI	
<i>subapennina</i> Rondani, 1862	M S	<i>Dufouria</i> Robineau-Desvoidy, 1830	
		<i>chalybeata</i> (Meigen, 1824)	B M S
VORIINI		<i>nigrita</i> (Fallén, 1810)	B M S
<i>Athrycia</i> Robineau-Desvoidy, 1830		<i>occlusa</i> (Robineau-Desvoidy, 1863)	B M S
<i>curvinervis</i> (Zetterstedt, 1844)	B M S	<i>Freraea</i> Robineau-Desvoidy, 1830	
<i>impressa</i> (Wulp, 1869)	M S	<i>gagatea</i> Robineau-Desvoidy, 1830	B S
<i>trepida</i> (Meigen, 1824)	B M S	<i>Chaetoptilia</i> Rondani, 1862	
<i>Blepharomyia</i> Brauer & Bergenstamm, 1889		<i>puella</i> (Rondani, 1862)	M
<i>pagana</i> (Meigen, 1824)	B M S	<i>Microsoma</i> Macquart, 1855	
<i>piliceps</i> (Zetterstedt, 1859)	S	<i>exiguum</i> (Meigen, 1824)	B M S
<i>Campylocheta</i> Rondani, 1859		<i>Rondania</i> Robineau-Desvoidy, 1850	
<i>fuscinervis</i> (Stein, 1924)	B S	<i>cucullata</i> Robineau-Desvoidy, 1850	B M S

<i>dimidiata</i> (Meigen, 1824)	S	LEUCOSTOMATINI	
<i>fasciata</i> (Macquart, 1834)	B M S	<i>Brullaea</i> Robineau-Desvoidy, 1863	
PHASIINAE		<i>ocypteroidea</i> Robineau-Desvoidy, 1863	M S
EUTHERINI		<i>Cinochira</i> Zetterstedt, 1845	
<i>Redtenbacheria</i> Schiner, 1861		<i>atra</i> Zetterstedt, 1845	B M
<i>insignis</i> Egger, 1861	M	<i>Clairvillia</i> Robineau-Desvoidy, 1830	
PHASIINI		<i>biguttata</i> (Meigen, 1824)	B
<i>Cistogaster</i> Latreille, 1829		<i>Dionaea</i> Robineau-Desvoidy, 1830	
<i>globosa</i> (Fabricius, 1775)	B M S	<i>aurifrons</i> (Meigen, 1824)	B M S
<i>Clytiomya</i> Robineau-Desvoidy, 1861		<i>Eulabidogaster</i> Belanovsky, 1951	
<i>continua</i> (Panzer, 1798)	B M S	<i>setifacies</i> (Rondani, 1861)	B M S
<i>Ectophasia</i> Townsend, 1912		<i>Labigastera</i> Macquart, 1834	
<i>crassipennis</i> (Fabricius, 1794)	B M S	<i>forcipata</i> (Meigen, 1824)	B M S
<i>oblonga</i> (Robineau-Desvoidy, 1830)	B M S	<i>nitidula</i> (Meigen, 1824)	S
<i>Elizeta</i> Rondani, 1856		<i>pauciseta</i> Rondani, 1861	M S
<i>helluo</i> (Fabricius, 1805)	B M S	<i>Leucostoma</i> Meigen, 1803	
<i>pellucens</i> (Fallén, 1820)	B M S	<i>anthracinum</i> (Meigen, 1824)	B M S
<i>Elomya</i> Robineau-Desvoidy, 1830		<i>simplex</i> (Fallén, 1815)	B M S
<i>lateralis</i> (Meigen, 1824)	M S	<i>tetraptera</i> (Meigen, 1824)	B M S
<i>Gymnosoma</i> Meigen, 1803		<i>turonicum</i> Dupuis, 1964	S
<i>brachypeltae</i> Dupuis, 1961	S	<i>Weberia</i> Robineau-Desvoidy, 1830	
<i>clavatum</i> (Rohdendorf, 1947)	B S	<i>digramma</i> (Meigen, 1824)	M
<i>costatum</i> (Panzer, 1800)	M S	CYLINDROMYIINI	
<i>desertorum</i> (Rohdendorf, 1947)	S	<i>Besseria</i> Robineau-Desvoidy, 1830	
<i>dolycoridis</i> Dupuis, 1961	B M S	<i>dimidiata</i> (Zetterstedt, 1844)	M S
<i>nitens</i> Meigen, 1824	B M S	<i>lateritia</i> (Meigen, 1824)	B
<i>nudifrons</i> Herting, 1966	B M S	<i>melanura</i> (Meigen, 1824)	B
<i>rotundatum</i> (Linnaeus, 1758)	B M S	<i>Cylindromyia</i> Meigen, 1803	
<i>Opesia</i> Robineau-Desvoidy, 1863		<i>auriceps</i> (Meigen, 1838)	B M S
<i>cana</i> (Meigen, 1824)	B M S	<i>bicolor</i> (Olivier, 1812)	M S
<i>semicana</i> (Egger, 1861)	B M S	<i>brassicaria</i> (Fabricius, 1775)	B M S
<i>Phasia</i> Latreille, 1804		<i>brevicornis</i> (Loew, 1844)	B M S
<i>aurigera</i> (Egger, 1860)	B M S	<i>crassa</i> (Loew, 1845)	S
<i>aurulans</i> Meigen, 1824	B M S	<i>intermedia</i> (Meigen, 1824)	B M S
<i>barbifrons</i> (Girschner, 1887)	B M S	<i>interrupta</i> (Meigen, 1824)	B M S
<i>hemiptera</i> (Fabricius, 1794)	B M S	<i>pilipes</i> (Loew, 1844)	B M S
<i>karczewskii</i> (Draber-Monko, 1965)	M	<i>pusilla</i> (Meigen, 1824)	B M S
<i>obesa</i> (Fabricius, 1798)	B M S	<i>rufifrons</i> (Loew, 1844)	S
<i>pandellei</i> (Dupuis, 1957)	S	<i>xylotina</i> (Egger, 1860)	B M S
<i>pusilla</i> Meigen, 1824	B M S	<i>Hemyda</i> Robineau-Desvoidy, 1830	
<i>subcoleopirata</i> (Linnaeus, 1767)	B M S	<i>obscuripennis</i> (Meigen, 1824)	B M S
<i>Subclytia</i> Pandellé, 1894		<i>vittata</i> (Meigen, 1824)	B M S
<i>rotundiventris</i> (Fallén, 1820)	B M S	<i>Lophosia</i> Meigen, 1824	
<i>Xysta</i> Meigen, 1824		<i>fasciata</i> Meigen, 1824	B M S
<i>holosericea</i> (Fabricius, 1805)	M S	<i>Phania</i> Meigen, 1824	
CATHAROSIINI		<i>curvicauda</i> (Fallén, 1820)	B M S
<i>Catharosia</i> Rondani, 1868		<i>funesta</i> (Meigen, 1824)	B M S
<i>pygmaea</i> (Fallén, 1815)	B M	<i>incrassata</i> Pandellé, 1894	B M S
<i>Litophasia</i> Girschner, 1887		<i>speculifrons</i> (Villeneuve, 1919)	B S
<i>hyalipennis</i> (Fallén, 1815)	B M S	<i>thoracica</i> Meigen, 1824	B M S
STRONGYGASTRINI		GASTEROPHILIDAE	
<i>Strongygaster</i> Macquart, 1834		Jan Minář	
<i>celer</i> (Meigen, 1838)	M S	<i>Gasterophilus</i> Leach, 1817	
<i>globula</i> (Meigen, 1824)	B M S	<i>haemorrhoidalis</i> (Linnaeus, 1758)	B S

<i>inermis</i> (Brauer, 1858)		S	<i>chloropus</i> Bergroth, 1901	M
<i>intestinalis</i> (De Geer, 1776)	B	M S	<i>fringilina</i> Curtis, 1836	B M S
<i>pecorum</i> (Fabricius, 1794)	B		<i>Ornithophila</i> Rondani, 1879	
			<i>metallica</i> (Schiner, 1864)	M
OESTRIDAE			<i>Olfersia</i> Wiedemann, 1830	
Jan Minář			<i>fumipennis</i> (Sahlberg, 1886)	M S
CEPHENEMYIINAE			<i>Icosta</i> Speiser, 1905	
<i>Cephenemyia</i> Latreille, 1818			<i>ardeae</i> (Macquart, 1835)	B M S
<i>auribarbis</i> (Meigen, 1824)	B	S	<i>minor</i> (Bigot, 1858)	M
<i>stimulator</i> (Clarck, 1815)	B	M S	<i>Pseudolynchia</i> Bequaert, 1926	
<i>Pharyngomyia</i> Schiner, 1861			<i>canariensis</i> (Macquart, 1840)	M S
<i>picta</i> (Meigen, 1824)	B	M S	<i>Crataerina</i> von Olfers, 1816	
			<i>pallida</i> (Latreille, 1812)	B M S
			<i>Stenepteryx</i> Leach, 1817	
			<i>hirundinis</i> (Linnaeus, 1758)	B M S
OESTRINAE			HIPPOBOSCINAE	
<i>Oestrus</i> Linnaeus, 1758			<i>Hippobosca</i> Linnaeus, 1758	
<i>ovis</i> Linnaeus, 1758	B	M S	<i>equina</i> Linnaeus, 1758	M S
<i>Rhinoestrus</i> Brauer, 1886			<i>longipennis</i> Fabricius, 1805	S
<i>purpureus</i> (Brauer, 1858)		S	<i>variegata</i> Megerle, 1803	S
HYPODERMATIDAE			LIPOPTENINAE	
Jan Minář			<i>Lipoptena</i> Nitsch, 1818	
OESTROMYIINAE			<i>cervi</i> (Linnaeus, 1758)	B M S
<i>Oestromyia</i> Brauer, 1860			<i>fortisetosa</i> Maa, 1965	B M S
<i>leporina</i> (Pallas, 1778)	B	M S	<i>Melophagus</i> Latreille, 1802	
			<i>ovinus</i> (Linnaeus, 1758)	B M S
			<i>rupicaprinus</i> Rondani, 1879	S
HYPODERMATINAE			NYCTERIBIIDAE	
<i>Hypoderma</i> Latreille, 1818			Karel Hürka	
<i>actaeon</i> Brauer, 1858	B	S	<i>Nycteribia</i> Latreille, 1796	
<i>bovis</i> (De Geer, 1776)	B	M S	<i>kolenatii</i> Theodor & Moscona, 1954	B M S
<i>diana</i> Brauer, 1858	B	M S	<i>latreillii</i> (Leach, 1817)	B M S
<i>lineatum</i> (De Villers, 1798)		M S	<i>schmidlii schmidlii</i> Schiner, 1853	S
HIPPOBOSCIDAE			<i>vexata</i> Westwood, 1835	B M S
Josef Chalupský & Dalibor Povolný			<i>Phthiridium</i> Hermann, 1804	
ORNITHOICINAE			<i>biarticulatum</i> Hermann, 1804	M S
<i>Ornithoica</i> Rondani, 1878			<i>Basilia</i> Miranda Ribeiro, 1903	
<i>turdi</i> (Latreille, 1811)		M S	<i>italica</i> Theodor, 1954	S
ORNITHOMYIINAE			<i>nana</i> Theodor & Moscona, 1954	B M S
<i>Ornithomyia</i> Latreille, 1802			<i>nattereri</i> (Kolenati, 1857)	M
<i>avicularia</i> (Linnaeus, 1758)	B	M S	<i>Penicillidia</i> Kolenati, 1863	
<i>biloba</i> Dufour, 1827	B	M S	<i>conspicua</i> Speiser, 1901	S
			<i>dufourii dufourii</i> (Westwood, 1835)	B M S
			<i>monoceros</i> Speiser, 1900	B M

- BARTÁK M., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Empididae, Lonchopteridae, Conopidae, Sepsidae, Piophilidae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*, Vol. 8, Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 228.
- BARTÁK M. & ROHÁČEK J., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Chyromyidae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*, Vol. 8, Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 217–218.
- BARTÁK M., TSCHORSNIG H. P. & VAŇHARA J., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Tachinidae (part 2). In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*, Vol. 8, Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 235–236.
- BARTÁK M., ZEMAN V. & ZUSKA J., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Sepsidae, Piophilidae, Chloropidae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*, Vol. 8, Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 226–227.
- BOSÁK J. 1996a: Faunistic records from the Czech Republic – 52. Diptera: Scenopinidae. *Klapalekiana*, 32: 270.
- BOSÁK J. 1996b: Zajímavé nálezy roupců (Diptera, Asilidae) na jižní Moravě. *Sbor. Přírodověd. klubu Uherské Hradiště*, 1: 78–79.
- BOSÁK J. 1997a: Scenopinidae (Diptera) of the Czech Republic and Slovak Republic. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*, Vol. 8, Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 23–25.
- BOSÁK J., 1997b: Faunistic Records from the Czech and Slovak Republics. Diptera: Bombyliidae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*, Vol. 8, Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 216–217.
- BOSÁK J. & HRADSKÝ M., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Asilidae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*, Vol. 8, Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 216.
- CHVÁLA M., 1997a: A revision of the European species of the *Hilara chorica*-complex (Diptera, Empididae), with new synonymy and description of a new species. *Studia dipterologica*, 4(1): 99–113.
- CHVÁLA M., 1997b: A revision of the European species of the *Hilara flavipes*-group (Diptera: Empididae) with new synonymies and descriptions of new species. *Studia dipterologica*, 4(2). (In press)
- CHVÁLA M., 1997c: A taxonomic revision of the *Hilara maura*-group (Diptera: Empididae) in Europe. *Syst. Entomol.*, 21(1996): 265–294.
- CHVÁLA M., 1997d: Additions and corrections to the faunistic data of the family Empididae (Diptera). In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*, Vol. 8, Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 43–46.
- CHVÁLA M., 1997e: Eleven new synonymies in European species of *Hilara* (Diptera: Empididae). *Acta Univ. Carol. – Biol.*, 41: 293–322.
- CHVÁLA M. & BARTÁK M., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Hybotidae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*, Vol. 8, Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 218–220.
- ČEPELÁK J., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Tachinidae (part 1). In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*, Vol. 8, Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 234–235.
- ČEPELÁK J., GREGOR F. & ROZKOŠNÝ R., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Calliphoridae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*, Vol. 8, Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 233.
- ČEPELÁK J. & ROZKOŠNÝ R., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Anthomyiidae, Muscidae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*, Vol. 8, Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 229.
- ČERNÝ M. & VÁLA M., 1996: Faunistic records of Agromyzidae (Diptera) from the Czech and Slovak Republics. – *Čas. Slez. Muz. Opava (A)*, 45: 157–169.
- EVENHUIS N. L. & BOSÁK J. 1997: New records of *Glabellula arctica* (Zetterstedt, 1838) (Mythicomyiidae, Diptera) from the Czech Republic and Slovak Republic. *Čas. Slez. Muz. Opava (A)*, 46: 95–96 (1997).
- GREGOR F., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Fanniidae, Muscidae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*, Vol. 8, Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 229–232.

- HALGOŠ J., PETRUS, O., 1996: First record of *Culex theileri* (Diptera, Culicidae) in Slovakia. *Biológia Bratislava*, 51: 190.
- HOLINKA J., 1997: Occurrence of hoverflies (Diptera, Syrphidae) in the environs of Pardubice. In: VANHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8. *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 47–55.
- HOLINKA J., MAZANEK L., VYSOKÝ V., MARTINOVSKÝ J. & KULA E., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Syrphidae. In: VANHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8. *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 224–226.
- HURKA K., 1997: New data on taxonomy and distribution of Palearctic, Oriental and Neotropical Ischnopsyllidae (Siphonaptera), Nycteribiidae and Streblidae (Diptera). *Acta Soc. Zool. Bohem.*, 61: 23–33.
- JEDLIČKA L., 1996: A synopsis of blackfly fauna of Slovakia (Diptera, Simuliidae). *Acta Zool. Univ. Comenianae*, 40: 49–67.
- JEŽEK J. (ed.), 1987: *Enumeratio insectorum Bohemoslovakiae*. Check list of Czechoslovak insects, Vol. 2 (Diptera). *Acta faun. ent. Mus. Nat. Pragae*, 18: 1–342.
- JEŽEK J., 1997a: Redescription and figures of males of two little known Tonnoir's species of moth flies (Diptera, Psychodidae). In: VANHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8. *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 63–71.
- JEŽEK J., 1997b: Moth flies (Diptera, Psychodidae) inhabiting Prague city and adjacent localities with descriptions of five new species. *Acta ent. Mus. Nat. Pragae*, 44 (1995): 125–165.
- KINKOROVÁ J. and CHVÁLA M., 1997: Additions and corrections of the Czech and Slovak list of Tephritidae (Diptera). *Acta Univ. Carol. – Biol.*, 41. (In press)
- KNOZ J., 1997: Ceratopogonidae (Diptera) recorded firstly in the Czech and Slovak Republics. In: VANHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8. *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 77–87.
- KOŠEL V. & LAŠTOVKA P., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Mycetophilidae. In: VANHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8. *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 214–215.
- KOŠEL, V., ZATWARNICKI, T., HORVATH, M., 1997: The first Ephydriidae (Diptera) from cave habitat in Slovakia. *Acta Zool. Univ. Comenianae*, 41. (In press)
- KOZÁNEK, M. & ROLLER, L., 1997: The study of Scatopsid (Diptera: Scatopsidae) communities in some localities in Slovakia. *Biológia Bratislava*. (In press)
- KUBIK Š. & ROHÁČEK J., 1997: New records of Chloropidae (Diptera) from the Czech and Slovak Republics. In: VANHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8. *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 93–95.
- LAPÁČEK V., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Chamaemyiidae. In: VANHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8. *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 227.
- LAUTERER P., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Pipunculidae. In: VANHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8. *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 222–224.
- MACA J., 1997a: New and interesting records of Lonchaeidae (Diptera) from the Czech and Slovak Republics. In: VANHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8. *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 107–110.
- MACA J., 1997b: Faunistic Records from the Czech and Slovak Republics. Diptera: Drosophilidae, Odiniidae. In: VANHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8. *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 227–228.
- MACA J., 1997c: Zajímavé nálezy dvoukřídleho hmyzu skupiny Cyclorrhapha v jižních Čechách. /Interesting findings of Diptera Cyclorrhapha from Southern Bohemia/. *Sbor. Jihočes. Muz. Č. Budějovice, přír. vědy*, 36 (2). (In press)
- MARTINEK V., 1996a: Dvoukřídlý hmyz (Diptera) na lesní louce v Orlických horách v letním a podzimním aspektu. /Dipterous insects (Diptera) in a forest meadow in the Orlické Mts. in summer and fall aspects/. *Lesnictví*, 42 (5): 193–212.

- MARTINEK V., 1996b: Společenstvo vyšších dvoukřídých (Diptera-Brachycera) na mokřadech u řeky Divoké Orlice. /The community of the higher Diptera (Diptera-Brachycera) on wetlands along the Divoká Orlice river/. *Lesnictví*, 42 (11): 518–536.
- MARTINEK V., 1997: The newest finds of some species of the group Acalyptrata (Diptera) in the territory of the Czech and Slovak Republics. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8, *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 111–114.
- MARTINOVSKÝ J., 1997a: Contribution to the knowledge of Scatopsidae (Diptera) from Czech and Slovak Republics, with descriptions of two new species. – *Čas. Slez. Muz. Opava (A)*, 45: 193–210.
- MARTINOVSKÝ J., 1997b: Faunistic Records from the Czech and Slovak Republics. Diptera: Tipulidae, Thaumaleidae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8, *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 212–213.
- MARTINOVSKÝ J. & ŠEVČÍK J., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Keroplatidae, Mycetophilidae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8, *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 213.
- MOCEK B., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Phoridae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8, *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 220–222.
- OLEJNÍČEK J., BARTÁK M., 1996: Diptera: Dolichopodidae. Faunistic records from the Czech Republic – 41. *Klapalekiana*, 32: 129–130.
- OLEJNÍČEK J. & BARTÁK M., 1997: Some faunistically interesting Dolichopodidae (Diptera) from Europe. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8, *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 137–139.
- OLEJNÍČEK J., KOZÁNEK M., 1996: Faunistical notes, Diptera, Dolichopodidae. *Biologia, Bratislava*, 51: 200.
- OLEJNÍČEK J., MOCEK B., 1996: Faunistic records from the Czech Republic – 47. Diptera: Dolichopodidae. *Klapalekiana*, 32: 200.
- ORSZAGH, I., MRÁZ, R., 1996: A vernal aspect of the *Culicoides* fauna of Little Carpathians Mts. (Diptera, Ceratopogonidae). *Entomofauna carpathica*, 8: 141–144.
- POVOLNÝ D., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Sarcophagidae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8, *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 233–234.
- ROHÁČEK J., 1996a: Fourth supplement to the acalyptrate Diptera fauna of the Czech Republic and Slovakia. – *Čas. Slez. Muz. Opava (A)*, 45: 17–28.
- ROHÁČEK J., 1996b: New records of Chloropidae (Diptera) from the Czech Republic and Slovakia. – *Čas. Slez. Muz. Opava (A)*, 45: 97–116.
- ROHÁČEK J., 1996c: Revision of Palaearctic *Stiphrosoma*, including the *Anthomyza laeta*-group (Diptera: Anthomyzidae). *Eur. J. Entomol.*, 93: 89–120.
- ROHÁČEK J., 1996d: Sphaeroceridae (Diptera) of the Czech Republic: corrections and additions to the faunal list, with taxonomical notes. – *Čas. Slez. Muz. Opava (A)*, 44 (1995): 219–240.
- ROHÁČEK J., 1997: *Borboropsis* and *Oldenbergiella* (Diptera, Heleomyzidae) in the Czech and Slovak Republics: faunistic records and notes on their biology. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8, *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 145–147.
- ROZKOŠNÝ R., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Stratiomyidae, Sciomyzidae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8, *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 95: 215.
- ROZKOŠNÝ R., GREGOR F. & PONT A.C., 1997: The European Fanniidae (Diptera). *Acta Sc. Nat. Brno*, 31 (2): 1–80.
- ROZKOŠNÝ R. & VAŇHARA J., 1997: Dvoukřídli (Diptera, Brachycera) Mohelenské hadcové stepi a přilehlého údolí řeky Jihlavy. /Diptera (Brachycera) of the Mohelno serpentine steppe and the adjoining Jihlava river valley/. *Čas. Západohor. Muz. Třebíč*. (In press)
- RUDZINSKI H. G., 1994: Trauermückenfunde aus Nord – Mähren (Diptera: Sciaridae). *Entomol. Probl.*, 25 (2): 11–23.
- RUDZINSKI, H. G., 1996: Beiträge zur Trauermückenfauna der Slowakei (Diptera: Sciaridae). Teil I: Gattungen *Phytosciara* und *Xylosciara*. *Entomol. Probl.*, 27: 57–63.

- SKUHRAVÁ M., 1997: Gall midges (Diptera, Cecidomyiidae) of the Czech and Slovak Republics as members of zoogeographical units in the Palaearctic Region. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8. Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 149–171.
- STARÝ J., 1996a: New records of Blephariceridae (Diptera) from the Czech Republic. *Čas. Slez. Muz. Opava (A)*, 45: 117–118.
- STARÝ J., 1996b: New records of Limoniidae and Pediciidae (Diptera) from the Czech and Slovak Republics. *Čas. Slez. Muz. Opava (A)*, 45: 119–124.
- STARÝ J., 1997a: Faunistic Records from the Czech and Slovak Republics. Diptera: Trichoceridae, Limoniidae, Pediciidae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8. Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 211–212.
- STARÝ J., 1997b: A new *Ula* from eastern Slovakia (Diptera, Pediciidae). *Čas. Slez. Muz. Opava (A)*. (In press)
- STARÝ J., 1998: *Trichocera* (s. str.) *rufulenta* Edwards, 1938, a valid species (Diptera, Trichoceridae). *Studia dipterologica*. (In press)
- STARÝ J. & MARTINOVSKÝ J., 1998: Little-known and new *Trichocera* s. str., related to *T. implicata* Dahl. 1976, with a revised checklist of the Czech and Slovak Trichoceridae (Diptera). *Entomol. probl.* (In press)
- SZADZIEWSKI R., 1984: On synonymy and morphology of some *Culicoides* species (Diptera, Ceratopogonidae). *Pol. Pis. Entomol.*, 53: 559–566.
- SZADZIEWSKI R., 1986: Redescriptions and notes on some Ceratopogonidae (Diptera). *Pol. Pis. Entomol.*, 56: 3–103.
- SZADZIEWSKI R., GILKA W., ANTHON H., 1995: Immature stages of two European species of the subgenus *Melochelca* (Diptera: Ceratopogonidae), with keys to the European subgenera of *Atrichopogon*. *Ent. scand.*, 26: 181–190.
- ŠIFNER F., 1997: Faunistic Records from the Czech and Slovak Republics. Diptera: Scathophagidae. In: VAŇHARA J. & ROZKOŠNÝ R. (eds), *Dipterologica bohemoslovaca*. Vol. 8. Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia, 95: 228–229.
- VAŇHARA J. & ROZKOŠNÝ R., 1997: Czech and Slovak dipterological literature 1986–1995. *Folia Fac. Sci. Univ. Masarykianae Brunensis, Biologia*, 96. (In press)
- ZATWARNICKI T., 1996: New records of Western Palaearctic shore flies (Diptera: Ephydriidae). *Annals Upper Sil. Mus., Entomol.*, 6–7: 107–125.

INDEX

This index includes the names of families, subfamilies, tribes, genera and subgenera.
Synonyms are in italics.

Ablabesmyia (Chironomidae)	35	Allodiopsis (Mycetophilidae)	22
Abrachyglossum (Conopidae)	68	Allognosta (Stratiomyidae)	44
Acanthiophilus (Tephritidae)	71	Allohelea (Ceratopogonidae)	40
Acanthocnema (Scatophagidae)	90	Allolimosina (Sphaeroceridae)	85
Acanthohelea (Ceratopogonidae)	40	Allopiophila (Piophilidae)	69
Acantholeria (Heleomyzidae)	82	Allopsectrocladius (Chironomidae)	37
Acaroletes (Cecidomyiidae)	28	Allotheobaldia (Culicidae)	34
ACARTOPHTHALMIDAE	75	Allotrichoma (Ephydriidae)	88
Acartophthalmus (Acartophthalmidae)	75	Alluaudomyia (Ceratopogonidae)	40
Acenya (Tachinidae)	101	Alsomyia (Tachinidae)	101
ACEMYINI	101	Amalopsis (Pediidae)	15
Acerocnema (Scatophagidae)	90	Amauromyza (Agromyzidae)	76
Acidia (Tephritidae)	71	Amelibaca (Tachinidae)	101
Acinia (Tephritidae)	71	Amerhapha (Cecidomyiidae)	25
Acklandia (Anthomyiidae)	90	Americina (Scatophagidae)	89
Acletoxenus (Drosophilidae)	86	Ametrodiplosis (Cecidomyiidae)	28
Aclonempis (Empididae)	50	Amiota (Drosophilidae)	86
Acnemia (Mycetophilidae)	19	Ammothereva (Therevidae)	46
Acodiplosis (Cecidomyiidae)	28	Amobia (Sarcophagidae)	98
Acricotopus (Chironomidae)	35	AMOBINI	98
Acrocera (Acroceridae)	45	Amphipogon (Piophilidae)	69
ACROCERIDAE	45	Amydroneura (Empididae)	50
Acrometopia (Chamaemyiidae)	73	Anabremia (Cecidomyiidae)	28
Acropsilus (Dolichopodidae)	54	Anaclileia (Mycetophilidae)	19
Acrosanthe (Therevidae)	46	Anacanthothecum (Sarcophagidae)	98
Actenoptera (Piophilidae)	70	Anacrostichus (Empididae)	49
Actia (Tachinidae)	104	Anagnota (Anthomyiidae)	78
Actina (Stratiomyidae)	44	Anapausis (Scatopsidae)	32
Actinoptera (Tephritidae)	71	Anasimyia (Syrphidae)	65
Acutipula (Tipulidae)	16	Anastoechus (Bombyliidae)	46
Acyglossa (Anthomyiidae)	90	Anatella (Mycetophilidae)	22
Adelphomyia (Limoniidae)	11	Anatopynia (Chironomidae)	35
Adia (Anthomyiidae)	90	ANATOPYNIINI	35
Adidocidia (Diadocidiidae)	18	Andrenosoma (Asilidae)	47
Admontia (Tachinidae)	100	ANDRENOSOMATINI	47
Adoxomyia (Stratiomyidae)	45	Androprosopa (Thaumaleidae)	42
Aedes (Culicidae)	34	Anepsiomyia (Dolichopodidae)	54
Aedimorphus (Culicidae)	34	Anevrina (Phoridae)	58
Aenigmatis (Phoridae)	58	Angiometopa (Sarcophagidae)	98
Afrolimonia (Limoniidae)	14	Angioneura (Calliphoridae)	97
Agathomyia (Platypezidae)	57	ANISOPODIDAE	32
Agria (Sarcophagidae)	98	Anisostephus (Cecidomyiidae)	29
Agrolimna (Ephydriidae)	88	Anomoia (Tephritidae)	71
Agromyza (Agromyzidae)	75	Anopheles (Culicidae)	34
AGROMYZIDAE	75	ANOPHELINAE	34
AGROMYZINAE	75	Anthalia (Hybotidae)	52
Achalcus (Dolichopodidae)	54	Anthepiscopus (Empididae)	48
Achanthiptera (Muscidae)	93	Anthomyia (Anthomyiidae)	90
ACHANTHIPTERINAE	93	ANTHOMYIIDAE	90
Achyrolimonia (Limoniidae)	14	Anthomyza (Anthomyzidae)	78
Alimosina (Sphaeroceridae)	85	ANTHOMYZIDAE	78
Allanthalia (Hybotidae)	52	ANTHRACINAE	45
Alliopsis (Anthomyiidae)	90	Anthrax (Bombyliidae)	45
Alloborborus (Sphaeroceridae)	84	<i>Antichaeta</i> (Sciomyzidae)	74
Allocotocera (Mycetophilidae)	19	Anticheta (Sciomyzidae)	74
Allodia (Mycetophilidae)	22	Antichiridium (Cecidomyiidae)	29

Antipalus (Asilidae)	47	Austrolimnophila (Limoniidae)	11
Antiphrisson (Asilidae)	47	Avaritia (Ceratopogonidae)	39
Antlemon (Keroplatidae)	18	AXYMYIIDAE	17
Antlemonopsis (Keroplatidae)	18	Axysta (Ephydriidae)	88
Antocha (Limoniidae)	14	Azana (Mycetophilidae)	19
Anypotacta (Heleomyzidae)	82	Azelia (Muscidae)	93
Aphaniosoma (Chyromyidae)	83	AZELIINI	93
Aphanotrigonum (Chloropidae)	80		
Aphidoletes (Cecidomyiidae)	29	Baccha (Syrphidae)	62
Aphria (Tachinidae)	104	BACCHINI	62
Apiloscatopse (Scatopsidae)	33	Bactromyia (Tachinidae)	101
Apiomyia (Cecidomyiidae)	25	Barraudius (Culicidae)	34
Aplomya (Tachinidae)	101	Basilia (Nycteribiidae)	107
Aplonyx (Cecidomyiidae)	25	Bathycranium (Dolichopodidae)	55
Apodacra (Sarcophagidae)	98	Baumhaueria (Tachinidae)	102
Apolephthisa (Mycetophilidae)	19	Bayeria (Cecidomyiidae)	25
Apotropina (Chloropidae)	80	Bayeriola (Cecidomyiidae)	25
Appendicia (Tachinidae)	103	Bazarella (Psychodidae)	32
Apsectrotanytus (Chironomidae)	35	Beckerias (Pipunculidae)	66
Apteromyia (Sphaeroceridae)	84	Beckerina (Phoridae)	58
Aptilotus (Sphaeroceridae)	84	Belida (Tachinidae)	100
Arachnidomyia (Sarcophagidae)	99	Bellardia (Calliphoridae)	96
Arctocnopa (Limoniidae)	12	Bellieriomima (Sarcophagidae)	99
Arctophila (Syrphidae)	64	Beltranmyia (Ceratopogonidae)	39
Argyra (Dolichopodidae)	55	Bercaea (Sarcophagidae)	99
Argyrandrus (Empididae)	49	Berdeniella (Psychodidae)	32
ARGYRINI	55	Bergenstammia (Empididae)	51
Archilimnophila (Limoniidae)	11	BERIDINAE	44
Arnoldiola (Cecidomyiidae)	25	Beringotipula (Tipulidae)	16
Arthrocnodax (Cecidomyiidae)	29	Beris (Stratiomyidae)	44
Ascelotella (Sarcophagidae)	99	Berkshiria (Stratiomyidae)	45
Aschistonyx (Cecidomyiidae)	29	Bessa (Tachinidae)	100
ASILIDAE	46	Besseria (Tachinidae)	106
ASILINAE	47	Bezzia (Ceratopogonidae)	40
Asilus (Asilidae)	47	Bibio (Bibionidae)	17
Asindulum (Keroplatidae)	18	BIBIONIDAE	17
Asphondylia (Cecidomyiidae)	28	Bicellaria (Hybotidae)	51
ASPHONDYLIIDI	28	Bifronsina (Sphaeroceridae)	85
Aspistes (Scatopsidae)	32	Billaea (Tachinidae)	105
ASPISTINAE	32	Billotia (Phoridae)	58
Asteia (Asteiidae)	79	Bithia (Tachinidae)	104
ASTEIIDAE	79	Blastodiplosis (Cecidomyiidae)	29
ASTEIINAE	79	Blastomyia (Cecidomyiidae)	25
Asynapta (Cecidomyiidae)	25	Blepharicera (Blephariceridae)	17
Asyndetus (Dolichopodidae)	55	BLEPHARICERIDAE	17
ATELESTIDAE	57	Blepharipa (Tachinidae)	102
Atelestus (Atelestidae)	57	Blepharomyia (Tachinidae)	105
ATHERICIDAE	43	Blera (Syrphidae)	65
Atherigona (Muscidae)	94	Blesoxipha (Sarcophagidae)	98
ATHERIGONINI	94	Blondelia (Tachinidae)	100
Atherix (Athericidae)	43	BLONDELIINI	100
Athrycia (Tachinidae)	105	Boettcherella (Sarcophagidae)	99
Athyroglossa (Ephydriidae)	88	Boletina (Mycetophilidae)	20
Atissa (Ephydriidae)	87	Bolitophila (Bolitophilidae)	18
ATISSINI	87	BOLITOPHILIDAE	18
Atrichopogon (Ceratopogonidae)	41	Bolopus (Platypezidae)	58
Atrichops (Athericidae)	43	BOMBYLIIDAE	45
Atrichosema (Cecidomyiidae)	29	BOMBYLIINAE	46
Atylomyia (Tachinidae)	101	BOMBYLIINI	46
Atylostoma (Tachinidae)	104	Bombylisoma (Bombyliidae)	46
Atylotus (Tabanidae)	43	Bombylius (Bombyliidae)	46
Atypophthalmus (Limoniidae)	14	Boophthora (Simuliidae)	42
Aulacigaster (Aulacigastriidae)	79	Borborillus (Sphaeroceridae)	84
AULACIGASTRIDAE	79	BORBOROPSIDINI	82
Aulagromyza (Agromyzidae)	76	Borboropsis (Heleomyzidae)	82
Aulogastromyia (Lauxaniidae)	72	Boreoclytocerus (Psychodidae)	32

Boreoheptagyia (Chironomidae)	35	Camptomyia (Cecidomyiidae)	25
BOREOHEPTAGYIINI	35	Camptoneuromyia (Cecidomyiidae)	25
Borophaga (Phoridae)	58	Campylocheta (Tachinidae)	105
Botanophila (Anthomyiidae)	90	Campylomyza (Cecidomyiidae)	25
Bothria (Tachinidae)	102	Camtochaeta (Sciaridae)	23
Bradysia (Sciaridae)	23	CANTHYLOSCELIDAE	33
Brachicoma (Sarcophagidae)	98	Carcelia (Tachinidae)	101
Brachicheta (Tachinidae)	102	Cardiocladius (Chironomidae)	36
Brachycampta (Mycetophilidae)	22	Caricea (Muscidae)	96
Brachydiplosis (Cecidomyiidae)	29	CARNIDAE	79
Brachylimnophila (Limoniidae)	11	Carnus (Carnidae)	79
BRACHYMERINI	104	Catagonia (Tachinidae)	101
Brachymyia (Syrphidae)	65	Catharosia (Tachinidae)	106
Brachyopa (Syrphidae)	64	CATHAROSIINI	106
Brachypalpoides (Syrphidae)	65	Catocha (Cecidomyiidae)	23
Brachypalpus (Syrphidae)	65	Cecidomyia (Cecidomyiidae)	29
Brachypeza (Mycetophilidae)	22	CECIDOMYIIDAE	25
Brachypogon (Ceratopogonidae)	40	CECIDOMYIID	28
Brachystoma (Empididae)	51	CECIDOMYIINAE	25
BRACHYSTOMATINAE	51	Centorisoma (Chloropidae)	81
Braula (Braulidae)	79	Centrophlebomyia (Piophilidae)	70
BRAULIDAE	79	Cephalia (Otitidae)	70
Bremia (Cecidomyiidae)	29	Cephalomyza (Agromyzidae)	76
Bremiola (Cecidomyiidae)	25	Cephalops (Pipunculidae)	66
Brevicornu (Mycetophilidae)	22	Cephalosphaera (Pipunculidae)	66
Brillia (Chironomidae)	36	Cephenemyia (Oestridae)	107
Brontaea (Muscidae)	95	CEPHENEMYIINAE	107
Brullaea (Tachinidae)	106	Cerajocera (Tephritidae)	71
Bryophaenocladius (Chironomidae)	36	Ceranthia (Tachinidae)	104
Buhriella (Cecidomyiidae)	29	Ceratoculicoides (Ceratopogonidae)	40
Buchonomyia (Chironomidae)	35	Ceratopogon (Ceratopogonidae)	40
BUCHONOMYIINAE	35	CERATOPOGONIDAE	39
Buquetia (Tachinidae)	101	CERATOPOGONINAE	39
Butomomyza (Agromyzidae)	76	CERATOPOGONINI	40
		Cerdistus (Asilidae)	47
Cacoxenus (Drosophilidae)	86	Ceriana (Syrphidae)	65
Cadurcia (Tachinidae)	102	CERIODINI	65
Cadurciella (Tachinidae)	101	Cerodontha (Agromyzidae)	76
Calamoncosis (Chloropidae)	80	Ceromasia (Tachinidae)	102
Caliprobola (Syrphidae)	65	Ceromya (Tachinidae)	104
Callicera (Syrphidae)	63	Cerotelion (Keroplastidae)	18
CALLICERINI	63	Ceroxys (Otitidae)	70
Calliopum (Lauxaniidae)	72	Cetema (Chloropidae)	81
Calliphora (Calliphoridae)	96	Chaetoapnea (Ephydriidae)	89
CALLIPHORIDAE	96	Chaetocladius (Chironomidae)	36
CALLIPHORINAE	96	Chaetolonchaea (Lonchaeidae)	69
Callixania (Lauxaniidae)	72	Chaetomus (Heleomyzidae)	82
Callomyia (Platypezidae)	58	Chaetopleurophora (Phoridae)	58
CALLOMYIINAE	57	Chaetopodella (Sphaeroceridae)	84
Calloptereella (Diatatidae)	87	Chaetoptilia (Tachinidae)	105
Calobata (Micropezidae)	67	Chaetorellia (Tephritidae)	71
Calobatella (Micropezidae)	67	Chaetosa (Scatophagidae)	90
CALOBATINAE	67	Chaetosciara (Sciaridae)	23
Caloforecipomyia (Ceratopogonidae)	41	Chaetostomella (Tephritidae)	71
Calycomyza (Agromyzidae)	76	CHALARINAE	66
Calythea (Anthomyiidae)	91	Chalarus (Pipunculidae)	66
Camarota (Chloropidae)	81	Chalcosyrphus (Syrphidae)	65
Camilla (Camillidae)	86	Chamaedipsia (Empididae)	51
CAMILLIDAE	86	Chamaemyia (Chamaemyiidae)	73
Campiglossa (Tephritidae)	71	CHAMAEMYIIDAE	73
Campichoeta (Diatatidae)	87	Chamaepsila (Psilidae)	67
CAMPICHOETINAE	87	Chamaesyrphus (Syrphidae)	63
CAMPSICNEMINI	55	CHAOBORIDAE	33
Campsicnemus (Dolichopodidae)	55	CHAOBORINAE	33
Camptocladius (Chironomidae)	36	CHAOBORINI	33
Camptochironomus (Chironomidae)	37	Chaoborus (Chaoboridae)	33

Cheilosia (Syrphidae)	63	Cnodacophora (Micropezidae)	67
CHEILOSINI	63	Coboldia (Scatopsidae)	33
Cheilotrichia (Limoniidae)	12	Coelophthiria (Mycetophilidae)	20
Chelifera (Empididae)	51	Coelosia (Mycetophilidae)	20
Chelipoda (Empididae)	51	COELOTANYPODINI	34
Chersodromia (Hybotidae)	53	Coenia (Ephydriidae)	89
Chetogena (Tachinidae)	100	Coenomyia (Coenomyiidae)	43
Chiastochacta (Anthomyiidae)	91	COENOMYIIDAE	43
Chionea (Limoniidae)	12	Coenosia (Muscidae)	96
CHIONEINAE	12	COENOSIINAE	95
CHIRONOMIDAE	34	COENOSIINI	96
CHIRONOMINAE	37	Cochliarium (Scatophagidae)	90
CHIRONOMINI	37	Collimosina (Sphaoceridae)	85
Chironomus (Chironomidae)	37	Colobaea (Sciomyzidae)	73
Chirosia (Anthomyiidae)	91	Colobostema (Scatopsidae)	33
Chloromyia (Stratiomyidae)	44	Columbiella (Clusiidae)	75
CHLOROPIDAE	80	Compsilura (Tachinidae)	100
CHLOROPINAE	81	Compsobata (Micropezidae)	67
Chlorops (Chloropidae)	81	Conchapelopia (Chironomidae)	35
Chodopsycha (Psychodidae)	31	Conicera (Phoridae)	58
Choerades (Asilidae)	46	Conioscinella (Chloropidae)	80
Chorisops (Stratiomyidae)	44	Coniosternum (Scatophagidae)	90
Chromatomyia (Agromyzidae)	77	Conogaster (Tachinidae)	100
Chrysogaster (Syrphidae)	64	CONOPHORINI	46
CHRYSOGASTERINI	64	Conophorus (Bombyliidae)	46
CHRYSOMYINAE	97	CONOPIDAE	68
Chrysopilus (Rhagionidae)	43	CONOPINAE	68
Chrysops (Tabanidae)	43	Conops (Conopidae)	68
CHRYSOPSINAE	43	Contarinia (Cecidomyiidae)	29
Chrysotimus (Dolichopodidae)	54	Cookella (Scatopsidae)	33
CHRYSOTOXINI	62	Coproica (Sphaoceridae)	84
Chrysotoxum (Syrphidae)	62	Copromyza (Sphaoceridae)	84
Chrysotus (Dolichopodidae)	55	COPROMYZINAE	84
Chyliza (Psilidae)	67	Copropsychoda (Psychodidae)	31
Chylizosoma (Scatophagidae)	89	Coptophlebia (Empididae)	49
Chymomyza (Drosophilidae)	86	Coquillettia (Culicidae)	34
Chyromya (Chyromyidae)	83	Coquillettomyia (Cecidomyiidae)	30
CHYROMYIDAE	83	Cordilura (Scatophagidae)	89
Cinochira (Tachinidae)	106	CORDILURINAE	89
Cistogaster (Tachinidae)	106	Cordyla (Mycetophilidae)	22
Cladopelma (Chironomidae)	38	Coremacera (Sciomyzidae)	74
Cladotanytarsus (Chironomidae)	39	Coreozelia (Limoniidae)	11
Clairvillia (Tachinidae)	106	Corynocera (Chironomidae)	39
Clanoneurum (Ephydriidae)	87	Corynoneura (Chironomidae)	36
Clastobasis (Mycetophilidae)	20	Corynoptera (Sciaridae)	23
Clausicella (Tachinidae)	104	Cosmetopus (Scatophagidae)	90
Cleigastria (Scatophagidae)	90	Cranciobia (Cecidomyiidae)	25
Clemelis (Tachinidae)	102	Crassivenula (Chloropidae)	80
Cleonice (Tachinidae)	103	Crataerina (Hippoboscidae)	107
Clitocera (Empididae)	51	Cratyna (Sciaridae)	24
CLINOCERINAE	51	Cremifania (Cremifaniidae)	73
Clinodiplosis (Cecidomyiidae)	29	CREMIFANIIDAE	73
Clinotanytus (Chironomidae)	34	Cricotopus (Chironomidae)	36
Clitopisa (Bolitophilidae)	18	Criorhina (Syrphidae)	65
Clitormia (Therevidae)	46	Crossopalpus (Hybotidae)	53
Clitellaria (Stratiomyidae)	45	Crumomyia (Sphaoceridae)	84
CLITELLARIINAE	45	Crunobia (Pediidae)	15
Clusia (Clusiidae)	75	Cryptaciura (Tephritidae)	72
Clusiaria (Clusiidae)	75	Crypteria (Limoniidae)	12
CLUSIIDAE	75	Cryptochironomus (Chironomidae)	38
Clusiodes (Clusiidae)	75	Cryptocladopehna (Chironomidae)	38
Clytiomyia (Tachinidae)	106	Cryptonevra (Chloropidae)	82
Clytocerus (Psychodidae)	32	Cryptotendipes (Chironomidae)	38
Cnemacantha (Lauxaniidae)	72	Ctenodasyptigia (Sarcophagidae)	99
Cnemoncosis (Tipulidae)	17	Ctenophora (Tipulidae)	17
Cnestrum (Ephydriidae)	87	CTENOPHORINAE	17

CTENOPHORINI	17	DIASTATIDAE	87
Ctenosciara (Sciaridae)	24	Diazosma (Trichoceridae)	11
Culex (Culicidae)	34	Dicerura (Cecidomyiidae)	25
Culicella (Culicidae)	34	Diclasiope (Ephydriidae)	88
CULICIDAE	34	Dicraeus (Chloropidae)	80
CULICINAE	34	Dicranomyia (Limoniidae)	14
Culicoides (Ceratopogonidae)	39	Dicranoptycha (Limoniidae)	14
CULICOIDINI	39	Dicranota (Pediidae)	15
Culiseta (Culicidae)	34	Dicrotendipes (Chironomidae)	38
Curranella (Sarcophagidae)	99	Dicryptoscena (Ceratopogonidae)	41
CURTONOTIDAE	86	Dictenidia (Tipulidae)	17
Curtonotum (Curtonotidae)	86	Dictya (Sciomyzidae)	74
Cylindromyia (Tachinidae)	106	Didea (Syrphidae)	61
CYLINDROMYIINI	106	Didymomyia (Cecidomyiidae)	26
Cylindrotoma (Cylindrotomidae)	15	Digitiventra (Sarcophagidae)	100
CYLINDROTOMIDAE	15	Dichaeta (Ephydriidae)	88
Cymomyia (Mycetophilidae)	19	Dichaetophora (Sciomyzidae)	74
Cynomyia (Calliphoridae)	97	Dichetophora (Sciomyzidae)	74
Cyrtophleba (Tachinidae)	105	Dichodiplosis (Cecidomyiidae)	30
Cyrtopogon (Asilidae)	47	Dilophus (Bibionidae)	17
Cystiphora (Cecidomyiidae)	25	Dinera (Tachinidae)	105
Cyzenis (Tachinidae)	102	Dioctria (Asilidae)	47
		DIOCTRIINI	47
DACTYLOLABINAE	11	Diodaulus (Cecidomyiidae)	30
Dactylolabis (Limoniidae)	11	Diogma (Cylindrotomidae)	15
Dahlmosina (Sphaeroceridae)	85	Dionaea (Tachinidae)	106
Dalmanina (Conopidae)	68	Dioxyna (Tephritidae)	71
DALMANNINAE	68	Diplocladius (Chironomidae)	36
Dasineura (Cecidomyiidae)	25	Diploneura (Phoridae)	58
DASIOPINAE	68	Diplostichus (Tachinidae)	100
Dasiops (Lonchaeidae)	68	Diplotoxa (Chloropidae)	82
Dasyhelea (Ceratopogonidae)	41	Diplotoxoides (Chloropidae)	82
DASYHELEINAE	41	Discachaeta (Sarcophagidae)	99
Dasymolophilus (Limoniidae)	13	Discobola (Limoniidae)	14
Dasya (Psilidae)	68	Discocerina (Ephydriidae)	88
Dasyopa (Chloropidae)	80	DISCOCERININI	88
Dasyphora (Muscidae)	94	Discomyza (Ephydriidae)	87
Dasyopogon (Asilidae)	47	DISCOMYZINAE	87
DASYPOGONINAE	47	DISCOMYZINI	87
DASYPOGONINI	47	DISCHISTINI	46
Dasyrhyphus (Syrphidae)	60	Ditacniella (Sciomyzidae)	73
Decembrina (Sciaridae)	24	Dithryca (Tephritidae)	71
Delia (Anthomyiidae)	91	DITHRYCINI	71
Delina (Sarcophagidae)	90	Ditomyia (Ditomyiidae)	18
DELININAE	90	DITOMYIIDAE	18
Demeijerea (Chironomidae)	38	Ditrichophora (Ephydriidae)	88
Demicryptochironomus (Chironomidae)	38	Dixa (Dixidae)	33
Demotus (Tachinidae)	104	Dixella (Dixidae)	33
Dendrotipula (Tipulidae)	16	DIXIDAE	33
Dentilimosina (Sphaeroceridae)	85	Dizygomyza (Agromyzidae)	76
Desmometopa (Milichiidae)	80	Docosia (Mycetophilidae)	20
Dexia (Tachinidae)	105	Dohrniphora (Phoridae)	58
DEXIINAE	105	Dolichocephala (Empididae)	51
DEXINI	105	Dolichocheila (Tipulidae)	15
Dexiosoma (Tachinidae)	105	DOLICHOPEZINAE	15
Diadocidia (Diadocidiidae)	18	DOLICHOPEZINI	15
DIADOCIDIIDAE	18	Dolichophorus (Dolichopodidae)	57
DIACHLORINI	44	DOLICHOPODIDAE	54
Dialineura (Therevidae)	46	DOLICHOPODINAE	55
Diamesa (Chironomidae)	35	DOLICHOPODINI	56
DIAMESINAE	35	Dolichopus (Dolichopodidae)	56
DIAMESINI	35	Dolichosciara (Sciaridae)	24
DIAPHORINAE	55	Doros (Syrphidae)	61
DIAPHORINI	55	Dorsilopa (Drosophilidae)	86
Diaphorus (Dolichopodidae)	55	Dorycera (Otitidae)	70
Diastata (Diastatidae)	87	Dorylomorpha (Pipunculidae)	66

Dorylomyia (Pipunculidae)	66	EPHYDRINAE	89
Dorylomyza (Pipunculidae)	66	EPHYDRINI	89
DRAPETINI	53	Epicampocera (Tachinidae)	101
Drapetis (Hybotidae)	53	Epicypta (Mycetophilidae)	20
Dratnalia (Chironomidae)	36	Epidapus (Sciaridae)	24
Drino (Tachinidae)	101	Epichlorops (Chloropidae)	82
Drisina (Cecidomyiidae)	30	Epiphragma (Limoniidae)	11
Drosophila (Drosophilidae)	86	Epistrophe (Syrphidae)	61
DROSOPHILIDAE	86	Epistrophella (Syrphidae)	61
DROSOPHILINAE	86	Episyrphus (Syrphidae)	61
Drymeia (Muscidae)	94	Epoicocladus (Chironomidae)	36
Dryodromia (Empididae)	51	Erax (Asilidae)	47
Dryomyia (Cecidomyiidae)	26	Eribolus (Chloropidae)	80
Dryomyza (Dryomyzidae)	73	Eriocera (Limoniidae)	11
DRYOMYZIDAE	73	Eriocnopa (Limoniidae)	12
Duckhousiella (Psychodidae)	31	Erioptera (Limoniidae)	12
Dufouria (Tachinidae)	105	Eriothrix (Tachinidae)	105
DUFOURIINI	105	Eriozona (Syrphidae)	61
Dynatosoma (Mycetophilidae)	20	ERISTALINI	65
Dysaletria (Hybotidae)	52	Eristalinus (Syrphidae)	65
Dysmachus (Asilidae)	47	Eristalis (Syrphidae)	65
Dziedzickia (Mycetophilidae)	20	Ernestia (Tachinidae)	103
		ERNESTIINI	103
Earomyia (Lonchaeidae)	69	Erycia (Tachinidae)	101
EAROMYIINI	69	ERYCIINI	101
Eccoptomera (Heleomyzidae)	83	Erycilla (Tachinidae)	102
Ectaetia (Scatopsidae)	32	Erythroceria (Tachinidae)	102
ECTAETIINAE	32	Estheria (Tachinidae)	105
Ectinocera (Sciomyzidae)	74	Ethilla (Tachinidae)	101
Ectophasia (Tachinidae)	106	ETHILLINI	101
Ectrepesthoneura (Mycetophilidae)	20	Euaresta (Tephritidae)	71
Eggisops (Calliphoridae)	97	Eucelidia (Empididae)	51
Eginia (Muscidae)	94	Eudactylocadius (Chironomidae)	36
EGINIINI	94	Eudasyphora (Muscidae)	94
Egle (Anthomyiidae)	91	Eudorylas (Pipunculidae)	66
Eclthistus (Asilidae)	47	Euempis (Empididae)	49
Einfeldia (Chironomidae)	38	Eukiefferiella (Chironomidae)	36
Elachiptera (Chloropidae)	80	Eulabidogaster (Tachinidae)	106
Elachisoma (Sphaeroceridae)	84	Euleia (Tephritidae)	72
Elaphropeza (Hybotidae)	53	Eulimosina (Sphaeroceridae)	85
Elephantomyia (Limoniidae)	14	Eumea (Tachinidae)	102
Elfia (Tachinidae)	104	EUMERINI	64
Elgiva (Sciomyzidae)	74	Eumerus (Syrphidae)	64
Elizeta (Tachinidae)	106	Eumetopiella (Sarcophagidae)	98
Elliptera (Limoniidae)	14	Euorthocladus (Chironomidae)	36
Ellipteroides (Limoniidae)	12	Eupachygaster (Stratiomyidae)	45
Eloceria (Tachinidae)	103	Eupeodes (Syrphidae)	61
Elodia (Tachinidae)	102	Euphranta (Tephritidae)	71
Elocophila (Limoniidae)	11	EUPHRANTINI	71
Elomya (Tachinidae)	106	Euphylidorea (Limoniidae)	11
Emmesomyia (Anthomyiidae)	91	Euprojoannisia (Ceratopogonidae)	41
Emodotipula (Tipulidae)	16	Euptilostena (Limoniidae)	12
Empeda (Limoniidae)	12	Eurina (Chloropidae)	82
EMPIDIDAE	48	Eurithia (Tachinidae)	103
EMPIDINAE	48	Eurygnathomyia (Pallopteridae)	69
EMPIDINI	49	EURYGNATHOMYIINAE	69
Empis (Empididae)	49	Eurychaeta (Calliphoridae)	97
Endochironomus (Chironomidae)	38	Eurysthaca (Tachinidae)	102
Endopsylla (Cecidomyiidae)	30	Eusapromyza (Lauxaniidae)	72
ENLININAE	54	Eusimulium (Simuliidae)	42
Enneastigma (Anthomyiidae)	91	Eustalomyia (Anthomyiidae)	91
Ensina (Tephritidae)	71	Eutaenionotum (Ephydriidae)	89
Entomophaga (Tachinidae)	104	EUTHERINI	106
Eoseristalis (Syrphidae)	65	Euthycera (Sciomyzidae)	74
Ephydra (Ephydriidae)	89	Euthyneura (Hybotidae)	52
EPHYDRIDAE	87	Eutolmus (Asilidae)	47

Eutonia (Limoniidae)	11	Greenomyia (Mycetophilidae)	20
Eutricheta (Anthomyiidae)	91	Grzegorzekia (Mycetophilidae)	20
Eutropha (Chloropidae)	82	Guttipelopia (Chironomidae)	35
Euxesta (Ulidiidae)	70	Gymnoclasiopa (Ephydriidae)	88
Exechia (Mycetophilidae)	22	Gymnocheta (Tachinidae)	103
EXECHIINI	22	Gymnochiromyia (Chyromyidae)	84
Exechiopsis (Mycetophilidae)	22	Gymnomera (Scatophagidae)	90
Exhyalanthrax (Bombyliidae)	46	Gymnometriocnemus (Chironomidae)	36
Exodontha (Stratiomyidae)	44	Gymnomus (Heleomyzidae)	83
Exoprosopa (Bombyliidae)	46	GYMNOMYZINAE	88
EXOPROSOPINAE	45	GYMNOMYZINI	88
EXOPROSOPINI	46	Gymnophora (Phoridae)	58
Exorista (Tachinidae)	100	Gymnoptera (Phoridae)	58
EXORISTINAE	100	Gymnosoma (Tachinidae)	106
EXORISTINI	100		
		Hackmanina (Sphaeroceridae)	85
Fagisyrphus (Syrphidae)	61	Hadrobremia (Cecidomyiidae)	30
Fannia (Fanniidae)	93	Haematobia (Muscidae)	94
FANNIIDAE	93	Haematobosca (Muscidae)	94
Fausta (Tachinidae)	103	Haematopota (Tabanidae)	44
Feltiella (Cecidomyiidae)	30	HAEMATOPOTINI	44
Ferdinandea (Syrphidae)	63	Halidayina (Sphaeroceridae)	84
Ferneiella (Scatopsidae)	33	Haloscatella (Ephydriidae)	89
Feuerborniella (Psychodidae)	31	Hammerschmidtia (Syrphidae)	64
Fiebrigella (Chloropidae)	81	Hapleginella (Chloropidae)	80
Finlaya (Culicidae)	34	Haplodiplosis (Cecidomyiidae)	30
Fleuria (Chironomidae)	38	Harmandia (Cecidomyiidae)	30
Forcipomyia (Ceratopogonidae)	41	Harnischia (Chironomidae)	38
FORCIPOMYIINAE	41	Harrisomyia (Cecidomyiidae)	27
Frendelia (Lauxaniidae)	72	Hartigiola (Cecidomyiidae)	27
Freræa (Tachinidae)	105	Hebecnema (Muscidae)	95
Frontina (Tachinidae)	102	Hebia (Tachinidae)	102
		HECAMEDINI	88
Gaedia (Tachinidae)	102	Hecamedoides (Ephydriidae)	88
Galiomyza (Agromyzidae)	77	Heleniella (Chironomidae)	36
Gampsocera (Chloropidae)	80	Heleodromia (Empididae)	51
GASTEROPHILIDAE	106	Heleomyza (Heleomyzidae)	82
Gasterophilus (Gasterophilidae)	106	HELEOMYZIDAE	82
Gastrolepta (Tachinidae)	100	HELEOMYZINAE	82
Gaurax (Chloropidae)	80	HELEOMYZINI	82
Geocrypta (Cecidomyiidae)	26	HELICOBOSCINAE	97
Geomyza (Opomyzidae)	78	Helicophagella (Sarcophagidae)	99
Gephyraulius (Cecidomyiidae)	26	Helina (Muscidae)	95
Germaria (Tachinidae)	103	Helius (Limoniidae)	14
Gigalimosina (Sphaeroceridae)	84	Helophilus (Syrphidae)	65
Giraudiella (Cecidomyiidae)	27	Hemerodromia (Empididae)	51
Gitona (Drosophilidae)	86	HEMERODROMIINAE	51
GITONINI	86	Hemeromyia (Carnidae)	79
Glabellula (Mythicomyiidae)	45	Hemimacquantia (Tachinidae)	100
Glaucops (Tabanidae)	43	Hemineurina (Sciaridae)	24
Glochina (Limoniidae)	14	Hemipenthes (Bombyliidae)	45
Gloma (Empididae)	48	Hemyda (Tachinidae)	106
Gluhhovia (Ceratopogonidae)	41	Hendelia (Clusiidae)	75
Glyptotendipes (Chironomidae)	38	Heptatoma (Tabanidae)	44
Gnophomyia (Limoniidae)	12	Hercostomus (Dolichopodidae)	56
Gnoriste (Mycetophilidae)	20	Herina (Otitidae)	70
GNORISTINI	19	Heringia (Syrphidae)	62
Gonarticus (Scatophagidae)	90	Heringina (Tephritidae)	71
Gonatherus (Scatophagidae)	90	Herniosina (Sphaeroceridae)	84
Gonempeda (Limoniidae)	12	HESPERINIDAE	17
Gonia (Tachinidae)	102	Hesperinus (Hesperinidae)	17
GONIINI	102	Heteromeringia (Clusiidae)	75
Goniocera (Tachinidae)	104	Heteromyza (Heleomyzidae)	83
Gonioneura (Sphaeroceridae)	84	HETEROMYZINAE	83
Gonomyia (Limoniidae)	12	Heteronychia (Sarcophagidae)	99
Graphomya (Muscidae)	95	Heteropeza (Cecidomyiidae)	25

Heterostylodes (Anthomyiidae)	91	Ilione (Sciomyzidae)	74
Heterotanytarsus (Chironomidae)	36	Ilisia (Limoniidae)	13
Heterotrissocladius (Chironomidae)	36	Ilythea (Ephydriidae)	88
Hexamitocera (Scatophagidae)	90	ILYTHEINAE	88
Hexatoma (Limoniidae)	11	ILYTHEINI	88
Hexomyza (Agromyzidae)	75	Incertella (Chloropidae)	81
Hilara (Empididae)	48	Insulomyia (Sphaeroceridae)	86
Hilarella (Sarcophagidae)	98	Ischiolepta (Sphaeroceridae)	84
Hilarimorpha (Hilarimorphidae)	46	Ischyrosyrphus (Syrphidae)	61
HILARIMORPHIDAE	46	Isocladius (Chironomidae)	36
HILARINI	48	Isohelea (Ceratopogonidae)	40
Hippobosca (Hippoboscidae)	107	Isoneuromyia (Keroplastidae)	18
HIPPOBOSCIDAE	107	ISOPOGONINI	47
HIPPOBOSCINAE	107	Istocheta (Tachinidae)	100
Hirtodrosophila (Drosophilidae)	87	Iteaphila (Empididae)	48
Holoclera (Empididae)	50	Iteomyia (Cecidomyiidae)	27
Holoneurus (Cecidomyiidae)	25		
Holoplagia (Scatopsidae)	33	Jaapiella (Cecidomyiidae)	27
Holopogon (Asilidae)	47	Janetia (Cecidomyiidae)	27
Holotanypus (Chironomidae)	34	Janetiella (Cecidomyiidae)	27
Homalura (Chloropidae)	82	Jantia (Sarcophagidae)	99
Homobezzia (Ceratopogonidae)	41	Jassidophaga (Pipunculidae)	66
Homoneura (Lauxaniidae)	72	JOHNSONIINI	99
HOMONEURINAE	72	Jovamormia (Psychodidae)	31
Hoplolabis (Limoniidae)	12	Jungiella (Psychodidae)	31
Huebneria (Tachinidae)	101		
Hyadina (Ephydriidae)	88	Kaltenbachiola (Cecidomyiidae)	27
HYADININI	88	KEROPLATIDAE	18
Hyalurgus (Tachinidae)	103	Keroplatus (Keroplastidae)	18
Hybolasiptera (Cecidomyiidae)	25	Kesselimyia (Platypezidae)	58
Hybomitra (Tabanidae)	43	Kiefferia (Cecidomyiidae)	28
Hybos (Hybotidae)	52	Kiefferulus (Chironomidae)	38
HYBOTIDAE	51	Kimosina (Sphaeroceridae)	85
HYBOTINAE	52	Klugia (Tachinidae)	105
Hydrellia (Ephydriidae)	87	Knutsonia (Sciomyzidae)	74
HYDRELLIINAE	87	Kolenohela (Ceratopogonidae)	40
HYDRELLIINI	87	Kowarzia (Empididae)	51
Hydrobaenus (Chironomidae)	36	Kramerea (Sarcophagidae)	99
Hydromya (Sciomyzidae)	74	Krameromyia (Sarcophagidae)	99
Hydromyza (Scatophagidae)	90	Krenopelopia (Chironomidae)	35
HYDROMYZINAE	90	Krenosmittia (Chironomidae)	36
Hydrophoria (Anthomyiidae)	91	Kritempis (Empididae)	49
HYDROPHORINAE	54		
HYDROPHORINI	54	Labigastera (Tachinidae)	106
Hydrophorus (Dolichopodidae)	54	Lamprochromus (Dolichopodidae)	55
Hydrotaea (Muscidae)	94	Lamproscatella (Ephydriidae)	89
Hygroceleuthus (Dolichopodidae)	56	Laphria (Asilidae)	47
Hygrodiplosis (Cecidomyiidae)	30	LAPHRIINAE	46
Hylemya (Anthomyiidae)	91	LAPHRIINI	46
Hylemyza (Anthomyiidae)	92	Laphystia (Asilidae)	47
Hyleorus (Tachinidae)	105	LAPHYSTIINAE	47
Hyperoscelis (Canthyluscelidae)	33	Lapposyrphus (Syrphidae)	61
Hypocera (Phoridae)	58	Larsia (Chironomidae)	35
Hypoderma (Hypodermatidae)	107	Lasiambia (Chloropidae)	81
HYPODERMATIDAE	107	Lasiohelea (Ceratopogonidae)	42
HYPODERMATINAE	107	Lasiomna (Anthomyiidae)	92
Hypophyllus (Dolichopodidae)	56	Lasiopa (Stratiomyidae)	45
Hyporites (Anthomyiidae)	92	Lasiopleura (Chloropidae)	80
		Lasiopogon (Asilidae)	47
Ibisia (Athericidae)	43	Lasioptera (Cecidomyiidae)	25
Icosta (Hippoboscidae)	107	LASIOPTERIDI	25
Ictericoides (Tephritidae)	71	LASIOPTERINI	25
Icteromyza (Agromyzidae)	77	Lasiosina (Chloropidae)	82
Idiocera (Limoniidae)	12	Lathyromyza (Cecidomyiidae)	27
Idioptera (Limoniidae)	11	Lathyrrophthalmus (Syrphidae)	65
Idiopyga (Limoniidae)	14	Lauterborniella (Chironomidae)	38

Lauxania (Lauxaniidae)	72	Liriomyza (Agromyzidae)	77
LAUXANIIDAE	72	Lispe (Muscidae)	96
LAUXANIINAE	72	Lispocephala (Muscidae)	96
Lecanipa (Tachinidae)	100	Lissempis (Empididae)	50
Leia (Mycetophilidae)	20	Lithophasia (Tachinidae)	106
LEIINI	20	Lobochironomus (Chironomidae)	38
Leiomyza (Asteiidae)	79	Loewia (Tachinidae)	103
Leiophora (Tachinidae)	100	Loewiola (Cecidomyiidae)	30
Lejogaster (Syrphidae)	64	Logima (Psychodidae)	31
Lejops (Syrphidae)	65	Lomachantha (Tachinidae)	100
Lejota (Syrphidae)	65	Lomatia (Bombyliidae)	45
Lengersdorfia (Sciaridae)	24	LOMATIINAE	45
Leopoldius (Conopidae)	68	Lonchaea (Lonchaeidae)	69
Lepimormia (Psychodidae)	31	LONCHAEIDAE	68
Lepiseodina (Psychodidae)	31	LONCHAEINAE	69
Leptarthrus (Asilidae)	47	LONCHAEINI	69
Leptempis (Empididae)	49	Lonchoptera (Lonchopteridae)	60
Leptocera (Sphaeroceridae)	84	LONCHOPTERIDAE	60
Leptogaster (Asilidae)	47	Lophomyidium (Ceratopogonidae)	41
LEPTOGASTRINAE	47	Lophosceles (Muscidae)	95
Leptometa (Milichiidae)	80	Lophosia (Tachinidae)	106
Leptomorphus (Mycetophilidae)	19	Lordiphosa (Drosophilidae)	87
Leptopa (Scatophagidae)	89	Lotobia (Sphaeroceridae)	84
Leptopeza (Hybotidae)	52	Lotophila (Sphaeroceridae)	84
Leptosciarella (Sciaridae)	24	Loxocera (Psilidae)	68
Leriola (Heleomyzidae)	83	Lucilia (Calliphoridae)	97
Leskia (Tachinidae)	104	LUCILIINAE	97
LESKIINI	104	Ludicia (Pediidae)	15
Lestodiplosis (Cecidomyiidae)	30	Lunaria (Limoniidae)	12
Lestremia (Cecidomyiidae)	25	Lunatipula (Tipulidae)	16
LESTREMIINAE	25	Lundstroemiella (Empididae)	50
Leucophenga (Drosophilidae)	86	Lyciella (Lauxaniidae)	72
Leucophora (Anthomyiidae)	92	Lycomya (Mycetophilidae)	19
Leucopis (Chamaemyiidae)	73	Lycoriella (Sciaridae)	24
Leucopomyia (Chamaemyiidae)	73	Lydella (Tachinidae)	101
Leucostoma (Tachinidae)	106	Lydina (Tachinidae)	103
LEUCOSTOMATINI	106	Lypha (Tachinidae)	103
Leucozona (Syrphidae)	61	Lytogaster (Ephydriidae)	88
Liancalus (Dolichopodidae)	54		
Libnotes (Limoniidae)	14	Macquartia (Tachinidae)	104
Ligeria (Tachinidae)	100	MACQUARTIINI	104
Ligeriella (Tachinidae)	100	Macrocera (Macroceridae)	18
Limnellia (Ephydriidae)	89	MACROCERIDAE	18
Limnia (Sciomyzidae)	74	Macrodiplosis (Cecidomyiidae)	30
Limnophila (Limoniidae)	11	Macrolabis (Cecidomyiidae)	27
LIMNOPHILINAE	11	Macronychia (Sarcophagidae)	98
Limnophora (Muscidae)	96	MACRONYCHIIINAE	98
LIMNOPHORINI	95	Macropelopia (Chironomidae)	35
Limnophyes (Chironomidae)	36	MACROPELOPIINI	35
Limnospila (Muscidae)	96	Macropeza (Ceratopogonidae)	40
Limonia (Limoniidae)	14	Macroprosopa (Tachinidae)	104
LIMONIIDAE	11	Macrorchis (Muscidae)	96
LIMONIINAE	14	Macrorrhyncha (Keroplastidae)	18
Limosina (Sphaeroceridae)	85	Madiza (Milichiidae)	80
LIMOSININAE	84	MADIZINAE	80
Lindneromyia (Platypezidae)	58	Machimus (Asilidae)	48
Linnaemya (Tachinidae)	103	Maillotia (Culicidae)	34
LINNAEMYINI	103	Mallochohelea (Ceratopogonidae)	40
Liopiophila (Piophilidae)	69	Mallota (Syrphidae)	65
Liopygia (Sarcophagidae)	99	Mansonia (Culicidae)	34
Liosarcophaga (Sarcophagidae)	99	Masicera (Tachinidae)	102
Lipara (Chloropidae)	81	Masistylum (Tachinidae)	102
Liponeura (Blephariceridae)	17	Massalongia (Cecidomyiidae)	30
Lipoptena (Hippoboscidae)	107	Mayetiola (Cecidomyiidae)	27
LIPOPTENINAE	107	Medetera (Dolichopodidae)	57
Lipsothrix (Limoniidae)	14	MEDETERINAE	57

MEDETERINI	57	Micromorphus (Dolichopodidae)	54
Medina (Tachinidae)	100	Micromya (Cecidomyiidae)	25
Mediotipula (Tipulidae)	16	Micropeza (Micropezidae)	67
Megacyttarus (Empididae)	50	MICROPEZIDAE	67
Megalopelma (Mycetophilidae)	19	MICROPEZINAE	67
Megamerina (Megamerinidae)	67	Microphor (Dolichopodidae)	57
MEGAMERINIDAE	67	Microphorella (Dolichopodidae)	57
Megaphthalma (Scatophagidae)	89	MICROPHORINAE	57
Megaphthalmoides (Scatophagidae)	90	MICROPHORINI	57
Megaselia (Phoridae)	58	Microphthalma (Tachinidae)	105
Megasyrphus (Syrphidae)	61	MICROPHTHALMINI	105
Meghyperus (Atelestidae)	57	Microprosopa (Scatophagidae)	90
Megophthalmidia (Mycetophilidae)	20	Micropsectra (Chironomidae)	39
Mehria (Sarcophagidae)	99	Micropselapha (Scatophagidae)	89
Meigenia (Tachinidae)	100	Microsania (Platypezidae)	57
Melanagromyza (Agromyzidae)	75	MICROSANIINAE	57
Melangyna (Syrphidae)	61	Microsoma (Tachinidae)	105
Melanogaster (Syrphidae)	64	Microtendipes (Chironomidae)	38
Melanochaeta (Chloropidae)	81	Mikiola (Cecidomyiidae)	27
Melanolimonia (Limoniidae)	14	Mikomya (Cecidomyiidae)	27
Melanomyia (Calliphoridae)	97	MILESIINAE	62
MELANOMYINAE	97	MILESIINI	65
Melanophora (Rhinophoridae)	97	Milichia (Milichiidae)	80
Melanosoma (Conopidae)	68	Milichiella (Milichiidae)	80
Melanostolus (Dolichopodidae)	55	MILICHIIDAE	80
Melanostoma (Syrphidae)	62	MILICHIINAE	80
MELANOSTOMATINI	62	Miltogramma (Sarcophagidae)	98
Melanum (Chloropidae)	82	Miltogrammatidium (Sarcophagidae)	98
Meliera (Otitidae)	70	MILTOGRAMMATINAE	98
Meligramma (Syrphidae)	61	MILTOGRAMMATINI	98
Melinda (Calliphoridae)	97	Minettia (Lauxaniidae)	72
Meliscaeva (Syrphidae)	61	Minilimosina (Sphaeroceridae)	85
Melisonera (Tachinidae)	105	Mintho (Tachinidae)	105
Melochelea (Ceratopogonidae)	41	MINTHOINI	105
Melophagus (Hippoboscidae)	107	Minuscula (Sphaeroceridae)	86
Menoziola (Phoridae)	60	Mmamaevia (Cecidomyiidae)	30
Meoneura (Carnidae)	79	Mochlonyx (Chaoboridae)	34
Merodon (Syrphidae)	64	MOCHLONYXINI	34
Meromyza (Chloropidae)	82	Molobratia (Asilidae)	47
Meroplus (Sepsidae)	74	Molophilus (Limoniidae)	13
Mesaxymyia (Axymyiidae)	17	Monarthropalpus (Cecidomyiidae)	30
Mesembrina (Muscidae)	94	Monoclona (Mycetophilidae)	19
Mesembrius (Syrphidae)	65	Monoculicoides (Ceratopogonidae)	39
Mesocyphona (Limoniidae)	12	Monodiamesa (Chironomidae)	35
Mesomelaena (Sarcophagidae)	98	Monodiplosis (Cecidomyiidae)	30
Mesopsectrocladius (Chironomidae)	37	Monohela (Ceratopogonidae)	40
Metalimnobia (Limoniidae)	15	Monopelopia (Chironomidae)	35
Metasyrphus (Syrphidae)	61	Monopsectrocladius (Chironomidae)	37
Metatrichocera (Trichoceridae)	11	Morellia (Muscidae)	94
Metopia (Sarcophagidae)	98	Morinia (Calliphoridae)	97
METOPINI	98	MORMIINI	31
Metopina (Phoridae)	60	Morpholeria (Heleomyzidae)	82
Metopodia (Sarcophagidae)	98	Moschusa (Sarcophagidae)	98
Metopomyza (Agromyzidae)	77	Mosillus (Ephydriidae)	88
Metopostigma (Chloropidae)	82	Musca (Muscidae)	94
Metriocnemus (Chironomidae)	36	MUSCIDAE	93
Microcephalops (Pipunculidae)	66	Muscina (Muscidae)	93
Microceris (Chloropidae)	81	MUSCINAE	93
Microdon (Syrphidae)	66	MUSCINI	94
MICRODONTINAE	66	Myathropa (Syrphidae)	65
Microdrosophila (Drosophilidae)	87	Mycetaulus (Piophilidae)	69
Microhelea (Ceratopogonidae)	42	Mycetobia (Mycetobiidae)	32
Microchironomus (Chironomidae)	38	MYCETOBIIDAE	32
Microchrysa (Stratiomyidae)	44	MYCETOBIIINAE	32
Microlasioptera (Cecidomyiidae)	25	Mycetophila (Mycetophilidae)	20
Microlimonia (Limoniidae)	14	MYCETOPHILIDAE	19

MYCETOPHILINAE	20	Neottiophilum (Piophilidae)	70
MYCETOPHILINI	20	Neozavrelia (Chironomidae)	39
Mycodiplosis (Cecidomyiidae)	30	NEPHROCERINAE	66
Mycodrosophila (Drosophilidae)	87	Nephrocerus (Pipunculidae)	66
Mycomya (Mycetophilidae)	19	Nephrotoma (Tipulidae)	15
MYCOMYINI	19	Neria (Micropezidae)	67
Mycomyopsis (Mycetophilidae)	19	Neuratelia (Mycetophilidae)	19
Mycophaga (Anthomyiidae)	92	Neurigona (Dolichopodidae)	54
Mycosciara (Sciaridae)	24	NEURIGONINAE	54
Mydaea (Muscidae)	95	Nevermannia (Simuliidae)	42
MYDAEINAE	95	Nigrotipula (Tipulidae)	16
Myennis (Otitidae)	70	Nilea (Tachinidae)	101
Myodris (Perisclididae)	79	Nilotanypus (Chironomidae)	35
Myoleja (Tephritidae)	72	Noeeta (Tephritidae)	71
Myolepta (Syrphidae)	64	Norellia (Scatophagidae)	89
Myopa (Conopidae)	68	NORELLINAE	89
Myopina (Anthomyiidae)	92	Norellisoma (Scatophagidae)	89
MYOPINAE	68	Norrbonia (Sphaeroceridae)	84
MYOPINI	68	Nostima (Ephydriidae)	88
Myopites (Tephritidae)	70	Notiphila (Ephydriidae)	88
MYOPITINAE	70	NOTIPHILINI	88
Myopota (Conopidae)	68	Notolopha (Mycetophilidae)	22
Myospila (Muscidae)	95	Nowickia (Tachinidae)	103
Myrosia (Mycetophilidae)	22	Nudopella (Sphaeroceridae)	86
MYTHICOMYIIDAE	45	Numantia (Limoniidae)	14
Myxexoristops (Tachinidae)	102	Nycteribia (Nycteribiidae)	107
		NYCTERIBIIDAE	107
		Nyctia (Sarcophagidae)	98
Nanna (Scatophagidae)	90		
Nanocladius (Chironomidae)	36		
Napomyza (Agromyzidae)	77	Obuchovia (Simuliidae)	42
Nartshukiella (Chloropidae)	81	Ocydromia (Hybotidae)	52
Nasiternella (Pedicidae)	15	OCYDROMIINAE	51
Natarsia (Chironomidae)	35	OCYDROMIINI	52
NATARSIINI	35	Ocytata (Tachinidae)	102
NEARINI	104	Odinia (Odiniidae)	75
Nematoproctus (Dolichopodidae)	54	ODINIINAE	75
Nemopoda (Sepsidae)	74	Odonatisca (Tipulidae)	16
Nemoraia (Tachinidae)	103	Odontomesa (Chironomidae)	35
NEMORAEINI	103	Odontomyia (Stratiomyidae)	44
Nemorilla (Tachinidae)	101	Oebalia (Sarcophagidae)	98
Nemorimyza (Agromyzidae)	77	OEBALIINI	98
Nemotelus (Stratiomyidae)	45	Oecacta (Ceratopogonidae)	39
Nealticomerus (Odiniidae)	75	Oecothia (Heleomyzidae)	83
Neoscia (Syrphidae)	64	OECOTHEINI	83
Neocnemodon (Syrphidae)	62	Oedalea (Hybotidae)	52
Neoculex (Culicidae)	34	OEDALEINI	52
Neoempheria (Mycetophilidae)	19	Oedesiella (Chloropidae)	80
Neopitriptus (Asilidae)	48	OESTRIDAE	107
Neohaplegis (Chloropidae)	82	OESTRINAE	107
Neoitamus (Asilidae)	48	Oestromyia (Hypodermatidae)	107
Neoleria (Heleomyzidae)	82	OESTROMYIINAE	107
Neoleucopis (Chamaemyiidae)	73	Oestrus (Oestridae)	107
Neolimnomyia (Limoniidae)	11	Ogcodes (Acroceridae)	45
Neolimnophila (Limoniidae)	13	Ochlerotatus (Culicidae)	34
Neolimonia (Limoniidae)	15	Ochthera (Ephydriidae)	88
Neomikiella (Cecidomyiidae)	27	OCHTHERINI	88
Neomochtherus (Asilidae)	48	Oldenbergiella (Heleomyzidae)	83
Neomycomya (Mycetophilidae)	19	Olfersia (Hippoboscidae)	107
Neomyia (Muscidae)	94	OLIGOTROPHINI	25
Neopachygaster (Stratiomyidae)	45	Oligotrophus (Cecidomyiidae)	27
Neophryxe (Tachinidae)	100	Omisis (Chironomidae)	38
Neophyllomyza (Milichiidae)	80	Oncopygius (Dolichopodidae)	54
Neoplatyura (Keroplatidae)	18	Onesia (Calliphoridae)	97
Neoscatella (Ephydriidae)	89	Onychogonia (Tachinidae)	102
Neostempellina (Chironomidae)	39	Oomormia (Psychodidae)	31
NEOTTIOPHILINAE	70	Opacifrons (Sphaeroceridae)	85

<i>Opacifrons</i> (Sphaeroceridae)	85	<i>Pappiella</i> (Sphaeroceridae)	85
<i>Opalimosina</i> (Sphaeroceridae)	85	<i>Parabazarella</i> (Psychodidae)	32
<i>Opesia</i> (Tachinidae)	106	<i>Parabellieria</i> (Sarcophagidae)	99
<i>Opetia</i> (Opetiidae)	57	<i>Paracacoxenus</i> (Drosophilidae)	86
OPETIIDAE	57	<i>Paracladius</i> (Chironomidae)	37
<i>Ophiomyia</i> (Agromyzidae)	76	<i>Paracladopelma</i> (Chironomidae)	38
<i>Oplisa</i> (Rhinophoridae)	97	<i>Paraclusia</i> (Clusiidae)	75
<i>Oplodontha</i> (Stratiomyidae)	44	<i>Paracoenia</i> (Ephydriidae)	89
<i>Opomyza</i> (Opomyzidae)	78	<i>Paracordyla</i> (Mycetophilidae)	22
OPOMYZIDAE	78	<i>Paracraspedothrix</i> (Tachinidae)	101
<i>Orbellia</i> (Heleomyzidae)	83	<i>Paracricotopus</i> (Chironomidae)	37
ORBELLIINI	83	<i>Paradelia</i> (Anthomyiidae)	92
<i>Orellia</i> (Tephritidae)	71	<i>Paradelphomyia</i> (Limoniidae)	11
<i>Oreogeton</i> (Empididae)	48	<i>Paradicranota</i> (Pediciidae)	15
OREOGETONINAE	48	<i>Paradiplosis</i> (Cecidomyiidae)	30
<i>Oreophila</i> (Limoniidae)	13	PARAGINI	62
<i>Orfelia</i> (Keroplatidae)	18	<i>Paragus</i> (Syrphidae)	62
<i>Orchisia</i> (Muscidae)	96	<i>Parachironomus</i> (Chironomidae)	38
<i>Orimarga</i> (Limoniidae)	15	<i>Parajungiella</i> (Psychodidae)	31
<i>Orimargula</i> (Limoniidae)	14	<i>Parakiefferiella</i> (Chironomidae)	37
<i>Ormosia</i> (Limoniidae)	13	<i>Paralimosina</i> (Sphaeroceridae)	85
<i>Ornithoica</i> (Hippoboscidae)	107	<i>Parallelodiplosis</i> (Cecidomyiidae)	30
ORNITHOICINAE	107	<i>Parallelomma</i> (Scatophagidae)	89
<i>Ornithomyia</i> (Hippoboscidae)	107	<i>Paramacronychia</i> (Sarcophagidae)	98
ORNITHOMYIINAE	107	PARAMACRONYCHIINAE	98
<i>Ornithophila</i> (Hippoboscidae)	107	PARAMACRONYCHIINI	98
<i>Oropezella</i> (Hybotidae)	52	<i>Paramerina</i> (Chironomidae)	35
<i>Orphnephilina</i> (Thaumaleidae)	42	<i>Parametricnemus</i> (Chironomidae)	37
<i>Orthacheta</i> (Scatophagidae)	90	<i>Paramormia</i> (Psychodidae)	31
ORTHOCLADIINAE	35	PARAMORMIINI	31
<i>Orthocladius</i> (Chironomidae)	36	<i>Paranthomyza</i> (Anthomyzidae)	79
<i>Orthonevra</i> (Syrphidae)	64	<i>Parapegomyia</i> (Anthomyiidae)	91
<i>Oscinella</i> (Chloropidae)	81	<i>Paraphaenocladius</i> (Chironomidae)	37
OSCINELLINAE	80	<i>Paraphylidorea</i> (Limoniidae)	11
<i>Oscinimorpha</i> (Chloropidae)	81	<i>Paraphytomyza</i> (Agromyzidae)	76
<i>Oscinisoma</i> (Chloropidae)	81	<i>Parapiophila</i> (Piophilidae)	69
<i>Oswaldia</i> (Tachinidae)	101	<i>Paraplatypeza</i> (Platypezidae)	58
<i>Otites</i> (Otitidae)	70	<i>Parapsectra</i> (Chironomidae)	39
OTITIDAE	70	<i>Pararhamphomyia</i> (Empididae)	50
<i>Oxycera</i> (Stratiomyidae)	45	<i>Parasarcophaga</i> (Sarcophagidae)	99
<i>Oxyna</i> (Tephritidae)	71	<i>Parascaptomyza</i> (Drosophilidae)	87
<i>Oxypsila</i> (Psilidae)	68	<i>Parasetigena</i> (Tachinidae)	100
<i>Oxyrhiza</i> (Limoniidae)	11	<i>Parasyrphus</i> (Syrphidae)	61
<i>Ozirhineus</i> (Cecidomyiidae)	25	<i>Paratanytarsus</i> (Chironomidae)	39
		<i>Paratendipes</i> (Chironomidae)	38
<i>Pachycerina</i> (Lauxaniidae)	72	PARATHALASSIINI	57
<i>Pachygaster</i> (Stratiomyidae)	45	<i>Paratinia</i> (Mycetophilidae)	19
PACHYGASTERINAE	45	<i>Paratrachocladius</i> (Chironomidae)	37
<i>Pachymeria</i> (Empididae)	50	<i>Paratrisocladius</i> (Chironomidae)	37
<i>Pachypericoma</i> (Psychodidae)	32	<i>Paratrixa</i> (Tachinidae)	101
<i>Pachysphlyria</i> (Syrphidae)	62	<i>Paratryphera</i> (Tachinidae)	101
<i>Palaeodocosia</i> (Mycetophilidae)	20	<i>Parectecephala</i> (Chloropidae)	82
<i>Palaeogonomyia</i> (Limoniidae)	13	<i>Paregle</i> (Anthomyiidae)	92
<i>Pales</i> (Tachinidae)	102	<i>Parepidosis</i> (Cecidomyiidae)	25
<i>Palloptera</i> (Pallopteridae)	69	<i>Parhelophilus</i> (Syrphidae)	65
PALLOPTERIDAE	69	<i>Parilisia</i> (Limoniidae)	12
PALLOPTERINAE	69	<i>Paritamus</i> (Asilidae)	48
<i>Palpomyia</i> (Ceratopogonidae)	41	<i>Paroecus</i> (Lauxaniidae)	72
PALPOMYIINI	40	<i>Paroedesiella</i> (Chloropidae)	80
<i>Pamponerus</i> (Asilidae)	48	<i>Parochthiphila</i> (Chamaemyiidae)	73
<i>Pandasyopthalmus</i> (Syrphidae)	62	<i>Parorthocladius</i> (Chironomidae)	37
<i>Pandelleana</i> (Sarcophagidae)	99	<i>Parydra</i> (Ephydriidae)	89
<i>Pandelleisca</i> (Sarcophagidae)	99	PARYDRINI	89
<i>Pandelleola</i> (Sarcophagidae)	99	<i>Paykullia</i> (Rhinophoridae)	97
<i>Pandivritlia</i> (Therevidae)	46	<i>Pedicia</i> (Pediciidae)	15
<i>Parumerus</i> (Psychodidae)	31	PEDICIIDAE	15

Pegomya (Anthomyiidae)	92	Phortica (Drosophilidae)	86
Pegoplata (Anthomyiidae)	92	Phronia (Mycetophilidae)	21
Pelatachina (Tachinidae)	104	Phrosia (Scatophagidae)	89
PELATACHININI	104	Phrosinella (Sarcophagidae)	98
Pelecocera (Syrphidae)	63	Phryno (Tachinidae)	103
Peleteria (Tachinidae)	103	Phryxe (Tachinidae)	102
Pelidnoptera (Phaeomyiidae)	73	Phthiria (Mycetophilidae)	19
Pelina (Ephydriidae)	88	Phthiria (Phthiriidae)	45
Pelomyia (Tethinidae)	79	Phthiridium (Nycteribiidae)	107
Pelomyiella (Tethinidae)	79	PHTHIRIIDAE	45
PELOROPEODINAE	54	Phthitia (Sphaeroceridae)	85
Pemphigocercis (Cecidomyiidae)	27	Phylidorea (Limoniidae)	11
Penicillidia (Nycteribiidae)	107	Phyllodromia (Empididae)	51
Pentaneurella (Chironomidae)	35	Phyllolabis (Limoniidae)	13
PENTANEURINI	35	Phyllomya (Tachinidae)	105
Pentapedilum (Chironomidae)	38	Phyllomyza (Milichiidae)	80
Penthetria (Pleciidae)	17	Phylloteles (Sarcophagidae)	98
Peodes (Dolichopodidae)	54	PHYLOTETELINI	98
Periarchielops (Tachinidae)	102	Physemocercis (Cecidomyiidae)	27
Peribaea (Tachinidae)	104	Physiphora (Ulidiidae)	70
Pericoma (Psychodidae)	32	Physocephala (Conopidae)	68
PERICOMINI	32	Phytagromyza (Agromyzidae)	77
Peripsychosa (Psychodidae)	31	Phyto (Rhinophoridae)	97
PERISCCELIDIDAE	79	Phytobia (Agromyzidae)	77
Periscelis (Periscelididae)	79	Phytoliriomyza (Agromyzidae)	77
Periscepsia (Tachinidae)	105	Phytomyptera (Tachinidae)	104
Petagnia (Tachinidae)	103	Phytomyza (Agromyzidae)	77
Petenia (Tachinidae)	105	PHYTOMYZINAE	76
Peusomyia (Chaoboridae)	34	Phytosciara (Sciaridae)	24
Pexopsis (Tachinidae)	103	Picconia (Tachinidae)	101
Peyerimhoffia (Sciaridae)	24	Pierretia (Sarcophagidae)	99
Phaenobezzia (Ceratopogonidae)	41	Piezura (Fanniidae)	93
Phaeobalia (Empididae)	51	Pilaria (Limoniidae)	11
PHAEOMYIIDAE	73	Piophila (Piophilidae)	69
Phalacroceras (Cylindrotomidae)	15	PIOPHILIDAE	69
Phalacrotophora (Phoridae)	60	PIOPHILINAE	69
Phania (Tachinidae)	106	Pipiza (Syrphidae)	62
Phaonia (Muscidae)	94	Pipizella (Syrphidae)	62
PHAONIINAE	94	PIPIZINI	62
PHAONIINI	94	PIPUNCULIDAE	66
Pharsoreichertella (Scatopsidae)	33	Pipunculina (Pipunculidae)	67
Pharyngomyia (Oestridae)	107	PIPUNCULINAE	66
Phasia (Tachinidae)	106	Pipunculus (Pipunculidae)	66
PHASIINAE	106	Placochela (Cecidomyiidae)	28
PHASIINI	106	Planetella (Cecidomyiidae)	30
Phebellia (Tachinidae)	102	Plastosciara (Sciaridae)	24
Phegomyia (Cecidomyiidae)	27	Platurocypta (Mycetophilidae)	21
Phenicellia (Tachinidae)	105	Platycephala (Chloropidae)	82
Pherbellia (Sciomyzidae)	73	Platycheirus (Syrphidae)	62
Pherbina (Sciomyzidae)	74	Platymya (Tachinidae)	103
Philipomyia (Tabanidae)	44	Platypalpus (Hybotidae)	52
Philocoprella (Sphaeroceridae)	85	Platyparea (Tephritidae)	72
Philolutra (Empididae)	51	Platypeza (Platypezidae)	58
Philonicus (Asilidae)	48	PLATYPEZIDAE	57
Philophylla (Tephritidae)	72	Platypozina (Platypezidae)	58
Philosepedon (Psychodidae)	31	PLATYPEZINAE	58
Philotelma (Ephydriidae)	89	Platyptera (Empididae)	50
Philygia (Ephydriidae)	88	Platystoma (Platystomatidae)	70
PHILYGRINI	88	PLATYSTOMATIDAE	70
Phonomyia (Tachinidae)	102	Platystyla (Psilidae)	68
Phora (Phoridae)	60	Platytipula (Tipulidae)	16
Phorbia (Anthomyiidae)	92	Platyura (Keroplatidae)	18
PHORIDAE	58	PLECIIDAE	17
Phorinia (Tachinidae)	100	Plectanocnema (Phoridae)	60
Phormia (Calliphoridae)	97	Plemeliella (Cecidomyiidae)	30
Phorocera (Tachinidae)	100	Pliorecepta (Tephritidae)	72

Pneumia (Psychodidae)	32	Pseudogaurax (Chloropidae)	81
Pocota (Syrphidae)	65	Pseudogonia (Tachinidae)	103
Poecilobothrus (Dolichopodidae)	56	PSEUDOCHIRONOMINI	39
Poemyza (Agromyzidae)	77	Pseudochironomus (Chironomidae)	39
Pogonosoma (Asilidae)	47	Pseudokiefferiella (Chironomidae)	35
Pogonota (Scatophagidae)	90	Pseudolimnophila (Limoniidae)	12
Polietes (Muscidae)	94	Pseudolynchia (Hippoboscidae)	107
Policheta (Tachinidae)	101	Pseudonapomyza (Agromyzidae)	78
Pollenia (Calliphoridae)	97	Pseudopachychaeta (Chloropidae)	82
POLLENIINAE	97	Pseudopachystylum (Tachinidae)	104
Polyblepharis (Empididae)	50	Pseudoperichaeta (Tachinidae)	102
Polylepta (Mycetophilidae)	19	Pseudopomyza (Pseudopomyzidae)	67
Polyodaspis (Chloropidae)	81	PSEUDOPOMYZIDAE	67
Polypedilum (Chironomidae)	38	Pseudorthocladus (Chironomidae)	37
Polyporivora (Platypezidae)	58	Pseudorymosia (Mycetophilidae)	23
Polystepha (Cecidomyiidae)	28	Pseudosmittia (Chironomidae)	37
Polytrichophora (Ephydriidae)	88	Pseudowiedemannia (Empididae)	51
Pontoculicoides (Ceratopogonidae)	40	Psila (Psilidae)	68
Porricondyla (Cecidomyiidae)	25	PSILIDAE	67
PORRICONDYLINAE	25	Psiloconopa (Limoniidae)	13
Portevinia (Syrphidae)	63	Psilokempia (Ceratopogonidae)	41
Potamia (Muscidae)	94	Psilopa (Ephydriidae)	87
Potthastia (Chironomidae)	35	PSILOPINI	87
Ppeplomyza (Lauaxiidae)	72	Psilosoma (Psilidae)	68
Prionocera (Tipulidae)	15	Psilota (Syrphidae)	64
PRIONOCERINI	15	Psilotanytus (Chironomidae)	35
Prionolabis (Limoniidae)	12	Psycmera (Psychodidae)	31
Probezzia (Ceratopogonidae)	40	Psycha (Psychodidae)	31
PROCLADIINI	34	Psychoda (Psychodidae)	31
Procladius (Chironomidae)	34	PSYCHODIDAE	30
Prodiamesa (Chironomidae)	35	PSYCHODINAE	31
PRODIAMESINAE	35	PSYCHODINI	31
Prochyliza (Piophilidae)	70	Psychodocha (Psychodidae)	32
Prokempia (Ceratopogonidae)	41	Psychodula (Psychodidae)	32
Prolipophleps (Limoniidae)	12	Psychogella (Psychodidae)	31
Promormia (Psychodidae)	31	Psychocha (Psychodidae)	31
Propillocerus (Chironomidae)	37	Psychomora (Psychodidae)	32
Prosciara (Sciaridae)	24	Psychomormia (Psychodidae)	31
Prosenia (Tachinidae)	105	Ptenotaenia (Anthomyzidae)	79
Prosethilla (Tachinidae)	101	Pterelachisus (Tipulidae)	16
PROSIMULIINI	42	Pterella (Sarcophagidae)	98
Prosimulium (Simuliidae)	42	Pteremis (Sphaeroceridae)	85
Prosopea (Tachinidae)	103	Pteromicra (Sciomyzidae)	74
Protearomyia (Lonchaeidae)	69	Ptesiomyia (Tachinidae)	102
Protocalliphora (Calliphoridae)	97	Ptilomyia (Ephydriidae)	87
Protoclythia (Platypezidae)	58	Ptiolina (Rhagionidae)	43
PROTODEXIINI	98	Ptychoneura (Sarcophagidae)	98
Protogonomyia (Limoniidae)	12	Ptychoptera (Ptychopteridae)	33
Protomiltogramma (Sarcophagidae)	98	PTYCHOPTERIDAE	33
Protophormia (Calliphoridae)	97	Pullimosina (Sphaeroceridae)	85
Protophophila (Piophilidae)	70	Puncticorpus (Sphaeroceridae)	85
Psacadina (Sciomyzidae)	74	Putoniella (Cecidomyiidae)	30
Psammopogon (Ceratopogonidae)	41	Pygobezzia (Ceratopogonidae)	41
PSARINI	64	Pyratula (Keroplastidae)	18
Psarus (Syrphidae)	64	Pyrellia (Muscidae)	94
Psectrocladius (Chironomidae)	37		
PSECTROSCIARINAE	32	Rabdophaga (Cecidomyiidae)	27
Psectrotanytus (Chironomidae)	35	Rachispoda (Sphaeroceridae)	84
Pseudacteon (Phoridae)	60	Rainieria (Micropezidae)	67
Pseudexechia (Mycetophilidae)	23	Ramonda (Tachinidae)	105
Pseudobrachypeza (Mycetophilidae)	23	Ravinia (Sarcophagidae)	99
Pseudocoenosia (Muscidae)	96	RAVINIINI	99
Pseudocollinella (Sphaeroceridae)	85	Redtenbacheria (Tachinidae)	106
Pseudoculicoides (Ceratopogonidae)	41	Reichertella (Scatopsidae)	33
Pseudodiamesa (Chironomidae)	35	REINWARDTIINI	93
Pseudoficalbia (Culicidae)	34	Renocera (Sciomyzidae)	74


Resseliella (Cecidomyiidae)	30	Scaptomyza (Drosophilidae)	87
Rhabdomastix (Limoniidae)	13	Scatella (Ephydriidae)	89
Rhacochlaena (Tephritidae)	71	SCATELLINI	89
Rhadiurgus (Asilidae)	48	Scatophaga (Scatophagidae)	90
Rhagio (Rhagionidae)	43	SCATOPHAGIDAE	89
RHAGIONIDAE	43	SCATOPHAGINAE	90
RHAGIONINAE	43	Scatophila (Ephydriidae)	89
Rhagoletis (Tephritidae)	72	Scatopsiara (Sciaridae)	24
Rhamphomyia (Empididae)	50	Scatopse (Scatopsidae)	33
Rhaphidolabis (Pediidae)	15	SCATOPSIDAE	32
RHAPHIINAE	54	SCATOPSINAE	33
Rhaphiochaeta (Tachinidae)	101	SCATOPSINI	33
Rhaphiopyga (Chloropidae)	80	Scellus (Dolichopodidae)	54
Rhaphium (Dolichopodidae)	54	SCENOPINIDAE	46
Rheocricotopus (Chironomidae)	37	Scenopinus (Scenopinidae)	46
Rheopelopia (Chironomidae)	35	Sceptonia (Mycetophilidae)	21
Rheosmittia (Chironomidae)	37	SCIAPODINAE	54
Rheotanytarsus (Chironomidae)	39	Sciapus (Dolichopodidae)	54
Rhexosa (Scatopsidae)	33	Sciara (Sciaridae)	24
Rhicnoessa (Tethinidae)	79	SCIARIDAE	23
Rhingia (Syrphidae)	63	Sciomyza (Sciomyzidae)	74
RHINIINAE	97	SCIOMYZIDAE	73
Rhinoestrus (Oestridae)	107	SCIOMYZINAE	73
Rhinomorinia (Rhinophoridae)	97	SCIOMYZINI	73
Rhinophora (Rhinophoridae)	97	Sciophila (Mycetophilidae)	19
RHINOPHORIDAE	97	SCIOPHILINAE	19
Rhiphidia (Limoniidae)	15	SCIOPHILINI	19
Rhizomyia (Cecidomyiidae)	28	Sclerophallus (Chloropidae)	82
Rhodesiella (Chloropidae)	81	Scleroprocta (Limoniidae)	13
Rhopalomyia (Cecidomyiidae)	28	Scoliaphleps (Scatophagidae)	89
Rhopalopterum (Chloropidae)	81	Scoliocentra (Heleomyzidae)	82
Rhypholophus (Limoniidae)	13	Sebessia (Ceratopogonidae)	41
Rivellia (Platystomatidae)	70	Seioptera (Otitidae)	70
Robineauella (Sarcophagidae)	100	Selachops (Agromyzidae)	78
Rocetelion (Keroplastidae)	18	Semudobia (Cecidomyiidae)	28
Rondania (Tachinidae)	105	Senometopia (Tachinidae)	102
Rondaniella (Mycetophilidae)	20	Senotainia (Sarcophagidae)	98
Rondaniola (Cecidomyiidae)	28	Sepedon (Sciomyzidae)	74
Rosellea (Sarcophagidae)	100	SEPSIDAE	74
Rudolfia (Sphaeroceridae)	85	Sepsis (Sepsidae)	74
Rudolfina (Sphaeroceridae)	85	Sergentia (Chironomidae)	38
Rutylapa (Keroplastidae)	18	Seri (Platypezidae)	58
Rymosia (Mycetophilidae)	23	Sericomyia (Syrphidae)	64
		SERICOMYIINI	64
Sacandaga (Limoniidae)	13	Serromyia (Ceratopogonidae)	40
Sackenomyia (Cecidomyiidae)	28	Servaisia (Sarcophagidae)	99
Saetheria (Chironomidae)	38	Setacera (Ephydriidae)	89
Saigusaia (Mycetophilidae)	20	Setisquamalonchaea (Lonchaeidae)	69
Saltella (Sepsidae)	74	Schineria (Tachinidae)	103
Salticella (Sciomyzidae)	73	Schizohelca (Ceratopogonidae)	40
SALTICELLINAE	73	Schizomyia (Cecidomyiidae)	28
Sapromyza (Lauxaniidae)	72	Schmidtella (Cecidomyiidae)	28
Sapromyzosoma (Lauxaniidae)	73	Schoenbaueria (Simuliidae)	42
Saraella (Psychodidae)	32	Schoenomyza (Muscidae)	96
Sarcophaga (Sarcophagidae)	100	Schoenophilus (Dolichopodidae)	54
SARCOPHAGIDAE	98	Schroederella (Heleomyzidae)	83
SARCOPHAGINAE	98	Schueziella (Cecidomyiidae)	28
SARCOPHAGINI	99	Schumannimyia (Lauxaniidae)	73
Sarcophila (Sarcophagidae)	98	Schummelia (Tipulidae)	17
Sarcotachinella (Sarcophagidae)	99	Schwenckfeldina (Sciaridae)	24
SARGINAE	44	SICINI	68
Sargus (Stratiomyidae)	44	Sicus (Conopidae)	68
Satchelliella (Psychodidae)	32	SIGALOESSINAE	79
Savtshenkia (Tipulidae)	16	Silvius (Tabanidae)	43
Scaeva (Syrphidae)	61	SIMULIIDAE	42
Scaptodrosophila (Drosophilidae)	87	SIMULIINAE	42

SIMULIINI	42	Stigmatomeria (Mycetophilidae)	22
Simulium (Simuliidae)	42	Stichilus (Phoridae)	60
Siphona (Tachinidae)	104	Stichopogon (Asilidae)	47
Siphonella (Chloropidae)	81	STICHOPOGONINI	47
SIPHONELLOPSINAE	80	Stilobezzia (Ceratopogonidae)	40
SIPHONINI	104	STILOBEZZIINI	40
Siphunculina (Chloropidae)	81	Stilocladius (Chironomidae)	37
Sitodiplosis (Cecidomyiidae)	30	Stilpnogaster (Asilidae)	48
Sivalimnobia (Limoniidae)	14	Stilpon (Hybotidae)	53
Smittia (Tachinidae)	101	Stiphrosoma (Anthomyzidae)	79
Smittia (Chironomidae)	37	Stomatosema (Cecidomyiidae)	28
Solieria (Tachinidae)	104	STOMATOSEMIDI	28
Solva (Xylomyidae)	44	Stomina (Tachinidae)	105
Sophophora (Drosophilidae)	86	Stomorhina (Calliphoridae)	97
Soudekia (Sciaridae)	24	STOMOXYINI	94
Spallanzania (Tachinidae)	103	Stomoxys (Muscidae)	94
Spania (Rhagionidae)	43	STRATIOMYIDAE	44
SPANIINAE	43	STRATIOMYINAE	44
Spanochaeta (Muscidae)	96	Stratiomys (Stratiomyidae)	44
Spanoparea (Heleomyzidae)	82	Strobilomyia (Anthomyiidae)	92
Spathephilus (Scatophagidae)	90	Strongygaster (Tachinidae)	106
Spathobdella (Sciaridae)	24	STRONGYGASTERINI	106
Spazigaster (Syrphidae)	62	Strongylophthalmyia (Strongylophthalmyiidae)	67
Spaziphora (Scatophagidae)	90	STRONGYLOPHTHALMYIIDAE	67
Speccafrons (Chloropidae)	81	Sturmia (Tachinidae)	103
Spelobia (Sphaeroceridae)	85	Subclytia (Tachinidae)	106
Speolepta (Mycetophilidae)	20	Subhylemyia (Anthomyiidae)	92
Sphaeconophilus (Limoniidae)	12	Suillia (Heleomyzidae)	83
Sphaerocera (Sphaeroceridae)	84	SUILLIINAE	83
SPHAEROCERIDAE	84	Svarciella (Sphaeroceridae)	85
SPHAEROCERINAE	84	Swammerdamella (Scatopsidae)	33
Sphaeromias (Ceratopogonidae)	40	SWAMMERDAMELLINI	33
SPHAEROMIINI	40	SYCORACINAE	30
Sphaerophoria (Syrphidae)	61	Sycorax (Psychodidae)	30
Sphegina (Syrphidae)	64	Sylvicola (Anisopodidae)	32
Sphenella (Tephritidae)	71	Symbalophthalmus (Hybotidae)	53
Sphenometopa (Sarcophagidae)	98	Symbiolcladius (Chironomidae)	37
Sphixapata (Sarcophagidae)	98	Symmerus (Ditomyiidae)	18
Sphiximorpha (Syrphidae)	65	Symphoromyia (Rhagionidae)	43
Spilogona (Muscidae)	95	Symplecta (Limoniidae)	13
Spilomyia (Syrphidae)	65	Symposiocladius (Chironomidae)	36
Spinilimosina (Sphaeroceridae)	85	Sympotthastia (Chironomidae)	35
Spiniphora (Phoridae)	60	SYMPYCNINAE	54
Spinodrosophila (Drosophilidae)	87	SYMPYCNINI	54
Spogostylum (Bombyliidae)	45	Sympycnus (Dolichopodidae)	54
Spurgia (Cecidomyiidae)	28	Synactia (Tachinidae)	104
Stackelbergcola (Sarcophagidae)	100	Synapha (Mycetophilidae)	20
Staurochaeta (Tachinidae)	101	Syndiamesa (Chironomidae)	35
Stearibia (Piophilidae)	70	Syndyas (Hybotidae)	52
Stefaniella (Cecidomyiidae)	25	Syneches (Hybotidae)	52
Stegana (Drosophilidae)	86	SYNNEURIDAE	33
Stegamina (Drosophilidae)	86	Synneuron (Synneuridae)	33
STEGANINAE	86	Synorthocladius (Chironomidae)	37
STEGANINI	86	Synplasta (Mycetophilidae)	23
Stemonocera (Tephritidae)	72	Syntemna (Mycetophilidae)	19
Stempellina (Chironomidae)	39	Synthyridomyia (Ceratopogonidae)	42
Stempellinella (Chironomidae)	39	Syntormon (Dolichopodidae)	55
Stenopteryx (Hippoboscidae)	107	Syritta (Syrphidae)	65
Stenochironomus (Chironomidae)	38	SYRPHIDAE	60
Stenomicra (Stenomicroidae)	79	SYRPHINAE	60
STENOMICRIDAE	79	SYRPHINI	60
Stenopogon (Asilidae)	47	Syrphus (Syrphidae)	61
STENOPOGONINAE	47	Systata (Otitidae)	70
STENOPOGONINI	47	SYSTEMINI	57
Stevenia (Rhizophoridae)	97	Systemus (Dolichopodidae)	57
Stictochironomus (Chironomidae)	38		

Systoechus (Bombyliidae)	46	Thelymorpha (Tachinidae)	103
Szaboicella (Psychodidae)	32	Thelymyia (Tachinidae)	102
TABANIDAE	43	Themira (Sepsidae)	74
TABANINAE	43	Thereva (Therevidae)	46
TABANINI	43	THEREVIDAE	46
Tabanus (Tabanidae)	44	Theriopectes (Tabanidae)	44
TAENIAPTERINAE	67	Therodiplosis (Cecidomyiidae)	30
Tachina (Tachinidae)	103	Thienemaniella (Chironomidae)	37
TACHINIDAE	100	Thienemannia (Chironomidae)	37
TACHININAE	103	Thienemannimyia (Chironomidae)	35
TACHININI	103	Thienemanniola (Chironomidae)	39
Tachydromia (Hybotidae)	53	THINOPHILINI	54
TACHYDROMIINAE	52	Thinophilus (Dolichopodidae)	54
TACHYDROMIINI	52	Thomsonea (Sarcophagidae)	99
Tachypeza (Hybotidae)	53	Thoracochaeta (Sphaeroceridae)	85
TACHYTRECHINI	55	Threticus (Psychodidae)	31
Tachytrechus (Dolichopodidae)	55	Thricops (Muscidae)	93
Tanypeza (Tanypezidae)	67	Thripomorpha (Scatopsidae)	33
TANYPEZIDAE	67	THRIPOMORPHINI	33
TANYPODINAE	34	THRYPTICINI	57
TANYPODINI	34	Thrypticus (Dolichopodidae)	57
Tanyptera (Tipulidae)	17	Thurauia (Cecidomyiidae)	30
TANYPTERINI	17	THYREOPHORINAE	70
Tanypus (Chironomidae)	34	Thyridanthrax (Bombyliidae)	46
TANYTARSINI	39	THYRIDANTHRAXINI	46
Tanytarsus (Chironomidae)	39	Thyridomyia (Ceratopogonidae)	42
Taramormia (Psychodidae)	31	Thyrsocnema (Sarcophagidae)	100
Tarnania (Mycetophilidae)	23	Timavia (Tachinidae)	101
Tasiocera (Limoniidae)	13	Tinearia (Psychodidae)	32
Taxigramma (Sarcophagidae)	98	Tipula (Tipulidae)	16
Taxomyia (Cecidomyiidae)	28	TIPULIDAE	15
Telmatopelopia (Chironomidae)	35	TIPULINAE	15
Telmatoscopus (Psychodidae)	31	TIPULINI	15
Telmaturgus (Dolichopodidae)	55	Tlephusa (Tachinidae)	102
Telomerina (Sphaeroceridae)	85	Tokungaia (Chironomidae)	37
Telopelopia (Chironomidae)	35	Tolmerus (Asilidae)	48
Temnosira (Pallopteridae)	69	Tomosvaryella (Pipunculidae)	67
Temnostoma (Syrphidae)	65	Tonnoiriella (Psychodidae)	32
TEPHRITIDAE	70	Townsendiellomyia (Tachinidae)	102
TEPHRITINAE	71	Toxoneura (Pallopteridae)	69
TEPHRITINI	71	Trafoia (Tachinidae)	105
Tephritis (Tephritidae)	71	Trachyopella (Sphaeroceridae)	86
Tephrochlaena (Heleomyzidae)	83	Trachysiphonella (Chloropidae)	80
Tephrochlamys (Heleomyzidae)	83	Triarthria (Tachinidae)	104
Tephromyia (Sarcophagidae)	99	TRIARTHRIINI	104
Terellia (Tephritidae)	71	Tribelos (Chironomidae)	38
TERELLIINAE	71	Tricimba (Chloropidae)	81
Terrilimosina (Sphaeroceridae)	85	Tricogena (Rhizophoridae)	97
Tetanocera (Sciomyzidae)	74	Tricyphona (Pediidae)	15
TETANOCERINI	74	Triglyphus (Syrphidae)	62
Tetanura (Sciomyzidae)	74	Trigonometopus (Lauxaniidae)	73
Tethina (Tethinidae)	79	Trichactia (Tachinidae)	104
TETHINIDAE	79	Trichieurina (Chloropidae)	82
Tetragoneura (Mycetophilidae)	20	Trichina (Hybotidae)	52
Tetrapsila (Psilidae)	67	TRICHININI	51
Teuchogonomyia (Limoniidae)	12	Trichinomyia (Hybotidae)	52
Teuchophorus (Dolichopodidae)	55	Trichocera (Trichoceridae)	11
Thaumalea (Thaumaleidae)	42	TRICHOCERIDAE	11
THAUMALEIDAE	42	Trichohelca (Ceratopogonidae)	42
Thaumastoptera (Limoniidae)	15	Tricholaba (Cecidomyiidae)	30
Thaumatomyia (Chloropidae)	82	Tricholauxania (Lauxaniidae)	73
Thecocarcelia (Tachinidae)	102	Trichomyia (Psychodidae)	31
Thecodiplosis (Cecidomyiidae)	30	TRICHOMYIINAE	31
Thecophora (Conopidae)	68	Trichonta (Mycetophilidae)	21
Thelaira (Tachinidae)	105	Trichopalpus (Scatophagidae)	90
		Trichopeza (Empididae)	51

TRICHOPEZINAE	51	Wiedemannia (Empididae)	51
Trichopsomyia (Syrphidae)	62	Wilhelmia (Simuliidae)	42
Trichopsychoda (Psychodidae)	31	Winnertzia (Cecidomyiidae)	25
<i>Trichoptera</i> (Diastatidae)	87	Winthemia (Tachinidae)	101
Trichosia (Sciaridae)	24	WINTHEMIINI	101
<i>Trilobomyza</i> (Agromyzidae)	76	Wohlfahrtia (Sarcophagidae)	98
<i>Trilophyrobata</i> (Micropezidae)	67		
Trimerina (Ephydriidae)	87	Xanthandrus (Syrphidae)	62
Trimicra (Limoniidae)	13	Xanthempis (Empididae)	50
Triogma (Cylindrotomidae)	15	Xanthogramma (Syrphidae)	61
Triphleba (Phoridae)	60	XANTHOCHLORINAE	54
Tripodura (Chironomidae)	38	Xanthochlorus (Dolichopodidae)	54
Trisopsis (Cecidomyiidae)	30	Xenexechia (Mycetophilidae)	23
Trissopelopia (Chironomidae)	35	Xenodiplosis (Cecidomyiidae)	30
Trixa (Tachinidae)	105	Xenochironomus (Chironomidae)	38
TRIXOSCELIDIDAE	83	Xenopelopia (Chironomidae)	35
Trixoscelis (Trixoscelididae)	83	Xenophyomyza (Agromyzidae)	77
Tropidia (Syrphidae)	65	Xenosepsis (Sepsidae)	75
Trotteria (Cecidomyiidae)	28	Xiphuromorpha (Tipulidae)	17
Trupanea (Tephritidae)	71	Xylodiplosis (Cecidomyiidae)	30
Trypeta (Tephritidae)	72	Xylomya (Xylomyidae)	44
TRYPETINAE	71	XYLOMYIDAE	44
TRYPETINI	71	XYLOPHAGIDAE	43
Trypetoptera (Sciomyzidae)	74	Xylophagus (Xylophagidae)	43
Trypocalliphora (Calliphoridae)	97	Xylopriona (Cecidomyiidae)	25
Tvetenia (Chironomidae)	37	Xylosciara (Sciaridae)	24
Twinnia (Simuliidae)	42	Xylota (Syrphidae)	65
Typhamyza (Anthomyzidae)	79	Xylotachina (Tachinidae)	102
		XYLOTINI	65
Ula (Pedicidae)	15	Xyphosia (Tephritidae)	71
Ulidia (Ulidiidae)	70	Xysta (Tachinidae)	106
ULIDIIDAE	70		
Ulomyia (Psychodidae)	32	Yamatotipula (Tipulidae)	17
Uranotaenia (Culicidae)	34	Ypsydocha (Psychodidae)	32
Urophora (Tephritidae)	70		
Urytalpa (Keroplastidae)	18	Zabrachia (Stratiomyidae)	45
		Zaira (Tachinidae)	101
Varrosellea (Sarcophagidae)	99	Zalutschia (Chironomidae)	37
Verrallia (Pipunculidae)	66	Zaphne (Anthomyiidae)	92
Vestiplex (Tipulidae)	17	Zavrelia (Chironomidae)	39
Vibrissina (Tachinidae)	101	Zavreliella (Chironomidae)	38
Villa (Bombyliidae)	45	Zavreliomyia (Chironomidae)	35
VILLINI	45	Zenillia (Tachinidae)	103
Vimmeria (Sciaridae)	24	Zeuxia (Tachinidae)	105
Volucella (Syrphidae)	63	Zeuxidiplosis (Cecidomyiidae)	30
VOLUCCELLINI	63	Zodion (Conopidae)	68
Voria (Tachinidae)	105	ZODIONINI	68
VORIINI	105	Zophomyia (Tachinidae)	104
		Zygiobia (Cecidomyiidae)	28
Wagneria (Tachinidae)	105	Zygomomyia (Mycetophilidae)	22
Wachtliella (Cecidomyiidae)	28	Zygoneura (Sciaridae)	24
Weberia (Tachinidae)	106		

Check List of Diptera (Insecta) of the Czech and Slovak Republics



Prof. RNDr. Milan Chvála, DrSc.
editor

Published by Karolínium – Charles University Press,
Ovocný trh 3, 116 36 Praha 1
Prague 1977

Vice-Rector-Editor Prof. MUDr. Pavel Klener, DrSc.
Cover Jiří Staněk

Cover Illustration Jindřich Roháček

Layout Jiří Staněk and Doc. RNDr. Jaromír Vaňhara, CSc.
Type matter ÚMV, Nerudova 3, 118 50 Praha 1

Printed by Calamarus Ltd., Prague 3, Roháčova 101

First Edition

ISBN 80-7184-476-4